

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 March 2002 (21.03.2002)

PCT

(10) International Publication Number
WO 02/22598 A1

(51) International Patent Classification⁷: **C07D 401/04**,
405/04, 409/04, A61K 31/4704

Center Street, Castro Valley, CA 94546 (US). **HARRIS, Alex** [US/US]; 14 MacPherson Place, Danville, CA 94526 (US).

(21) International Application Number: PCT/US01/42131

(74) Agent: **FRIEDRICHSEN, Bernard**; Foley & Lardner, 150 East Gilman Street, P.O. Box 1497, Madison, WI 53701-1497 (US).

(22) International Filing Date:
11 September 2001 (11.09.2001)

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(26) Publication Language: English

(30) Priority Data:
60/232,159 11 September 2000 (11.09.2000) US

(71) Applicant (*for all designated States except US*): **CHIRON CORPORATION** [US/US]; 4560 Horton Street, Emeryville, CA 94608-2916 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **RENHOWE, Paul** [US/US]; 262 Stetson Drive, Danville, CA 94506 (US). **PECCHI, Sabina** [IT/US]; 5834 Merriewood Drive, Oakland, CA 94611 (US). **MACHAJEWSKI, Tim** [US/US]; 2514 Norwalk Court, Martinez, CA 94553 (US). **SHAFFER, Cynthia** [US/US]; 4868 El Grande Place, El Sobrante, CA 94803 (US). **TAYLOR, Clarke** [US/US]; 612 Adams Street, Albany, CA 94706 (US). **McCREA, Bill** [US/US]; 1040 Amito Drive, Berkeley, CA 94705 (US). **McBRIDE, Chris** [US/US]; 4504 Lawton Avenue, Oakland, CA 94609 (US). **JAZAN, Elisa** [US/US]; 520 McLaughlin Avenue, Richmond, CA 94805 (US). **WERNETTE-HAMMOND, Mary-Ellen** [US/US]; 17893

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: QUINOLINONE DERIVATIVES

(57) Abstract: Organic compounds having the formulas I and II are provided where the variables have the values described herein. Pharmaceutical formulations include the organic compounds or pharmaceutically acceptable salts thereof and a pharmaceutically acceptable carrier and may be prepared by mixing the organic compounds or pharmaceutically acceptable salts of the organic compounds with a carrier and water. A method of treating a patient includes administering a pharmaceutical formulation according to the invention to a patient in need thereof.

WO 02/22598 A1

QUINOLINONE DERIVATIVES AS TYROSINE KINASE INHIBITORS

FIELD OF THE INVENTION

This invention pertains generally to treating diseases characterized by angiogenesis including cancer. More specifically, the invention described herein pertains to treating diseases characterized by activity of vascular endothelial growth factor receptor tyrosine kinases. The present invention provides, small molecule inhibitors of vascular endothelial growth factor receptor tyrosine kinase, pharmaceutical formulations containing such inhibitors, methods of treating patients with such pharmaceutical formulations, and to methods of preparing such pharmaceutical formulations and inhibitors.

BACKGROUND OF THE INVENTION

Capillaries reach into almost all tissues of the human body and supply tissues with oxygen and nutrients as well as removing waste products. Under typical conditions, the endothelial cells lining the capillaries do not divide, and capillaries, therefore, do not normally increase in number or size in a human adult. Under certain normal conditions, however, such as when a tissue is damaged, or during certain parts of the menstrual cycle, the capillaries begin to proliferate rapidly. This process of forming new capillaries from pre-existing blood vessels is known as angiogenesis or neovascularization. See Folkman, J. Scientific American 275, 150-154 (1996). Angiogenesis during wound healing is an example of pathophysiological neovascularization during adult life. During wound healing, the additional capillaries provide a supply of oxygen and nutrients, promote granulation tissue, and aid in waste removal. After termination of the healing process, the capillaries normally regress. Lymboussaki, A. "Vascular Endothelial Growth Factors and their Receptors in Embryos, Adults, and in Tumors" Academic

Dissertation, University of Helsinki, Molecular/Cancer Biology Laboratory and Department of Pathology, Haartman Institute, (1999).

Angiogenesis also plays an important role in the growth of cancer cells. It is known that once a nest of cancer cells reaches a certain size, roughly 1 to 2 mm in diameter, the cancer cells must develop a blood supply in order for the tumor to grow larger as diffusion will not be sufficient to supply the cancer cells with enough oxygen and nutrients. Thus, inhibition of angiogenesis is expected to halt the growth of cancer cells.

Receptor tyrosine kinases (RTKs) are transmembrane polypeptides that regulate developmental cell growth and differentiation, remodeling and regeneration of adult tissues. Mustonen, T. et al., *J. Cell Biology* 129, 895-898 (1995); van der Geer, P. et al. *Ann Rev. Cell Biol.* 10, 251-337 (1994). Polypeptide ligands known as growth factors or cytokines, are known to activate RTKs. Signaling RTKs involves ligand binding and a shift in conformation in the external domain of the receptor resulting in its dimerization. Lymboussaki, A. "Vascular Endothelial Growth Factors and their Receptors in Embryos, Adults, and in Tumors" Academic Dissertation, University of Helsinki, Molecular/Cancer Biology Laboratory and Department of Pathology, Haartman Institute, (1999); Ullrich, A. et al., *Cell* 61, 203-212 (1990). Binding of the ligand to the RTK results in receptor trans-phosphorylation at specific tyrosine residues and subsequent activation of the catalytic domains for the phosphorylation of cytoplasmic substrates. Id.

Two subfamilies of RTKs are specific to the vascular endothelium. These include the vascular endothelial growth factor (VEGF) subfamily and the Tie receptor subfamily. Class III RTKs include VEGFR-1, VEGFR-2, and VEGFR-3. Shibuya, M. et al., *Oncogene* 5, 519-525 (1990); Terman, B. et al., *Oncogene* 6, 1677-1683 (1991); Aprelikova, O. et al., *Cancer Res.* 52, 746-748 (1992).

Members of the VEGF subfamily have been described as being able to induce vascular permeability and endothelial cell proliferation and further

identified as a major inducer of angiogenesis and vasculogenesis. Ferrara, N. et al., *Endocrinol. Rev.* 18, 4-25 (1997). VEGF is known to specifically bind to RTKs including VEGFR-1 and VEGFR-2. DeVries, C. et al., *Science* 255, 989-991 (1992); Quinn, T. et al., *Proc. Natl. Acad. Sci.* 90, 7533-7537 (1993). VEGF stimulates the migration and proliferation of endothelial cells and induces angiogenesis both *in vitro* and *in vivo*. Connolly, D. et al., *J. Biol. Chem.* 264, 20017-20024 (1989); Connolly, D. et al., *J. Clin. Invest.* 84, 1470-1478 (1989); Ferrara, N. et al., *Endocrino. Rew.* 18, 4-25 (1997); Leung, D. et al., *Science* 246, 1306-1309 (1989); Plouet, J. et al., *EMBO J* 8, 3801-3806 (1989).

10 Because angiogenesis is known to be critical to the growth of cancer and to be controlled by VEGF and VEGF-RTK, substantial efforts have been undertaken to develop therapeutics that are antagonists of VEGF-RTK to thereby inhibit or retard angiogenesis, and, hopefully, interfere or stop tumor proliferation.

A wide variety of chemical compounds and compositions have been reported as having activity against one or more of the VEGF-RTKs. Examples include quinoline derivatives such as described in WO 98/13350, aminonicotinamide derivatives (see, *e.g.*, WO 01/55114), antisense compounds (see, *e.g.*, WO 01/52904), peptidomimetics (see, *e.g.*, WO 01/52875), quinazoline derivatives (see, *e.g.*, U.S. Patent No. 6,258,951) monoclonal antibodies (see, *e.g.*, EP 1 086 705 A1), various 5,10,15,20-tetraaryl-porphyrins and 5,10,15-triaryl-corroles (see, *e.g.*, WO 00/27379), heterocyclic alkanesulfonic and alkane carboxylic acid derivatives (see, *e.g.*, DE19841985), oxindolylquinazoline derivatives (see, *e.g.*, WO 99/10349), 1,4-diazaanthracene derivatives (see, *e.g.*, U.S. Patent No. 5,763,441), and cinnoline derivatives (see, *e.g.*, WO 97/34876), and various indazole compounds (see, *e.g.*, WO 01/02369 and WO 01/53268).

Various indolyl substituted compounds have recently been disclosed in WO 01/29025, WO 01/62251, and WO 01/62252, and various benzimidazolyl compounds have recently been disclosed in WO 01/28993. These compounds are reportedly capable of inhibiting, modulating, and/or regulating signal transduction

-4-

of both receptor-type and non-receptor tyrosine kinases. Some of the disclosed compounds contain a quinolone fragment bonded to the indolyl or benzimidazolyl group.

The synthesis of 4-hydroxy quinolone and 4-hydroxy quinoline derivatives is disclosed in a number of references. For example, Ukrainets et al. have disclosed the synthesis of 3-(benzimidazol-2-yl)-4-hydroxy-2-oxo-1,2-dihydroquinoline. Ukrainets, I. et al., Tet. Lett. 42, 7747-7748 (1995); Ukrainets, I. et al., Khimiya Geterotsiklicheskikh Soedinii, 2, 239-241(1992). Ukrainets has also disclosed the synthesis, anticonvulsive and antithyroid activity of other 4-hydroxy quinolones and thio analogs such as 1H-2-oxo-3-(2-benzimidazolyl)-4-hydroxyquinoline. Ukrainets, I. et al., Khimiya Geterotsiklicheskikh Soedinii, 1, 105-108 (1993); Ukrainets, I. et al., Khimiya Geterotsiklicheskikh Soedinii, 8, 1105-1108 (1993); Ukrainets, I. et al., Chem. Heterocyclic Comp. 33, 600-604, (1997).

The synthesis of various quinoline derivatives is disclosed in WO 97/48694. These compounds are disclosed as capable of binding to nuclear hormone receptors and being useful for stimulating osteoblast proliferation and bone growth. The compounds are also disclosed as being useful in the treatment or prevention of diseases associated with nuclear hormone receptor families.

Various quinoline derivatives in which the benzene ring of the quinolone is substituted with a sulfur group are disclosed in WO 92/18483. These compounds are disclosed as being useful in pharmaceutical formulations and as medicaments.

Quinolone and coumarin derivatives have been disclosed as having use in a variety of applications unrelated to medicine and pharmaceutical formulations. References that describe the preparation of quinolone derivatives for use in photopolymerizable compositions or for luminescent properties include: U.S. Patent No. 5,801,212 issued to Okamoto et al.; JP 8-29973; JP 7-43896; JP 6-9952; JP 63-258903; EP 797376; and DE 23 63 459.


-5-

Despite the exploration of a variety of chemistries to provide VEGF-RTK-antagonist therapies, a continuing need exists for compounds that inhibit the proliferation of capillaries, inhibit the growth of tumors, and/or inhibit vascular endothelial growth factor receptor tyrosine kinase and pharmaceutical formulations that contain such compounds. A need also exists for methods for administering such compounds and pharmaceutical formulations to patients in need thereof.

SUMMARY OF THE INVENTION

The present invention provides compounds, pharmaceutical formulations including the compounds, methods of preparing the pharmaceutical formulations, and methods of treating patients with the pharmaceutical formulations and compounds.

The present invention provides a first group of compounds having the structure I. The invention also provides tautomers of the compounds, pharmaceutically acceptable salts of the compounds, and pharmaceutically acceptable salts of the tautomers. Structure I has the following formula:



I

where, in the first group of compounds:

Y is selected from $-OR^{10}$ groups, $-C(=O)-R^{11}$ groups, $-NR^{12}R^{13}$ groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted

heterocyclalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclaminoalkyl groups, substituted or unsubstituted saturated heterocycl groups, substituted or unsubstituted heterocycloxyalkyl groups, substituted or unsubstituted hydroxyalkyl groups, or substituted or unsubstituted aryloxyalkyl groups;

Z is selected from O, S, or NR^{14} groups;

R^1 , R^2 , R^3 , and R^4 may be the same or different and are independently selected from H, Cl, Br, F, I, -CN, -NO₂, -OH, -OR¹⁵ groups, -NR¹⁶R¹⁷ groups, substituted or unsubstituted amidinyl groups, substituted or unsubstituted guanidinyl groups, substituted or unsubstituted primary, secondary, or tertiary alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted heterocycl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclalkyl groups, substituted or unsubstituted diheterocyclaminoalkyl groups, substituted or unsubstituted (heterocycl)(alkyl)aminoalkyl groups, substituted or unsubstituted (heterocycl)(aryl)aminoalkyl groups, or -C(=O)R¹⁸ groups;

R^5 , R^6 , R^7 , and R^8 may be the same or different and are independently selected from H, Cl, Br, F, I, -NO₂, -OH, -OR¹⁹ groups, -NR²⁰R²¹ groups, -SH, -SR²² groups, -S(=O)R²³ groups, -S(=O)₂R²⁴ groups, -CN, substituted or unsubstituted amidinyl groups, substituted or unsubstituted guanidinyl groups, substituted or unsubstituted primary, secondary, or tertiary alkyl groups, substituted

or unsubstituted aryl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted heterocyclylalkyl groups, $-C(=O)R^{25}$ groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted diheterocyclylaminoalkyl groups, substituted or unsubstituted (heterocyclyl)(alkyl)aminoalkyl groups, substituted or unsubstituted (heterocyclyl)(aryl)aminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocycloxyalkyl groups;

R^9 and R^{14} may be the same or different and are independently selected from H, -OH, substituted or unsubstituted alkoxy groups, substituted or unsubstituted aryloxy groups, $-NH_2$, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, $-C(=O)H$, $-C(=O)$ -alkyl groups, or $-C(=O)$ -aryl groups;

R^{10} is selected from substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, $-C(=O)H$, $-C(=O)$ -alkyl groups, $-C(=O)$ -aryl groups, $-C(=O)O$ -alkyl groups, $-C(=O)O$ -aryl groups, $-C(=O)NH_2$, $-C(=O)NH$ (alkyl) groups, $-C(=O)NH$ (aryl) groups, $-C(=O)N$ (alkyl)₂ groups, $-C(=O)N$ (aryl)₂ groups, $-C(=O)N$ (alkyl)(aryl) groups, $-NH_2$, $-NH$ (alkyl) groups, $-NH$ (aryl) groups, $-N$ (alkyl)₂ groups, $-N$ (alkyl)(aryl) groups, $-N$ (aryl)₂ groups, $-NH$ (heterocyclyl) groups, $-N$ (heterocyclyl)₂ groups, $-N$ (alkyl)(heterocyclyl) groups, $-N$ (aryl)(heterocyclyl), $-C(=O)NH$ (heterocyclyl) groups, $-C(=O)N$ (heterocyclyl)₂ groups, $-C(=O)N$ (alkyl)(heterocyclyl) groups,

-C(=O)N(aryl)(heterocyclyl) groups, or substituted or unsubstituted heterocyclylalkyl groups;

R¹¹ is selected from H, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups, -NH(heterocyclyl) groups, -N(heterocyclyl)₂ groups, -N(alkyl)(heterocyclyl) groups, -N(aryl)(heterocyclyl) groups, -O-alkyl groups, O-aryl groups, heterocyclylalkoxyalkyl groups, or substituted or unsubstituted aryl groups;

R¹² is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups;

R¹³ is selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, -OH, alkoxy groups, aryloxy groups, -NH₂, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -C(=O)-heterocyclyl groups, -C(=O)-O-heterocyclyl groups, -C(=O)NH(heterocyclyl) groups, -C(=O)-N(heterocyclyl)₂ groups, -C(=O)N(aryl)(heterocyclyl) groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or

unsubstituted aryloxyalkyl groups, substituted or unsubstituted heterocyclyloxyalkyl groups, or $-C(=O)-N(\text{alkyl})(\text{heterocyclyl})$ groups;

- R^{15} and R^{19} may be the same or different and are independently selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted heterocyclylalkyl groups, $-C(=O)H$, $-C(=O)-\text{alkyl}$ groups, $-C(=O)-\text{aryl}$ groups, $-C(=O)NH_2$, $-C(=O)NH(\text{alkyl})$ groups, $-C(=O)NH(\text{aryl})$ groups, $-C(=O)N(\text{alkyl})_2$ groups, $-C(=O)N(\text{aryl})_2$ groups, $-C(=O)N(\text{alkyl})(\text{aryl})$ groups, $-NH(\text{heterocyclyl})$ groups, $-N(\text{heterocyclyl})_2$ groups, $-N(\text{alkyl})(\text{heterocyclyl})$ groups, $-N(\text{aryl})(\text{heterocyclyl})$ groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl, substituted or unsubstituted diheterocyclylaminoalkyl, substituted or unsubstituted (heterocyclyl)(alkyl)aminoalkyl, substituted or unsubstituted (heterocyclyl)(aryl)aminoalkyl, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocyclyloxyalkyl groups;

R^{16} and R^{20} may be the same or different and are independently selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups;

- R^{17} and R^{21} may be the same or different and are independently selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, $-C(=O)H$, $-C(=O)-\text{alkyl}$ groups, $-C(=O)-\text{aryl}$ groups, $-C(=O)NH_2$, $-C(=O)NH(\text{alkyl})$ groups, $-C(=O)NH(\text{aryl})$ groups, $-C(=O)N(\text{alkyl})_2$ groups, $-C(=O)N(\text{aryl})_2$ groups, $-C(=O)N(\text{alkyl})(\text{aryl})$ groups, $-C(=O)O-\text{alkyl}$ groups,

- C(=O)O-aryl groups, substituted or unsubstituted heterocyclalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted
- 5 diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, -C(=O)-heterocyclalkyl groups, -C(=O)-O-heterocyclalkyl groups, -C(=O)NH(heterocyclalkyl) groups, -C(=O)-N(heterocyclalkyl)₂ groups, -C(=O)N(aryl)(heterocyclalkyl) groups, substituted or unsubstituted heterocyclalkylaminoalkyl groups, substituted or unsubstituted
- 10 diheterocyclalkylaminoalkyl groups, substituted or unsubstituted (heterocyclalkyl)(alkyl)aminoalkyl groups, substituted or unsubstituted (heterocyclalkyl)(aryl)aminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, substituted or unsubstituted heterocyclalkyloxyalkyl groups, or
- 15 -C(=O)-N(alkyl)(heterocyclalkyl) groups;

- R¹⁸, R²³, R²⁴, and R²⁵ may be the same or different and are independently selected from H, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups, -NH(heterocyclalkyl) groups, -N(heterocyclalkyl)(alkyl) groups, -N(heterocyclalkyl)(aryl) groups,
- 20 -N(heterocyclalkyl)₂ groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, -OH, substituted or unsubstituted alkoxy groups, substituted or unsubstituted heterocyclalkyl groups, substituted or unsubstituted aryloxy groups, heterocyclalkyloxy groups, -NHOH, -N(alkyl)OH groups, -N(aryl)OH groups, -N(alkyl)O-alkyl groups, -N(aryl)O-alkyl groups, -N(alkyl)O-aryl groups, or
- 25 -N(aryl)O-aryl groups; and

R²² is selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclalkyl groups.

The invention provides a second group of compounds including compounds having the structure I, tautomers of the compounds, pharmaceutically acceptable salts of the compounds, and pharmaceutically acceptable salts of the tautomers.

5 In the second group of compounds:

Y is selected from $-OR^{10}$ groups, $-C(=O)-R^{11}$ groups, $-NR^{12}R^{13}$ groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted heterocyclalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted
10 arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclaminoalkyl groups, substituted or unsubstituted saturated heterocycl groups, substituted or unsubstituted heterocycloxyalkyl groups, substituted or unsubstituted hydroxyalkyl groups, or substituted or unsubstituted
15 aryloxyalkyl groups;

Z is selected from O, S, or NR^{14} groups;

R^1 , R^2 , R^3 , and R^4 may be the same or different and are independently selected from H, Cl, Br, F, I, $-CN$, $-NO_2$, $-OH$, $-OR^{15}$ groups, $-NR^{16}R^{17}$ groups, substituted or unsubstituted amidinyl groups, substituted or
20 unsubstituted guanidinyl groups, substituted or unsubstituted primary, secondary, and tertiary alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted heterocycl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted
25 or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclalkyl groups, or $-C(=O)R^{18}$ groups;

- R^5 , R^6 , R^7 , and R^8 may be the same or different and are independently selected from H, Cl, Br, F, I, $-\text{NO}_2$, $-\text{OH}$, $-\text{OR}^{19}$ groups, $-\text{NR}^{20}\text{R}^{21}$ groups, $-\text{SH}$, $-\text{SR}^{22}$ groups, $-\text{S}(=\text{O})\text{R}^{23}$ groups, $-\text{S}(=\text{O})_2\text{R}^{24}$ groups, $-\text{CN}$, substituted or unsubstituted amidinyl groups, substituted or unsubstituted guanidinyl groups,
- 5 substituted or unsubstituted primary, secondary, and tertiary alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted heterocyclalkyl groups, $-\text{C}(=\text{O})\text{R}^{25}$ groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted
- 10 alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups,
- 15 substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocycloxyalkyl groups;

- R^9 is selected from the group consisting of $-\text{OH}$, substituted or unsubstituted alkoxy groups, substituted or unsubstituted aryloxy groups, $-\text{NH}_2$, substituted or unsubstituted alkylamino groups, substituted or unsubstituted
- 20 arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, $-\text{C}(=\text{O})\text{H}$, $-\text{C}(=\text{O})$ -alkyl groups, or $-\text{C}(=\text{O})$ -aryl groups;

- R^{10} is selected from substituted or unsubstituted aryl groups,
- 25 substituted or unsubstituted heterocyclyl groups, $-\text{C}(=\text{O})\text{H}$, $-\text{C}(=\text{O})$ -alkyl groups, $-\text{C}(=\text{O})$ -aryl groups, $-\text{C}(=\text{O})\text{O}$ -alkyl groups, $-\text{C}(=\text{O})\text{O}$ -aryl groups, $-\text{C}(=\text{O})\text{NH}_2$, $-\text{C}(=\text{O})\text{NH}(\text{alkyl})$ groups, $-\text{C}(=\text{O})\text{NH}(\text{aryl})$ groups, $-\text{C}(=\text{O})\text{N}(\text{alkyl})_2$ groups, $-\text{C}(=\text{O})\text{N}(\text{aryl})_2$ groups, $-\text{C}(=\text{O})\text{N}(\text{alkyl})(\text{aryl})$ groups, $-\text{NH}_2$, $-\text{NH}(\text{alkyl})$ groups, $-\text{NH}(\text{aryl})$ groups, $-\text{N}(\text{alkyl})_2$ groups, $-\text{N}(\text{alkyl})(\text{aryl})$ groups, $-\text{N}(\text{aryl})_2$ groups,
- 30 $-\text{C}(=\text{O})\text{NH}(\text{heterocyclyl})$ groups, $-\text{C}(=\text{O})\text{N}(\text{heterocyclyl})_2$ groups,

-C(=O)N(alkyl)(heterocyclyl) groups, -C(=O)N(aryl)(heterocyclyl) groups, or substituted or unsubstituted heterocyclylalkyl groups;

R¹¹ is selected from H, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups, -NH(heterocyclyl) groups, -N(heterocyclyl)₂ groups, -N(alkyl)(heterocyclyl) groups, -O-alkyl groups, O-aryl groups, substituted or unsubstituted alkyl groups, or substituted or unsubstituted aryl groups;

R¹² is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups;

R¹³ is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, -OH, alkoxy groups, aryloxy groups, -NH₂, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, -C(=O)-heterocyclyl groups, -C(=O)-O-heterocyclyl groups, -C(=O)NH(heterocyclyl) groups, -C(=O)-N(heterocyclyl)₂ groups, -C(=O)N(aryl)(heterocyclyl) groups, -C(=O)-N(alkyl)(heterocyclyl) groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups,

substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocycloxyalkyl groups;

R¹⁴ is selected from H, -OH, substituted or unsubstituted alkoxy groups, substituted or unsubstituted aryloxy groups, -NH₂, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, -C(=O)H, -C(=O)-alkyl groups, or -C(=O)-aryl groups;

R¹⁵ and R¹⁹ may be the same or different and are independently selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted heterocyclylalkyl groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl, substituted or unsubstituted diheterocyclylaminoalkyl, substituted or unsubstituted (heterocyclyl)(alkyl)aminoalkyl, substituted or unsubstituted (heterocyclyl)(aryl)aminoalkyl, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocycloxyalkyl groups;

R¹⁶ and R²⁰ may be the same or different and are independently selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups;

- R^{17} and R^{21} may be the same or different and are independently selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, $-C(=O)H$, $-C(=O)$ -alkyl groups, $-C(=O)$ -aryl groups, $-C(=O)NH_2$,
 5 $-C(=O)NH$ (alkyl) groups, $-C(=O)NH$ (aryl) groups, $-C(=O)N$ (alkyl)₂ groups, $-C(=O)N$ (aryl)₂ groups, $-C(=O)N$ (alkyl)(aryl) groups, $-C(=O)O$ -alkyl groups, $-C(=O)O$ -aryl groups, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups,
 10 substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, $-C(=O)$ -heterocyclyl groups, $-C(=O)-O$ -heterocyclyl groups, $-C(=O)NH$ (heterocyclyl) groups, $-C(=O)-N$ (heterocyclyl)₂ groups, $-C(=O)N$ (aryl)(heterocyclyl) groups, $-C(=O)-N$ (alkyl)(heterocyclyl) groups,
 15 substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocyclioxyalkyl groups;

- R^{18} , R^{23} , R^{24} , and R^{25} may be the same or different and are
 20 independently selected from H, $-NH_2$, $-NH$ (alkyl) groups, $-NH$ (aryl) groups, $-N$ (alkyl)₂ groups, $-N$ (aryl)₂ groups, $-N$ (alkyl)(aryl) groups, $-NH$ (heterocyclyl) groups, $-N$ (heterocyclyl)(alkyl) groups, $-N$ (heterocyclyl)(aryl) groups, $-N$ (heterocyclyl)₂ groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, $-OH$, substituted or unsubstituted alkoxy groups,
 25 substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted aryloxy groups, $-NHOH$, $-N$ (alkyl)OH groups, $-N$ (aryl)OH groups, $-N$ (alkyl)O-alkyl groups, $-N$ (aryl)O-alkyl groups, $-N$ (alkyl)O-aryl groups, or $-N$ (aryl)O-aryl groups;
 and

-16-

R^{22} is selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups.

The invention provides a third group of compounds including
5 compounds having the structure I, tautomers of the compounds, pharmaceutically acceptable salts of the compounds, and pharmaceutically acceptable salts of the tautomers.

In the third group of compounds:

Y is selected from -OH, SH, alkylthio groups, arylthio groups, -OR¹⁰
10 groups, -C(=O)-R¹¹ groups, -NR¹²R¹³ groups, -CN, substituted or unsubstituted alkyl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted aralkyl groups, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups,
15 substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyoxyalkyl groups, substituted or
20 unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, or substituted or unsubstituted aryloxyalkyl groups;

Z is selected from O, S, or NR¹⁴ groups;

R¹, R², R³, and R⁴ may be the same or different and are independently selected from H, Cl, Br, F, I, -CN, -NO₂, -OH, -OR¹⁵ groups,
25 -NR¹⁶R¹⁷ groups, substituted or unsubstituted amidinyl groups, substituted or unsubstituted guanidinyl groups, substituted or unsubstituted primary, secondary, or tertiary alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups,

substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylalkyl groups, or $-C(=O)R^{18}$ groups;

R^5 , R^6 , R^7 , and R^8 may be the same or different and are independently selected from H, Cl, Br, F, I, $-NO_2$, $-OH$, $-OR^{19}$ groups, $-NR^{20}R^{21}$ groups, $-SH$, $-SR^{22}$ groups, $-S(=O)R^{23}$ groups, $-S(=O)_2R^{24}$ groups, $-CN$, substituted or unsubstituted amidinyl groups, substituted or unsubstituted guanidinyl groups, substituted or unsubstituted primary, secondary, or tertiary alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylalkyl groups, $-C(=O)R^{25}$ groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocycloxyalkyl groups;

R^9 and R^{14} may be the same or different and are independently selected from H, $-OH$, substituted or unsubstituted alkoxy groups, substituted or unsubstituted aryloxy groups, $-NH_2$, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, $-C(=O)H$, $-C(=O)$ -alkyl groups, or $-C(=O)$ -aryl groups;

R¹⁰ is selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted heterocyclylalkyl groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(alkyl)(aryl) groups, -N(aryl)₂ groups, -C(=O)NH(heterocyclyl) groups, -C(=O)N(heterocyclyl)₂ groups, -C(=O)N(alkyl)(heterocyclyl) groups, or -C(=O)N(aryl)(heterocyclyl) groups;

R¹¹ is selected from H, -OH, alkoxy groups, aryloxy groups, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups, substituted or unsubstituted alkyl groups, -NH(heterocyclyl) groups, -N(heterocyclyl)₂ groups, -N(alkyl)(heterocyclyl) groups, or substituted or unsubstituted aryl groups;

R¹² is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups;

R¹³ is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, -OH, alkoxy groups, aryloxy groups, -NH₂, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, -C(=O)H, -C(=O)-alkyl groups,

-C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, -C(=O)NH₂,
 -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups,
 -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -C(=O)-heterocyclyl
 groups, -C(=O)-O-heterocyclyl groups, -C(=O)NH(heterocyclyl) groups,
 5 -C(=O)N(heterocyclyl)₂ groups, -C(=O)N(alkyl)(heterocyclyl) groups,
 -C(=O)N(aryl)(heterocyclyl) groups, substituted or unsubstituted
 heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups,
 substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted
 aryloxyalkyl groups, or substituted or unsubstituted heterocycliloxyalkyl groups;

10 R¹⁵ and R¹⁹ may be the same or different and are independently
 selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted
 aryl groups, substituted or unsubstituted heterocyclyl groups, substituted or
 unsubstituted heterocyclylalkyl groups, -C(=O)H, -C(=O)-alkyl groups,
 -C(=O)-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl)
 15 groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl)
 groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted
 alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups,
 substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted
 diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl
 20 groups, substituted or unsubstituted heterocyclylaminoalkyl, substituted or
 unsubstituted diheterocyclylaminoalkyl, substituted or unsubstituted
 (heterocyclyl)(alkyl)aminoalkyl, substituted or unsubstituted
 (heterocyclyl)(aryl)aminoalkyl, substituted or unsubstituted alkoxyalkyl groups,
 substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted
 25 aryloxyalkyl groups, or substituted or unsubstituted heterocycliloxyalkyl groups;

R¹⁶ and R²⁰ may be the same or different and are independently
 selected from H, substituted or unsubstituted alkyl groups, substituted or
 unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups;

-20-

R¹⁷ and R²¹ may be the same or different and are independently selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)NH₂,
 5 -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups,
 10 substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, -C(=O)-heterocyclyl groups, -C(=O)-O-heterocyclyl groups, -C(=O)NH(heterocyclyl) groups, -C(=O)-N(heterocyclyl)₂ groups, -C(=O)N(alkyl)(heterocyclyl) groups, -C(=O)-N(aryl)(heterocyclyl) groups,
 15 substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocyclyloxyalkyl groups;

R¹⁸, R²³, R²⁴, and R²⁵ may be the same or different and are
 20 independently selected from H, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups, -NH(heterocyclyl) groups, -N(heterocyclyl)(alkyl) groups, -N(heterocyclyl)(aryl) groups, -N(heterocyclyl)₂ groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, -OH, substituted or unsubstituted alkoxy groups,
 25 substituted or unsubstituted aryloxy groups, substituted or unsubstituted heterocyclyl groups, -NHOH, -N(alkyl)OH groups, -N(aryl)OH groups, -N(alkyl)O-alkyl groups, -N(aryl)O-alkyl groups, -N(alkyl)O-aryl groups, or -N(aryl)O-aryl groups;
 and

R^{22} is selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups.

- In the third group of compounds, at least one of R^5 , R^6 , R^7 , or R^8 is
- 5 selected from substituted or unsubstituted amidinyl groups, substituted or unsubstituted guanidinyl groups, substituted or unsubstituted saturated heterocyclyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or
 - 10 unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocyclyloxyalkyl groups; $-OR^{19}$ groups where R^{19}
 - 15 is selected from substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted heterocyclylalkyl groups, $-C(=O)H$, $-C(=O)$ -aryl groups, $-C(=O)NH_2$, $-C(=O)NH(alkyl)$ groups, $-C(=O)NH(aryl)$ groups, $-C(=O)N(alkyl)_2$ groups, $-C(=O)N(aryl)_2$ groups, $-C(=O)N(alkyl)(aryl)$ groups, substituted or unsubstituted aminoalkyl groups,
 - 20 substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted diheterocyclylaminoalkyl groups, substituted or
 - 25 unsubstituted (heterocyclyl)(alkyl)aminoalkyl groups, substituted or unsubstituted (heterocyclyl)(aryl)aminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted and unsubstituted heterocyclyloxyalkyl groups; $-NR^{20}R^{21}$ groups where R^{20} is selected from substituted or unsubstituted heterocyclyl
 - 30 groups; $-NR^{20}R^{21}$ groups where R^{21} is selected from substituted or unsubstituted heterocyclyl groups, $-C(=O)H$, $-C(=O)$ -aryl groups, $-C(=O)NH_2$,

- C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, substituted or unsubstituted heterocyclylalkyl groups, or substituted or unsubstituted heterocycloxyalkyl groups; or -C(=O)R²⁵ groups where R²⁵ is selected from H, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups, -NH(heterocyclyl) groups, -N(heterocyclyl)(alkyl) groups, -N(heterocyclyl)(aryl) groups, -N(heterocyclyl)₂ groups, substituted or unsubstituted aryl groups, substituted or unsubstituted aryloxy groups, or substituted or unsubstituted heterocyclyl groups.

The invention provides a fourth group of compounds having the structure I, tautomers of the compounds, pharmaceutically acceptable salts of the compounds, and pharmaceutically acceptable salts of the tautomers.

- In the fourth group of compounds:

- Y is selected from -OH, SH, alkylthio groups, arylthio groups, -OR¹⁰ groups, -C(=O)-R¹¹ groups, -NR¹²R¹³ groups, -CN, substituted or unsubstituted alkyl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted aralkyl groups, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or

unsubstituted heterocyclyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocycloxyalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, or substituted or unsubstituted aryloxyalkyl groups;

5 Z is selected from O, S, or NR^{14} groups;

R^1 , R^2 , R^3 , and R^4 may be the same or different and are independently selected from H, Cl, Br, F, I, -CN, -NO₂, -OH, -OR¹⁵ groups, -NR¹⁶R¹⁷ groups, substituted or unsubstituted amidinyl groups, substituted or unsubstituted guanidinyl groups, substituted or unsubstituted primary, secondary, or tertiary alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylalkyl groups, or -C(=O)R¹⁸ groups;

R^5 , R^6 , R^7 , and R^8 may be the same or different and are independently selected from H, Cl, Br, F, I, -NO₂, -OH, -OR¹⁹ groups, -NR²⁰R²¹ groups, -SH, -SR²² groups, -S(=O)R²³ groups, -S(=O)₂R²⁴ groups, -CN, substituted or unsubstituted amidinyl groups, substituted or unsubstituted guanidinyl groups, substituted or unsubstituted primary, secondary, or tertiary alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylalkyl groups, -C(=O)R²⁵ groups, substituted or unsubstituted aminoalkyl groups,

substituted or unsubstituted heterocyclaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocycloxyalkyl groups;

- 5 R^9 and R^{14} may be the same or different and are independently selected from H, -OH, substituted or unsubstituted alkoxy groups, substituted or unsubstituted aryloxy groups, -NH₂, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, -C(=O)H, -C(=O)-alkyl groups, or -C(=O)-aryl groups;

- R^{10} is selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocycl groups, substituted or unsubstituted heterocyclalkyl groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(alkyl)(aryl) groups, -N(aryl)₂ groups, -C(=O)NH(heterocycl) groups, -C(=O)N(heterocycl)₂ groups, -C(=O)N(alkyl)(heterocycl) groups, or -C(=O)N(aryl)(heterocycl) groups;

- R^{11} is selected from H, -OH, alkoxy groups, aryloxy groups, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups, substituted or unsubstituted alkyl groups, -NH(heterocycl) groups, -N(heterocycl)₂ groups, -N(alkyl)(heterocycl) groups, or substituted or unsubstituted aryl groups;

R^{12} is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups;

R^{13} is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, -OH, alkoxy groups, aryloxy groups, -NH₂, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -C(=O)-heterocyclyl groups, -C(=O)-O-heterocyclyl groups, -C(=O)NH(heterocyclyl) groups, -C(=O)N(heterocyclyl)₂ groups, -C(=O)N(alkyl)(heterocyclyl) groups, -C(=O)N(aryl)(heterocyclyl) groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocyclyloxyalkyl groups;

R^{15} and R^{19} may be the same or different and are independently selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted heterocyclylalkyl groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted

alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl, substituted or
 5 unsubstituted diheterocyclylaminoalkyl, substituted or unsubstituted (heterocyclyl)(alkyl)aminoalkyl, substituted or unsubstituted (heterocyclyl)(aryl)aminoalkyl, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocyclyloxyalkyl groups;

10 R^{16} and R^{20} may be the same or different and are independently selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups;

R^{17} and R^{21} may be the same or different and are independently selected from H, substituted or unsubstituted alkyl groups, substituted or
 15 unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, $-C(=O)H$, $-C(=O)$ -alkyl groups, $-C(=O)$ -aryl groups, $-C(=O)NH_2$, $-C(=O)NH$ (alkyl) groups, $-C(=O)NH$ (aryl) groups, $-C(=O)N$ (alkyl)₂ groups, $-C(=O)N$ (aryl)₂ groups, $-C(=O)N$ (alkyl)(aryl) groups, $-C(=O)O$ -alkyl groups, $-C(=O)O$ -aryl groups, substituted or unsubstituted heterocyclylalkyl groups,
 20 substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, $-C(=O)$ -heterocyclyl groups, $-C(=O)$ -O-heterocyclyl groups,
 25 $-C(=O)NH$ (heterocyclyl) groups, $-C(=O)$ -N(heterocyclyl)₂ groups, $-C(=O)$ -N(alkyl)(heterocyclyl) groups, $-C(=O)$ -N(aryl)(heterocyclyl) groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted
 30 heterocyclyloxyalkyl groups;

-27-

R^{18} , R^{23} , R^{24} , and R^{25} may be the same or different and are independently selected from H, $-NH_2$, $-NH(\text{alkyl})$ groups, $-NH(\text{aryl})$ groups, $-N(\text{alkyl})_2$ groups, $-N(\text{aryl})_2$ groups, $-N(\text{alkyl})(\text{aryl})$ groups, $-NH(\text{heterocyclyl})$ groups, $-N(\text{heterocyclyl})(\text{alkyl})$ groups, $-N(\text{heterocyclyl})(\text{aryl})$ groups, $-N(\text{heterocyclyl})_2$ groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, $-OH$, substituted or unsubstituted alkoxy groups, substituted or unsubstituted aryloxy groups, substituted or unsubstituted heterocyclyl groups, $-NHOH$, $-N(\text{alkyl})OH$ groups, $-N(\text{aryl})OH$ groups, $-N(\text{alkyl})O\text{-alkyl}$ groups, $-N(\text{aryl})O\text{-alkyl}$ groups, $-N(\text{alkyl})O\text{-aryl}$ groups, or $-N(\text{aryl})O\text{-aryl}$ groups;

5
10 and

R^{22} is selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups.

In the fourth group of compounds, at least one of R^1 , R^2 , R^3 , or R^4 is an $-OR^{15}$ group and R^{15} is selected from substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted diheterocyclylaminoalkyl groups, substituted or unsubstituted (heterocyclyl)(alkyl)aminoalkyl groups, or substituted or unsubstituted (heterocyclyl)(aryl)aminoalkyl groups.

15
20

Preferred compounds in any of the first, second, or third groups described above are provided in which Z is an $-NR^{14}$. Preferred compound of the fourth group are also provided in which Z is an $-NR^{10}$ group.

25

Preferred compounds according to the first, second, third, and fourth groups of compounds are provided in which Y is an $-OR^{10}$ group, an $-NR^{12}R^{13}$ group, or a substituted or unsubstituted alkynyl group.


5 Other preferred compounds of the first, second, third, and fourth groups are provided in which R^1 is selected from H, substituted or unsubstituted alkoxy groups, substituted or unsubstituted heterocyclalkoxy groups, substituted or unsubstituted heterocycloxy groups, or substituted or unsubstituted heterocycl groups.

10 Still further provided are compounds of the first, second, third, and fourth groups in which R^2 is selected from the group consisting of H, F, Cl, $-NO_2$, substituted and unsubstituted heterocyclalkoxy groups, and substituted and unsubstituted heterocycl groups.

15 Still further provided are compounds of the first, second, third, and fourth groups in which R^6 or R^7 is an alkyl group. Still further preferred compounds of the four groups are those in which R^6 or R^7 is an $-OR^{19}$ group and R^{19} is an alkyl group, an aryl group, a heterocycl group, or a heterocyclalkyl group.

20 Preferred compounds of the fourth group of compounds are provided in which R^1 is an $-OR^{15}$ group and R^{15} is selected from substituted or unsubstituted heterocyclalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocycl groups, 25 substituted or unsubstituted heterocyclaminoalkyl groups, substituted or unsubstituted diheterocyclaminoalkyl groups, substituted or unsubstituted (heterocycl)(alkyl)aminoalkyl groups, or substituted or unsubstituted (heterocycl)(aryl)aminoalkyl groups.

The invention further provides compounds having the structure II. The invention provides tautomers of the compounds, pharmaceutically acceptable salts of the compounds, and pharmaceutically acceptable salts of the tautomers. Structure II has the following formula



II

where:

- Y is selected from H, -OH, -OR¹⁰ groups, -SH, -SR¹¹ groups, -NR¹²R¹³ groups, -CN, -C(=O)-R¹⁴ groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted aralkyl groups, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocyclyloxyalkyl groups;
- X¹, X², X³, and X⁴ are selected from C or N, and at least one of X¹, X², X³, and X⁴ is N;

-30-

R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 may be the same or different and are independently selected from H, Cl, Br, F, I, $-\text{NO}_2$, $-\text{CN}$, $-\text{OH}$, $-\text{OR}^{15}$ groups, $-\text{NR}^{16}\text{R}^{17}$ groups, $-\text{C}(=\text{O})\text{R}^{18}$ groups, $-\text{SH}$, $-\text{SR}^{19}$ groups, $-\text{S}(=\text{O})\text{R}^{20}$ groups, $\text{S}(=\text{O})_2\text{R}^{21}$ groups, substituted or unsubstituted amidinyl groups, substituted or unsubstituted guanidinyl groups, substituted or unsubstituted primary, secondary, or tertiary alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocyclyloxyalkyl groups; R^5 is absent or is H if X^1 is N; R^6 is absent or is H if X^2 is N; R^7 is absent or is H if X^3 is N; and R^8 is absent or is H if X^4 is N;

R^9 is selected from H, $-\text{OH}$, substituted or unsubstituted alkoxy groups, substituted or unsubstituted aryloxy groups, $-\text{NH}_2$, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, $-\text{C}(=\text{O})\text{H}$, $-\text{C}(=\text{O})$ -alkyl groups, or $-\text{C}(=\text{O})$ -aryl groups;

R^{10} is selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted heterocyclylalkyl groups, $-\text{C}(=\text{O})\text{H}$, $-\text{C}(=\text{O})$ -alkyl groups, $-\text{C}(=\text{O})$ -aryl groups, $-\text{C}(=\text{O})\text{O}$ -alkyl groups, $-\text{C}(=\text{O})\text{O}$ -aryl groups, $-\text{C}(=\text{O})\text{NH}_2$, $-\text{C}(=\text{O})\text{NH}(\text{alkyl})$ groups, $-\text{C}(=\text{O})\text{NH}(\text{aryl})$ groups,

- C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(alkyl)(aryl) groups, -N(aryl)₂ groups, -C(=O)NH(heterocyclyl) groups, -C(=O)N(heterocyclyl)₂ groups, -C(=O)N(alkyl)(heterocyclyl) groups, or
- 5 -C(=O)N(aryl)(heterocyclyl) groups;

R¹¹ and R¹⁹ may be the same or different and are independently selected from substituted or unsubstituted alkyl groups, or substituted or unsubstituted aryl groups;

- R¹² is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups;
- 10

- R¹³ is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, -OH, alkoxy groups, aryloxy groups, -NH₂, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -C(=O)-heterocyclyl groups, -C(=O)-O-heterocyclyl groups, -C(=O)NH(heterocyclyl) groups, -C(=O)N(heterocyclyl)₂ groups, -C(=O)N(alkyl)(heterocyclyl) groups, -C(=O)N(aryl)(heterocyclyl) groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups,
- 15
- 20
- 25

substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocycloxyalkyl groups;

- R^{14} is selected from H, -OH, alkoxy groups, aryloxy groups, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, -NH(heterocyclyl) groups, -N(heterocyclyl)₂ groups, -N(alkyl)(heterocyclyl) groups, or -N(aryl)(heterocyclyl) groups;

R^{12} and R^{13} may join together to form a 5 to 7 membered saturated or unsaturated, substituted or unsubstituted N-containing ring;

- R^{15} is selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted heterocyclylalkyl groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted diheterocyclylaminoalkyl groups, substituted or unsubstituted (heterocyclyl)(alkyl)aminoalkyl groups, substituted or unsubstituted (heterocyclyl)(aryl)aminoalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, substituted or unsubstituted hydroxyalkyl groups, or substituted or unsubstituted heterocycloxyalkyl groups;

R^{16} is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups;

R¹⁷ is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, 5 -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted 10 (aryl)(alkyl)aminoalkyl groups, substituted or unsubstituted heterocyclylalkyl groups, -C(=O)-heterocyclyl groups, -C(=O)-O-heterocyclyl groups, -C(=O)NH(heterocyclyl) groups, -C(=O)-N(heterocyclyl)₂ groups, -C(=O)-N(alkyl)(heterocyclyl) groups, -C(=O)-N(aryl)(heterocyclyl) groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or 15 unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, substituted or unsubstituted heterocyclyoxyalkyl groups, -OH, substituted or unsubstituted alkoxy groups, substituted or unsubstituted aryloxy groups, or -NH₂ groups;

R¹⁶ and R¹⁷ may join together to form a 5 to 7 membered saturated or 20 unsaturated, substituted or unsubstituted N-containing ring; and

R¹⁸, R²⁰, and R²¹ may be the same or different and are independently selected from H, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, -OH, substituted or unsubstituted alkoxy 25 groups, substituted or unsubstituted aryloxy groups, substituted or unsubstituted heterocyclyl groups, -NHOH, -N(alkyl)OH groups, -N(aryl)OH groups, -N(alkyl)O-alkyl groups, -N(aryl)O-alkyl groups, -N(alkyl)O-aryl groups, or -N(aryl)O-aryl groups.

Preferred compounds having structure II are also provided where Y is selected from H, -OH, -OR¹⁰ groups, or -NR¹²R¹³ groups.

Still other preferred compounds having structure II are provided where at least two of X¹, X², X³, and X⁴ are C and the corresponding substituents
5 R⁵, R⁶, R⁷, and R⁸ are hydrogen, and at least one of X¹, X², X³, and X⁴ is N, and the rest of the compound is consistent with any of the above-described compounds.

Still other more preferred compounds of structure II are provided in which R⁶ or R⁷ is an alkyl group and the rest of the compound is consistent with any of the above-described compounds.

10 Still other compounds of structure II are provided in which R⁶ or R⁷ is an -OR¹⁵ group and R¹⁵ is an alkyl, aryl, heterocyclyl, or heterocyclylalkyl group and the rest of the molecule is consistent with any of the above-described compounds.

Still further compounds having the formula of structure II are
15 provided in which R¹ is selected from H, substituted or unsubstituted alkoxy groups, substituted or unsubstituted heterocyclylalkoxy groups, substituted or unsubstituted heterocycloxy groups, or substituted or unsubstituted heterocyclyl groups.

Still other compounds having the Structure II are provided in which
In other compounds having the structure II, R² is selected from H, F, Cl, -NO₂,
20 substituted or unsubstituted heterocyclyl groups, or substituted or unsubstituted heterocyclylalkoxy groups.

Pharmaceutical formulations according to the present invention are provided which include any of the compounds described above in combination with a pharmaceutically acceptable carrier.

25 A method of treating a patient in need of an inhibitor of vascular endothelial growth factor receptor tyrosine kinase is provided which includes

-35-

administering an effective amount of the pharmaceutical formulation according to the present invention to a patient in need thereof.

Further objects, features and advantages of the invention will be apparent from the following detailed description.

5

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides novel compounds that act as antagonists of receptor tyrosine kinases, and, more particularly, as inhibitors of bFGF and/or VEGF-RTK function. The compounds provided herein can be formulated into pharmaceutical formulations that are useful in treating patients with
10 a need for an inhibitor of VEGF-RTK, especially, in particular embodiments, to provide compositions and methods for reducing capillary proliferation and in the treatment of cancer.

The following abbreviations and definitions are used throughout this application:

15

“VEGF” is an abbreviation that stands for vascular endothelial growth factor.

“RTK” is an abbreviation that stands for receptor tyrosine kinase.

“VEGF-RTK” is an abbreviation that stands for vascular endothelial growth factor receptor tyrosine kinase.

20

“Flt-1” is an abbreviation that stands for fms-like tyrosine kinase-1, also known as vascular endothelial growth factor receptor-1 or “VEGFR1”.

“KDR” is an abbreviation that stands for kinase- insert domain-containing receptor, also known as vascular endothelial growth factor receptor-2 or “VEGFR2”.

"bFGF" is an abbreviation that stands for basic fibroblast growth factor.

"bFGFR" is an abbreviation that stands for basic fibroblast growth factor receptor.

- 5 Generally, reference to a certain element such as hydrogen or H is meant to include all isotopes of that element. For example, if an R group is defined to include hydrogen or H, it also includes deuterium and tritium.

 The phrase "unsubstituted alkyl" refers to alkyl groups that do not contain heteroatoms. Thus the phrase includes straight chain alkyl groups such as methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl and the like. The phrase also includes branched chain isomers of straight chain alkyl groups, including but not limited to, the following which are provided by way of example: $-\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, $-\text{CH}(\text{CH}_2\text{CH}_3)_2$, $-\text{C}(\text{CH}_3)_3$, $-\text{C}(\text{CH}_2\text{CH}_3)_3$, $-\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}_2\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, $-\text{CH}_2\text{CH}(\text{CH}_2\text{CH}_3)_2$, $-\text{CH}_2\text{C}(\text{CH}_3)_3$, $-\text{CH}_2\text{C}(\text{CH}_2\text{CH}_3)_3$, $-\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_2\text{CH}_3)_2$, $-\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)_3$, $-\text{CH}_2\text{CH}_2\text{C}(\text{CH}_2\text{CH}_3)_3$, $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_2\text{CH}_3)\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, and others. The phrase also includes cyclic alkyl groups such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl and such rings substituted with straight and branched chain alkyl groups as defined above. The phrase also includes polycyclic alkyl groups such as, but not limited to, adamantyl, norbornyl, and bicyclo[2.2.2]octyl and such rings substituted with straight and branched chain alkyl groups as defined above. Thus, the phrase unsubstituted alkyl groups includes primary alkyl groups, secondary alkyl groups, and tertiary alkyl groups. Unsubstituted alkyl groups may be bonded to one or more carbon atom(s), oxygen atom(s), nitrogen atom(s), and/or sulfur atom(s) in the parent compound. Preferred unsubstituted alkyl groups include straight and branched chain alkyl groups and cyclic alkyl groups having 1 to 20 carbon atoms. More preferred such

10

15

20

25

unsubstituted alkyl groups have from 1 to 10 carbon atoms while even more preferred such groups have from 1 to 5 carbon atoms. Most preferred unsubstituted alkyl groups include straight and branched chain alkyl groups having from 1 to 3 carbon atoms and include methyl, ethyl, propyl, and $-\text{CH}(\text{CH}_3)_2$.

- 5 The phrase "substituted alkyl" refers to an unsubstituted alkyl group as defined above in which one or more bonds to a carbon(s) or hydrogen(s) are replaced by a bond to non-hydrogen and non-carbon atoms such as, but not limited to, a halogen atom in halides such as F, Cl, Br, and I; and oxygen atom in groups such as hydroxyl groups, alkoxy groups, aryloxy groups, and ester groups; a sulfur
- 10 atom in groups such as thiol groups, alkyl and aryl sulfide groups, sulfone groups, sulfonyl groups, and sulfoxide groups; a nitrogen atom in groups such as amines, amides, alkylamines, dialkylamines, arylamines, alkylarylamines, diarylamines, N-oxides, imides, and enamines; a silicon atom in groups such as in trialkylsilyl groups, dialkylarylsilyl groups, alkylarylsilyl groups, and triarylsilyl groups; and
- 15 other heteroatoms in various other groups. Substituted alkyl groups also include groups in which one or more bonds to a carbon(s) or hydrogen(s) atom is replaced by a bond to a heteroatom such as oxygen in carbonyl, carboxyl, and ester groups; nitrogen in groups such as imines, oximes, hydrazones, and nitriles. Preferred substituted alkyl groups include, among others, alkyl groups in which one or more
- 20 bonds to a carbon or hydrogen atom is/are replaced by one or more bonds to fluorine atoms. One example of a substituted alkyl group is the trifluoromethyl group and other alkyl groups that contain the trifluoromethyl group. Other alkyl groups include those in which one or more bonds to a carbon or hydrogen atom is replaced by a bond to an oxygen atom such that the substituted alkyl group contains
- 25 a hydroxyl, alkoxy, aryloxy group, or heterocycloxy group. Still other alkyl groups include alkyl groups that have an amine, alkylamine, dialkylamine, arylamine, (alkyl)(aryl)amine, diarylamine, heterocyclylamine, (alkyl)(heterocyclyl)amine, (aryl)(heterocyclyl)amine, or diheterocyclylamine group.

The phrase "unsubstituted aryl" refers to aryl groups that do not contain heteroatoms. Thus the phrase includes, but is not limited to, groups such as phenyl, biphenyl, anthracenyl, naphthenyl by way of example. Although the phrase "unsubstituted aryl" includes groups containing condensed rings such as naphthalene, it does not include aryl groups that have other groups such as alkyl or halo groups bonded to one of the ring members, as aryl groups such as tolyl are considered herein to be substituted aryl groups as described below. A preferred unsubstituted aryl group is phenyl. Unsubstituted aryl groups may be bonded to one or more carbon atom(s), oxygen atom(s), nitrogen atom(s), and/or sulfur atom(s) in the parent compound, however.

The phrase "substituted aryl group" has the same meaning with respect to unsubstituted aryl groups that substituted alkyl groups had with respect to unsubstituted alkyl groups. However, a substituted aryl group also includes aryl groups in which one of the aromatic carbons is bonded to one of the non-carbon or non-hydrogen atoms described above and also includes aryl groups in which one or more aromatic carbons of the aryl group is bonded to a substituted and/or unsubstituted alkyl, alkenyl, or alkynyl group as defined herein. This includes bonding arrangements in which two carbon atoms of an aryl group are bonded to two atoms of an alkyl, alkenyl, or alkynyl group to define a fused ring system (e.g. dihydronaphthyl or tetrahydronaphthyl). Thus, the phrase "substituted aryl" includes, but is not limited to tolyl, and hydroxyphenyl among others.

The phrase "unsubstituted alkenyl" refers to straight and branched chain and cyclic groups such as those described with respect to unsubstituted alkyl groups as defined above, except that at least one double bond exists between two carbon atoms. Examples include, but are not limited to vinyl, $-\text{CH}=\text{C}(\text{H})(\text{CH}_3)$, $-\text{CH}=\text{C}(\text{CH}_3)_2$, $-\text{C}(\text{CH}_3)=\text{C}(\text{H})_2$, $-\text{C}(\text{CH}_3)=\text{C}(\text{H})(\text{CH}_3)$, $-\text{C}(\text{CH}_2\text{CH}_3)=\text{CH}_2$, cyclohexenyl, cyclopentenyl, cyclohexadienyl, butadienyl, pentadienyl, and hexadienyl among others.

The phrase "substituted alkenyl" has the same meaning with respect to unsubstituted alkenyl groups that substituted alkyl groups had with respect to unsubstituted alkyl groups. A substituted alkenyl group includes alkenyl groups in which a non-carbon or non-hydrogen atom is bonded to a carbon double bonded to another carbon and those in which one of the non-carbon or non-hydrogen atoms is bonded to a carbon not involved in a double bond to another carbon.

The phrase "unsubstituted alkynyl" refers to straight and branched chain groups such as those described with respect to unsubstituted alkyl groups as defined above, except that at least one triple bond exists between two carbon atoms. Examples include, but are not limited to $-\text{C}\equiv\text{C}(\text{H})$, $-\text{C}\equiv\text{C}(\text{CH}_3)$, $-\text{C}\equiv\text{C}(\text{CH}_2\text{CH}_3)$, $-\text{C}(\text{H})_2\text{C}\equiv\text{C}(\text{H})$, $-\text{C}(\text{H})_2\text{C}\equiv\text{C}(\text{CH}_3)$, and $-\text{C}(\text{H})_2\text{C}\equiv\text{C}(\text{CH}_2\text{CH}_3)$ among others.

The phrase "substituted alkynyl" has the same meaning with respect to unsubstituted alkynyl groups that substituted alkyl groups had with respect to unsubstituted alkyl groups. A substituted alkynyl group includes alkynyl groups in which a non-carbon or non-hydrogen atom is bonded to a carbon triple bonded to another carbon and those in which a non-carbon or non-hydrogen atom is bonded to a carbon not involved in a triple bond to another carbon.

The phrase "unsubstituted aralkyl" refers to unsubstituted alkyl groups as defined above in which a hydrogen or carbon bond of the unsubstituted alkyl group is replaced with a bond to an aryl group as defined above. For example, methyl ($-\text{CH}_3$) is an unsubstituted alkyl group. If a hydrogen atom of the methyl group is replaced by a bond to a phenyl group, such as if the carbon of the methyl were bonded to a carbon of benzene, then the compound is an unsubstituted aralkyl group (*i.e.*, a benzyl group). Thus the phrase includes, but is not limited to, groups such as benzyl, diphenylmethyl, and 1-phenylethyl ($-\text{CH}(\text{C}_6\text{H}_5)(\text{CH}_3)$) among others.

The phrase "substituted aralkyl" has the same meaning with respect to unsubstituted aralkyl groups that substituted aryl groups had with respect to unsubstituted aryl groups. However, a substituted aralkyl group also includes

groups in which a carbon or hydrogen bond of the alkyl part of the group is replaced by a bond to a non-carbon or a non-hydrogen atom. Examples of substituted aralkyl groups include, but are not limited to, $-\text{CH}_2\text{C}(=\text{O})(\text{C}_6\text{H}_5)$, and $-\text{CH}_2(2\text{-methylphenyl})$ among others.

- 5 The phrase "unsubstituted heterocyclyl" refers to both aromatic and nonaromatic ring compounds including monocyclic, bicyclic, and polycyclic ring compounds such as, but not limited to, quinuclidyl, containing 3 or more ring members of which one or more is a heteroatom such as, but not limited to, N, O, and S. Although the phrase "unsubstituted heterocyclyl" includes condensed
- 10 heterocyclic rings such as benzimidazolyl, it does not include heterocyclyl groups that have other groups such as alkyl or halo groups bonded to one of the ring members as compounds such as 2-methylbenzimidazolyl are substituted heterocyclyl groups. Examples of heterocyclyl groups include, but are not limited to: unsaturated 3 to 8 membered rings containing 1 to 4 nitrogen atoms such as, but not
- 15 limited to pyrrolyl, pyrrolinyl, imidazolyl, pyrazolyl, pyridyl, dihydropyridyl, pyrimidyl, pyrazinyl, pyridazinyl, triazolyl (e.g. 4H-1,2,4-triazolyl, 1H-1,2,3-triazolyl, 2H-1,2,3-triazolyl etc.), tetrazolyl, (e.g. 1H-tetrazolyl, 2H tetrazolyl, etc.); saturated 3 to 8 membered rings containing 1 to 4 nitrogen atoms such as, but not limited to, pyrrolidinyl, imidazolidinyl, piperidinyl, piperazinyl; condensed
- 20 unsaturated heterocyclic groups containing 1 to 4 nitrogen atoms such as, but not limited to, indolyl, isoindolyl, indolynyl, indolizynyl, benzimidazolyl, quinolyl, isoquinolyl, indazolyl, benzotriazolyl; unsaturated 3 to 8 membered rings containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms such as, but not limited to, oxazolyl, isoxazolyl, oxadiazolyl (e.g. 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-
- 25 oxadiazolyl, etc.); saturated 3 to 8 membered rings containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms such as, but not limited to, morpholinyl; unsaturated condensed heterocyclic groups containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms, for example, benzoxazolyl, benzoxadiazolyl, benzoxazinyl (e.g. 2H-1,4-benzoxazinyl etc.); unsaturated 3 to 8 membered rings containing 1 to 3 sulfur
- 30 atoms and 1 to 3 nitrogen atoms such as, but not limited to, thiazolyl, isothiazolyl, thiadiazolyl (e.g. 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-

- thiadiazolyl, etc.); saturated 3 to 8 membered rings containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms such as, but not limited to, thiazolodiny; saturated and unsaturated 3 to 8 membered rings containing 1 to 2 sulfur atoms such as, but not limited to, thienyl, dihydrodithiynyl, dihydrodithionyl, tetrahydrothiophene, 5 tetrahydrothiopyran; unsaturated condensed heterocyclic rings containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms such as, but not limited to, benzothiazolyl, benzothiadiazolyl, benzothiazinyl (e.g. 2H-1,4-benzothiazinyl, etc.), dihydrobenzothiazinyl (e.g. 2H-3,4-dihydrobenzothiazinyl, etc.), unsaturated 3 to 8 membered rings containing oxygen atoms such as, but not limited to furyl; 10 unsaturated condensed heterocyclic rings containing 1 to 2 oxygen atoms such as benzodioxolyl (e.g. 1,3-benzodioxolyl, etc.); unsaturated 3 to 8 membered rings containing an oxygen atom and 1 to 2 sulfur atoms such as, but not limited to, dihydrooxathiinyl; saturated 3 to 8 membered rings containing 1 to 2 oxygen atoms and 1 to 2 sulfur atoms such as 1,4-oxathiane; unsaturated condensed rings 15 containing 1 to 2 sulfur atoms such as benzothieryl, benzodithiynyl; and unsaturated condensed heterocyclic rings containing an oxygen atom and 1 to 2 oxygen atoms such as benzoxathiinyl. Heterocyclyl group also include those described above in which one or more S atoms in the ring is double-bonded to one or two oxygen atoms (sulfoxides and sulfones). For example, heterocyclyl groups include 20 tetrahydrothiophene, tetrahydrothiophene oxide, and tetrahydrothiophene 1,1-dioxide. Preferred heterocyclyl groups contain 5 or 6 ring members. More preferred heterocyclyl groups include morpholine, piperazine, piperidine, pyrrolidine, imidazole, pyrazole, 1,2,3-triazole, 1,2,4-triazole, tetrazole, thiomorpholine, thiomorpholine in which the S atom of the thiomorpholine is 25 bonded to one or more O atoms, pyrrole, homopiperazine, oxazolidin-2-one, pyrrolidin-2-one, oxazole, quinuclidine, thiazole, isoxazole, furan, and tetrahydrofuran.

- The phrase "substituted heterocyclyl" refers to an unsubstituted heterocyclyl group as defined above in which one of the ring members is bonded to 30 a non-hydrogen atom such as described above with respect to substituted alkyl groups and substituted aryl groups. Examples, include, but are not limited to, 2-

methylbenzimidazolyl, 5-methylbenzimidazolyl, 5-chlorobenzthiazolyl, 1-methyl piperazinyl, and 2-chloropyridyl among others.

The phrase "unsubstituted heterocyclalkyl" refers to unsubstituted alkyl groups as defined above in which a hydrogen or carbon bond of the
5 unsubstituted alkyl group is replaced with a bond to a heterocycl group as defined above. For example, methyl ($-\text{CH}_3$) is an unsubstituted alkyl group. If a hydrogen atom of the methyl group is replaced by a bond to a heterocycl group, such as if the carbon of the methyl were bonded to carbon 2 of pyridine (one of the carbons bonded to the N of the pyridine) or carbons 3 or 4 of the pyridine, then the
10 compound is an unsubstituted heterocyclalkyl group.

The phrase "substituted heterocyclalkyl" has the same meaning with respect to unsubstituted heterocyclalkyl groups that substituted aralkyl groups had with respect to unsubstituted aralkyl groups. However, a substituted heterocyclalkyl group also includes groups in which a non-hydrogen atom is
15 bonded to a heteroatom in the heterocycl group of the heterocyclalkyl group such as, but not limited to, a nitrogen atom in the piperidine ring of a piperidinylalkyl group.

The phrase "unsubstituted alkylaminoalkyl" refers to an unsubstituted alkyl group as defined above in which a carbon or hydrogen bond is replaced by a
20 bond to a nitrogen atom that is bonded to a hydrogen atom and an unsubstituted alkyl group as defined above. For example, methyl ($-\text{CH}_3$) is an unsubstituted alkyl group. If a hydrogen atom of the methyl group is replaced by a bond to a nitrogen atom that is bonded to a hydrogen atom and an ethyl group, then the resulting compound is $-\text{CH}_2-\text{N}(\text{H})(\text{CH}_2\text{CH}_3)$ which is an unsubstituted alkylaminoalkyl
25 group.

The phrase "substituted alkylaminoalkyl" refers to an unsubstituted alkylaminoalkyl group as defined above except where one or more bonds to a carbon or hydrogen atom in one or both of the alkyl groups is replaced by a bond to a non-carbon or non-hydrogen atom as described above with respect to substituted

alkyl groups except that the bond to the nitrogen atom in all alkylaminoalkyl groups does not by itself qualify all alkylaminoalkyl groups as being substituted. However, substituted alkylaminoalkyl groups does include groups in which the hydrogen bonded to the nitrogen atom of the group is replaced with a non-carbon and non-
5 hydrogen atom.

The phrase "unsubstituted dialkylaminoalkyl" refers to an unsubstituted alkyl group as defined above in which a carbon bond or hydrogen bond is replaced by a bond to a nitrogen atom which is bonded to two other similar or different unsubstituted alkyl groups as defined above.

10 The phrase "substituted dialkylaminoalkyl" refers to an unsubstituted dialkylaminoalkyl group as defined above in which one or more bonds to a carbon or hydrogen atom in one or more of the alkyl groups is replaced by a bond to a non-carbon and non-hydrogen atom as described with respect to substituted alkyl groups. The bond to the nitrogen atom in all dialkylaminoalkyl groups does not by
15 itself qualify all dialkylaminoalkyl groups as being substituted.

The phrase "unsubstituted heterocycloxyalkyl" refers to an unsubstituted alkyl group as defined above in which a carbon bond or hydrogen bond is replaced by a bond to an oxygen atom which is bonded to an unsubstituted heterocyclyl group as defined above.

20 The phrase "substituted heterocycloxyalkyl" refers to an unsubstituted heterocycloxyalkyl group as defined above in which a bond to a carbon or hydrogen group of the alkyl group of the heterocycloxyalkyl group is bonded to a non-carbon and non-hydrogen atom as described above with respect to substituted alkyl groups or in which the heterocyclyl group of the
25 heterocycloxyalkyl group is a substituted heterocyclyl group as defined above.

The phrase "unsubstituted arylaminoalkyl" refers to an unsubstituted alkyl group as defined above in which a carbon bond or hydrogen bond is replaced

by a bond to a nitrogen atom which is bonded to at least one unsubstituted aryl group as defined above.

The phrase "substituted arylaminoalkyl" refers to an unsubstituted arylaminoalkyl group as defined above except where either the alkyl group of the
5 arylaminoalkyl group is a substituted alkyl group as defined above or the aryl group of the arylaminoalkyl group is a substituted aryl group except that the bonds to the nitrogen atom in all arylaminoalkyl groups does not by itself qualify all arylaminoalkyl groups as being substituted. However, substituted arylaminoalkyl groups does include groups in which the hydrogen bonded to the nitrogen atom of
10 the group is replaced with a non-carbon and non-hydrogen atom.

The phrase "unsubstituted heterocyclaminoalkyl" refers to an unsubstituted alkyl group as defined above in which a carbon or hydrogen bond is replaced by a bond to a nitrogen atom which is bonded to at least one unsubstituted heterocyclyl group as defined above.

15 The phrase "substituted heterocyclaminoalkyl" refers to unsubstituted heterocyclaminoalkyl groups as defined above in which the heterocyclyl group is a substituted heterocyclyl group as defined above and/or the alkyl group is a substituted alkyl group as defined above. The bonds to the nitrogen atom in all heterocyclaminoalkyl groups does not by itself qualify all
20 heterocyclaminoalkyl groups as being substituted. However, substituted heterocyclaminoalkyl groups do include groups in which the hydrogen bonded to the nitrogen atom of the group is replaced with a non-carbon and non-hydrogen atom.

The phrase "unsubstituted alkylaminoalkoxy" refers to an
25 unsubstituted alkyl group as defined above in which a carbon or hydrogen bond is replaced by a bond to an oxygen atom which is bonded to the parent compound and in which another carbon or hydrogen bond of the unsubstituted alkyl group is bonded to a nitrogen atom which is bonded to a hydrogen atom and an unsubstituted alkyl group as defined above.

The phrase "substituted alkylaminoalkoxy" refers to unsubstituted alkylaminoalkoxy groups as defined above in which a bond to a carbon or hydrogen atom of the alkyl group bonded to the oxygen atom which is bonded to the parent compound is replaced by one or more bonds to a non-carbon and non-hydrogen atoms as discussed above with respect to substituted alkyl groups and/or if the hydrogen bonded to the amino group is bonded to a non-carbon and non-hydrogen atom and/or if the alkyl group bonded to the nitrogen of the amine is bonded to a non-carbon and non-hydrogen atom as described above with respect to substituted alkyl groups. The presence of the amine and alkoxy functionality in all alkylaminoalkoxy groups does not by itself qualify all such groups as substituted alkylaminoalkoxy groups.

The phrase "unsubstituted dialkylaminoalkoxy" refers to an unsubstituted alkyl group as defined above in which a carbon or hydrogen bond is replaced by a bond to an oxygen atom which is bonded to the parent compound and in which another carbon or hydrogen bond of the unsubstituted alkyl group is bonded to a nitrogen atom which is bonded to two other similar or different unsubstituted alkyl groups as defined above.

The phrase "substituted dialkylaminoalkoxy" refers to an unsubstituted dialkylaminoalkoxy group as defined above in which a bond to a carbon or hydrogen atom of the alkyl group bonded to the oxygen atom which is bonded to the parent compound is replaced by one or more bonds to a non-carbon and non-hydrogen atoms as discussed above with respect to substituted alkyl groups and/or if one or more of the alkyl groups bonded to the nitrogen of the amine is bonded to a non-carbon and non-hydrogen atom as described above with respect to substituted alkyl groups. The presence of the amine and alkoxy functionality in all dialkylaminoalkoxy groups does not by itself qualify all such groups as substituted dialkylaminoalkoxy groups.

The phrase "unsubstituted heterocyclyloxy" refers to a hydroxyl group (-OH) in which the bond to the hydrogen atom is replaced by a bond to a ring atom of an otherwise unsubstituted heterocyclyl group as defined above.


5 The phrase "substituted heterocyclyloxy" refers to a hydroxyl group (-OH) in which the bond to the hydrogen atom is replaced by a bond to a ring atom of an otherwise substituted heterocyclyl group as defined above.

The term "protected" with respect to hydroxyl groups, amine groups, and sulfhydryl groups refers to forms of these functionalities which are protected from undesirable reaction with a protecting group known to those skilled in the art
10 such as those set forth in Protective Groups in Organic Synthesis, Greene, T.W.; Wuts, P. G. M., John Wiley & Sons, New York, NY, (3rd Edition, 1999) which can be added or removed using the procedures set forth therein. Examples of protected hydroxyl groups include, but are not limited to, silyl ethers such as those obtained by reaction of a hydroxyl group with a reagent such as, but not limited to,
15 t-butyldimethyl-chlorosilane, trimethylchlorosilane, triisopropylchlorosilane, triethylchlorosilane; substituted methyl and ethyl ethers such as, but not limited to methoxymethyl ether, methythiomethyl ether, benzyloxymethyl ether, t-butoxymethyl ether, 2-methoxyethoxymethyl ether, tetrahydropyranyl ethers, 1-ethoxyethyl ether, allyl ether, benzyl ether; esters such as, but not limited to,
20 benzoylformate, formate, acetate, trichloroacetate, and trifluoroacetate. Examples of protected amine groups include, but are not limited to, amides such as, formamide, acetamide, trifluoroacetamide, and benzamide; imides, such as phthalimide, and dithiosuccinimide; and others. Examples of protected sulfhydryl groups include, but are not limited to, thioethers such as S-benzyl thioether, and S-4-picolyl
25 thioether; substituted S-methyl derivatives such as hemithio, dithio and aminothio acetals; and others.

A "pharmaceutically acceptable salt" includes a salt with an inorganic base, organic base, inorganic acid, organic acid, or basic or acidic amino acid. As salts of inorganic bases, the invention includes, for example, alkali metals

such as sodium or potassium; alkaline earth metals such as calcium and magnesium or aluminum; and ammonia. As salts of organic bases, the invention includes, for example, trimethylamine, triethylamine, pyridine, picoline, ethanolamine, diethanolamine, and triethanolamine. As salts of inorganic acids, the instant
5 invention includes, for example, hydrochloric acid, hydroboric acid, nitric acid, sulfuric acid, and phosphoric acid. As salts of organic acids, the instant invention includes, for example, formic acid, acetic acid, trifluoroacetic acid, fumaric acid, oxalic acid, tartaric acid, maleic acid, citric acid, succinic acid, malic acid, methanesulfonic acid, benzenesulfonic acid, and p-toluenesulfonic acid. As salts of
10 basic amino acids, the instant invention includes, for example, arginine, lysine and ornithine. Acidic amino acids include, for example, aspartic acid and glutamic acid.

Generally, the invention provides compounds including having the structure I. The invention also provides tautomers of the compounds,
15 pharmaceutically acceptable salts of the compounds, and pharmaceutically acceptable salts of the tautomers. Structure I has the following formula:



I

Preferred compounds having structure I are those within one of four groups.

In the first group of compounds:

Y is selected from $-OR^{10}$ groups, $-C(=O)-R^{11}$ groups, $-NR^{12}R^{13}$ groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted saturated heterocyclyl groups, substituted or unsubstituted heterocyclyloxyalkyl groups, substituted or unsubstituted hydroxyalkyl groups, or substituted or unsubstituted aryloxyalkyl groups;

Z is selected from O, S, or NR^{14} groups;

R^1 , R^2 , R^3 , and R^4 may be the same or different and are independently selected from H, Cl, Br, F, I, $-CN$, $-NO_2$, $-OH$, $-OR^{15}$ groups, $-NR^{16}R^{17}$ groups, substituted or unsubstituted amidinyl groups, substituted or unsubstituted guanidinyl groups, substituted or unsubstituted primary, secondary, or tertiary alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted diheterocyclylaminoalkyl groups, substituted or unsubstituted (heterocyclyl)(alkyl)aminoalkyl groups, substituted or unsubstituted (heterocyclyl)(aryl)aminoalkyl groups, or $-C(=O)R^{18}$ groups;

R^5 , R^6 , R^7 , and R^8 may be the same or different and are independently selected from H, Cl, Br, F, I, $-NO_2$, $-OH$, $-OR^{19}$ groups, $-NR^{20}R^{21}$

groups, -SH, -SR²² groups, -S(=O)R²³ groups, -S(=O)₂R²⁴ groups, -CN, substituted or unsubstituted amidinyl groups, substituted or unsubstituted guanidinyl groups, substituted or unsubstituted primary, secondary, or tertiary alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted heterocyclylalkyl groups, -C(=O)R²⁵ groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted diheterocyclylaminoalkyl groups, substituted or unsubstituted (heterocyclyl)(alkyl)aminoalkyl groups, substituted or unsubstituted (heterocyclyl)(aryl)aminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocycliloxyalkyl groups;

R⁹ and R¹⁴ may be the same or different and are independently selected from H, -OH, substituted or unsubstituted alkoxy groups, substituted or unsubstituted aryloxy groups, -NH₂, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, -C(=O)H, -C(=O)-alkyl groups, or -C(=O)-aryl groups;

R¹⁰ is selected from substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(alkyl)(aryl) groups, -N(aryl)₂ groups,

-NH(heterocyclyl) groups, -N(heterocyclyl)₂ groups, -N(alkyl)(heterocyclyl) groups, -N(aryl)(heterocyclyl), -C(=O)NH(heterocyclyl) groups, -C(=O)N(heterocyclyl)₂ groups, -C(=O)N(alkyl)(heterocyclyl) groups, -C(=O)N(aryl)(heterocyclyl) groups, or substituted or unsubstituted
 5 heterocyclylalkyl groups;

R¹¹ is selected from H, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups, -NH(heterocyclyl) groups, -N(heterocyclyl)₂ groups, -N(alkyl)(heterocyclyl) groups, -N(aryl)(heterocyclyl) groups, -O-alkyl groups, O-aryl groups, heterocycloxyalkyl
 10 groups, or substituted or unsubstituted aryl groups;

R¹² is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups;

R¹³ is selected from substituted or unsubstituted alkyl groups,
 15 substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, -OH, alkoxy groups, aryloxy groups, -NH₂, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups,
 20 substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, -C(=O)H, -C(=O)-alkyl groups,
 25 -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -C(=O)-heterocyclyl groups, -C(=O)-O-heterocyclyl groups, -C(=O)NH(heterocyclyl) groups, -C(=O)-N(heterocyclyl)₂ groups, -C(=O)N(aryl)(heterocyclyl) groups, substituted

or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, substituted or unsubstituted heterocyclyloxyalkyl groups, or $-C(=O)-N(\text{alkyl})(\text{heterocyclyl})$ groups;

- 5 R^{15} and R^{19} may be the same or different and are independently selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted heterocyclylalkyl groups, $-C(=O)H$, $-C(=O)-\text{alkyl}$ groups, $-C(=O)-\text{aryl}$ groups, $-C(=O)NH_2$, $-C(=O)NH(\text{alkyl})$ groups, $-C(=O)NH(\text{aryl})$ groups, $-C(=O)N(\text{alkyl})_2$ groups, $-C(=O)N(\text{aryl})_2$ groups, $-C(=O)N(\text{alkyl})(\text{aryl})$ groups, $-NH(\text{heterocyclyl})$ groups, $-N(\text{heterocyclyl})_2$ groups, $-N(\text{alkyl})(\text{heterocyclyl})$ groups, $-N(\text{aryl})(\text{heterocyclyl})$ groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl, substituted or unsubstituted diheterocyclylaminoalkyl, substituted or unsubstituted (heterocyclyl)(alkyl)aminoalkyl, substituted or unsubstituted (heterocyclyl)(aryl)aminoalkyl, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocyclyloxyalkyl groups;
- 10
- 15
- 20

- R^{16} and R^{20} may be the same or different and are independently selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups;
- 25

R^{17} and R^{21} may be the same or different and are independently selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, $-C(=O)H$, $-C(=O)-\text{alkyl}$ groups, $-C(=O)-\text{aryl}$ groups, $-C(=O)NH_2$,

- C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, -C(=O)-heterocyclyl groups, -C(=O)-O-heterocyclyl groups, -C(=O)NH(heterocyclyl) groups, -C(=O)-N(heterocyclyl)₂ groups, -C(=O)N(aryl)(heterocyclyl) groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted diheterocyclylaminoalkyl groups, substituted or unsubstituted (heterocyclyl)(alkyl)aminoalkyl groups, substituted or unsubstituted (heterocyclyl)(aryl)aminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, substituted or unsubstituted heterocyclyloxyalkyl groups, or -C(=O)-N(alkyl)(heterocyclyl) groups;

- R¹⁸, R²³, R²⁴, and R²⁵ may be the same or different and are independently selected from H, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups, -NH(heterocyclyl) groups, -N(heterocyclyl)(alkyl) groups, -N(heterocyclyl)(aryl) groups, -N(heterocyclyl)₂ groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, -OH, substituted or unsubstituted alkoxy groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted aryloxy groups, heterocyclyloxy groups, -NHOH, -N(alkyl)OH groups, -N(aryl)OH groups, -N(alkyl)O-alkyl groups, -N(aryl)O-alkyl groups, -N(alkyl)O-aryl groups, or -N(aryl)O-aryl groups; and

- R²² is selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups.

In other embodiments of the invention, compounds of formula I above include a second group of compounds having the substituents described below:

Y is selected from $-OR^{10}$ groups, $-C(=O)-R^{11}$ groups, $-NR^{12}R^{13}$ groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted heterocyclalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclaminoalkyl groups, substituted or unsubstituted saturated heterocycl groups, substituted or unsubstituted heterocycloxyalkyl groups, substituted or unsubstituted hydroxyalkyl groups, or substituted or unsubstituted aryloxyalkyl groups;

Z is selected from O, S, or NR^{14} groups;

R^1 , R^2 , R^3 , and R^4 may be the same or different and are independently selected from H, Cl, Br, F, I, $-CN$, $-NO_2$, $-OH$, $-OR^{15}$ groups, $-NR^{16}R^{17}$ groups, substituted or unsubstituted amidinyl groups, substituted or unsubstituted guanidinyl groups, substituted or unsubstituted primary, secondary, and tertiary alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted heterocycl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclalkyl groups, or $-C(=O)R^{18}$ groups;

R^5 , R^6 , R^7 , and R^8 may be the same or different and are independently selected from H, Cl, Br, F, I, $-NO_2$, $-OH$, $-OR^{19}$ groups, $-NR^{20}R^{21}$ groups, $-SH$, $-SR^{22}$ groups, $-S(=O)R^{23}$ groups, $-S(=O)_2R^{24}$ groups, $-CN$, substituted

or unsubstituted amidinyl groups, substituted or unsubstituted guanidinyl groups, substituted or unsubstituted primary, secondary, and tertiary alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted heterocyclylalkyl groups, -C(=O)R²⁵ groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocyclyoxyalkyl groups;

R⁹ is selected from the group consisting of -OH, substituted or unsubstituted alkoxy groups, substituted or unsubstituted aryloxy groups, -NH₂, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, -C(=O)H, -C(=O)-alkyl groups, or -C(=O)-aryl groups;

R¹⁰ is selected from substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(alkyl)(aryl) groups, -N(aryl)₂ groups, -C(=O)NH(heterocyclyl) groups, -C(=O)N(heterocyclyl)₂ groups, -C(=O)N(alkyl)(heterocyclyl) groups, -C(=O)N(aryl)(heterocyclyl) groups, or substituted or unsubstituted heterocyclylalkyl groups;

R¹¹ is selected from H, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups, -NH(heterocyclyl) groups, -N(heterocyclyl)₂ groups, -N(alkyl)(heterocyclyl) groups, -O-alkyl groups, O-aryl groups, substituted or unsubstituted alkyl groups, or substituted or
 5 unsubstituted aryl groups;

R¹² is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups;

R¹³ is selected from H, substituted or unsubstituted alkyl groups,
 10 substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, -OH, alkoxy groups, aryloxy groups, -NH₂, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, -C(=O)H, -C(=O)-alkyl
 15 groups, -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or
 20 unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, -C(=O)-heterocyclyl groups, -C(=O)-O-heterocyclyl groups, -C(=O)NH(heterocyclyl) groups, -C(=O)N(heterocyclyl)₂ groups, -C(=O)N(aryl)(heterocyclyl) groups,
 25 -C(=O)N(alkyl)(heterocyclyl) groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocycliloxyalkyl groups;

R¹⁴ is selected from H, -OH, substituted or unsubstituted alkoxy groups, substituted or unsubstituted aryloxy groups, -NH₂, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, -C(=O)H, -C(=O)-alkyl groups, or -C(=O)-aryl groups;

R¹⁵ and R¹⁹ may be the same or different and are independently selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted heterocyclylalkyl groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl, substituted or unsubstituted diheterocyclylaminoalkyl, substituted or unsubstituted (heterocyclyl)(alkyl)aminoalkyl, substituted or unsubstituted (heterocyclyl)(aryl)aminoalkyl, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocycloxyalkyl groups;

R¹⁶ and R²⁰ may be the same or different and are independently selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups;

R¹⁷ and R²¹ may be the same or different and are independently selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups,

-57-

- C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)NH₂,
 -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups,
 -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -C(=O)O-alkyl groups,
 -C(=O)O-aryl groups, substituted or unsubstituted heterocyclylalkyl groups,
 5 substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted
 alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups,
 substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted
 diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl
 groups, -C(=O)-heterocyclyl groups, -C(=O)-O-heterocyclyl groups,
 10 -C(=O)NH(heterocyclyl) groups, -C(=O)-N(heterocyclyl)₂ groups,
 -C(=O)N(aryl)(heterocyclyl) groups, -C(=O)-N(alkyl)(heterocyclyl) groups,
 substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or
 unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups,
 substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted
 15 heterocyclyloxyalkyl groups;

- R¹⁸, R²³, R²⁴, and R²⁵ may be the same or different and are
 independently selected from H, -NH₂, -NH(alkyl) groups, -NH(aryl) groups,
 -N(alkyl)₂ groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups, -NH(heterocyclyl)
 groups, -N(heterocyclyl)(alkyl) groups, -N(heterocyclyl)(aryl) groups,
 20 -N(heterocyclyl)₂ groups, substituted or unsubstituted alkyl groups, substituted or
 unsubstituted aryl groups, -OH, substituted or unsubstituted alkoxy groups,
 substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted aryloxy
 groups, -NHOH, -N(alkyl)OH groups, -N(aryl)OH groups, -N(alkyl)O-alkyl
 groups, -N(aryl)O-alkyl groups, -N(alkyl)O-aryl groups, or -N(aryl)O-aryl groups;
 25 and

R²² is selected from substituted or unsubstituted alkyl groups,
 substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl
 groups.

In another embodiment, the present invention provides a third group of compounds having the general formula I above with substituents selected from the following:

Y is selected from -OH, SH, alkylthio groups, arylthio groups, -OR¹⁰ groups, -C(=O)-R¹¹ groups, -NR¹²R¹³ groups, -CN, substituted or unsubstituted alkyl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted aralkyl groups, substituted or unsubstituted heterocyclalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclaminoalkyl groups, substituted or unsubstituted heterocycl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocycloxyalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, or substituted or unsubstituted aryloxyalkyl groups;

Z is selected from O, S, or NR¹⁴ groups;

R¹, R², R³, and R⁴ may be the same or different and are independently selected from H, Cl, Br, F, I, -CN, -NO₂, -OH, -OR¹⁵ groups, -NR¹⁶R¹⁷ groups, substituted or unsubstituted amidinyl groups, substituted or unsubstituted guanidinyl groups, substituted or unsubstituted primary, secondary, or tertiary alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted heterocycl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclalkyl groups, or -C(=O)R¹⁸;

R^5 , R^6 , R^7 , and R^8 may be the same or different and are independently selected from H, Cl, Br, F, I, $-NO_2$, $-OH$, $-OR^{19}$ groups, $-NR^{20}R^{21}$ groups, $-SH$, $-SR^{22}$ groups, $-S(=O)R^{23}$ groups, $-S(=O)_2R^{24}$ groups, $-CN$, substituted or unsubstituted amidinyl groups, substituted or unsubstituted guanidinyl groups, substituted or unsubstituted primary, secondary, or tertiary alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylalkyl groups, $-C(=O)R^{25}$ groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocyclioxyalkyl groups;

R^9 and R^{14} may be the same or different and are independently selected from H, $-OH$, substituted or unsubstituted alkoxy groups, substituted or unsubstituted aryloxy groups, $-NH_2$, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, $-C(=O)H$, $-C(=O)$ -alkyl groups, or $-C(=O)$ -aryl groups;

R^{10} is selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted heterocyclylalkyl groups, $-C(=O)H$, $-C(=O)$ -alkyl groups, $-C(=O)$ -aryl groups, $-C(=O)O$ -alkyl groups, $-C(=O)O$ -aryl groups, $-C(=O)NH_2$, $-C(=O)NH$ (alkyl) groups, $-C(=O)NH$ (aryl) groups, $-C(=O)N$ (alkyl)₂ groups, $-C(=O)N$ (aryl)₂ groups, $-C(=O)N$ (alkyl)(aryl) groups,

-NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(alkyl)(aryl) groups, -N(aryl)₂ groups, -C(=O)NH(heterocyclyl) groups, -C(=O)N(heterocyclyl)₂ groups, -C(=O)N(alkyl)(heterocyclyl) groups, or -C(=O)N(aryl)(heterocyclyl) groups;

5 R¹¹ is selected from H, -OH, alkoxy groups, aryloxy groups, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups, substituted or unsubstituted alkyl groups, -NH(heterocyclyl) groups, -N(heterocyclyl)₂ groups, -N(alkyl)(heterocyclyl) groups, or substituted or unsubstituted aryl groups;

10 R¹² is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups;

R¹³ is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, -OH, alkoxy groups, aryloxy groups, -NH₂, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -C(=O)-heterocyclyl groups, -C(=O)-O-heterocyclyl groups, -C(=O)NH(heterocyclyl) groups, -C(=O)N(heterocyclyl)₂ groups, -C(=O)N(alkyl)(heterocyclyl) groups, -C(=O)N(aryl)(heterocyclyl) groups, substituted or unsubstituted

15

20

25

heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocyclyloxyalkyl groups;

- R¹⁵ and R¹⁹ may be the same or different and are independently
- 5 selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted heterocyclylalkyl groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl)
- 10 groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl, substituted or
- 15 unsubstituted diheterocyclylaminoalkyl, substituted or unsubstituted (heterocyclyl)(alkyl)aminoalkyl, substituted or unsubstituted (heterocyclyl)(aryl)aminoalkyl, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocyclyloxyalkyl groups;

- 20 R¹⁶ and R²⁰ may be the same or different and are independently selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups;

- R¹⁷ and R²¹ may be the same or different and are independently selected from H, substituted or unsubstituted alkyl groups, substituted or
- 25 unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, substituted or unsubstituted heterocyclylalkyl groups,

substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted
 alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups,
 substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted
 diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl
 5 groups, -C(=O)-heterocyclyl groups, -C(=O)-O-heterocyclyl groups,
 -C(=O)NH(heterocyclyl) groups, -C(=O)N(heterocyclyl)₂ groups,
 -C(=O)N(alkyl)(heterocyclyl) groups, -C(=O)N(aryl)(heterocyclyl) groups,
 substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or
 unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups,
 10 substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted
 heterocyclioxyalkyl groups;

R¹⁸, R²³, R²⁴, and R²⁵ may be the same or different and are
 independently selected from H, -NH₂, -NH(alkyl) groups, -NH(aryl) groups,
 -N(alkyl)₂ groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups, -NH(heterocyclyl)
 15 groups, -N(heterocyclyl)(alkyl) groups, -N(heterocyclyl)(aryl) groups,
 -N(heterocyclyl)₂ groups, substituted or unsubstituted alkyl groups, substituted or
 unsubstituted aryl groups, -OH, substituted or unsubstituted alkoxy groups,
 substituted or unsubstituted aryloxy groups, substituted or unsubstituted heterocyclyl
 groups, -NHOH, -N(alkyl)OH groups, -N(aryl)OH groups, -N(alkyl)O-alkyl
 20 groups, -N(aryl)O-alkyl groups, -N(alkyl)O-aryl groups, or -N(aryl)O-aryl groups;
 and

R²² is selected from substituted or unsubstituted alkyl groups,
 substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl
 groups.

25 In the third group of compounds, at least one of R⁵, R⁶, R⁷, or R⁸ is
 selected from substituted or unsubstituted amidinyl groups, substituted or
 unsubstituted guanidinyl groups, substituted or unsubstituted saturated heterocyclyl
 groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or
 unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl

- groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclalkyl groups, substituted or unsubstituted heterocyclaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or
- 5 unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocycloxyalkyl groups; $-OR^{19}$ groups where R^{19} is selected from substituted or unsubstituted aryl groups, substituted or unsubstituted heterocycl groups, substituted or unsubstituted heterocyclalkyl groups, $-C(=O)H$, $-C(=O)$ -aryl groups, $-C(=O)NH_2$, $-C(=O)NH(alkyl)$ groups,
- 10 $-C(=O)NH(aryl)$ groups, $-C(=O)N(alkyl)_2$ groups, $-C(=O)N(aryl)_2$ groups, $-C(=O)N(alkyl)(aryl)$ groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted
- 15 (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclaminoalkyl groups, substituted or unsubstituted diheterocyclaminoalkyl groups, substituted or unsubstituted (heterocycl)(alkyl)aminoalkyl groups, substituted or unsubstituted (heterocycl)(aryl)aminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted
- 20 aryloxyalkyl groups, or substituted and unsubstituted heterocycloxyalkyl groups; $-NR^{20}R^{21}$ groups where R^{20} is selected from substituted or unsubstituted heterocycl groups; $-NR^{20}R^{21}$ groups where R^{21} is selected from substituted or unsubstituted heterocycl groups, $-C(=O)H$, $-C(=O)$ -aryl groups, $-C(=O)NH_2$, $-C(=O)NH(alkyl)$ groups, $-C(=O)NH(aryl)$ groups, $-C(=O)N(alkyl)_2$ groups,
- 25 $-C(=O)N(aryl)_2$ groups, $-C(=O)N(alkyl)(aryl)$ groups, $-C(=O)O$ -alkyl groups, $-C(=O)O$ -aryl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted
- 30 (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or

unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, substituted or unsubstituted heterocyclalkyl groups, or substituted or unsubstituted heterocycloxyalkyl groups; or $-C(=O)R^{25}$ groups where R^{25} is selected from H, $-NH_2$, $-NH(alkyl)$ groups, $-NH(aryl)$ groups, $-N(alkyl)_2$ groups, $-N(aryl)_2$ groups, $-N(alkyl)(aryl)$ groups, $-NH(heterocycl)$ groups, $-N(heterocycl)(alkyl)$ groups, $-N(heterocycl)(aryl)$ groups, $-N(heterocycl)_2$ groups, substituted or unsubstituted aryl groups, substituted or unsubstituted aryloxy groups, or substituted or unsubstituted heterocycl groups.

In yet another embodiment, the present invention encompasses compounds of formula I in which the substituents described below define a fourth group of compounds:

Y is selected from $-OH$, SH , alkylthio groups, arylthio groups, $-OR^{10}$ groups, $-C(=O)-R^{11}$ groups, $-NR^{12}R^{13}$ groups, $-CN$, substituted or unsubstituted alkyl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted aralkyl groups, substituted or unsubstituted heterocyclalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted $(alkyl)(aryl)aminoalkyl$ groups, substituted or unsubstituted heterocyclaminoalkyl groups, substituted or unsubstituted heterocycl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocycloxyalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, or substituted or unsubstituted aryloxyalkyl groups;

Z is selected from O, S, or NR^{14} groups;

R^1 , R^2 , R^3 , and R^4 may be the same or different and are independently selected from H, Cl, Br, F, I, $-CN$, $-NO_2$, $-OH$, $-OR^{15}$ groups, $-NR^{16}R^{17}$ groups, substituted or unsubstituted amidinyl groups, substituted or unsubstituted guanidinyl groups, substituted or unsubstituted primary, secondary, or

tertiary alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylalkyl groups, or $-C(=O)R^{18}$ groups;

R^5 , R^6 , R^7 , and R^8 may be the same or different and are independently selected from H, Cl, Br, F, I, $-NO_2$, $-OH$, $-OR^{19}$ groups, $-NR^{20}R^{21}$ groups, $-SH$, $-SR^{22}$ groups, $-S(=O)R^{23}$ groups, $-S(=O)_2R^{24}$ groups, $-CN$, substituted or unsubstituted amidinyl groups, substituted or unsubstituted guanidinyl groups, substituted or unsubstituted primary, secondary, or tertiary alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylalkyl groups, $-C(=O)R^{25}$ groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocycliloxyalkyl groups;

R^9 and R^{14} may be the same or different and are independently selected from H, $-OH$, substituted or unsubstituted alkoxy groups, substituted or unsubstituted aryloxy groups, $-NH_2$, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, substituted or unsubstituted alkyl groups,

substituted or unsubstituted aryl groups, -C(=O)H, -C(=O)-alkyl groups, or -C(=O)-aryl groups;

R¹⁰ is selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted heterocyclylalkyl groups, -C(=O)H, 5 -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(alkyl)(aryl) 10 groups, -N(aryl)₂ groups, -C(=O)NH(heterocyclyl) groups, -C(=O)N(heterocyclyl)₂ groups, -C(=O)N(alkyl)(heterocyclyl) groups, or -C(=O)N(aryl)(heterocyclyl) groups;

R¹¹ is selected from H, -OH, alkoxy groups, aryloxy groups, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(aryl)₂ groups, 15 -N(alkyl)(aryl) groups, substituted or unsubstituted alkyl groups, -NH(heterocyclyl) groups, -N(heterocyclyl)₂ groups, -N(alkyl)(heterocyclyl) groups, or substituted or unsubstituted aryl groups;

R¹² is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl 20 groups;

R¹³ is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, -OH, alkoxy groups, aryloxy groups, -NH₂, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted 25 or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted

dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, 5 -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -C(=O)-heterocyclyl groups, -C(=O)-O-heterocyclyl groups, -C(=O)NH(heterocyclyl) groups, -C(=O)N(heterocyclyl)₂ groups, -C(=O)-N(alkyl)(heterocyclyl) groups, -C(=O)N(aryl)(heterocyclyl) groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, 10 substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocycloxyalkyl groups;

R¹⁵ and R¹⁹ may be the same or different and are independently selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, substituted or 15 unsubstituted heterocyclylalkyl groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, 20 substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl, substituted or unsubstituted diheterocyclylaminoalkyl, substituted or unsubstituted (heterocyclyl)(alkyl)aminoalkyl, substituted or unsubstituted 25 (heterocyclyl)(aryl)aminoalkyl, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocycloxyalkyl groups;

R¹⁶ and R²⁰ may be the same or different and are independently selected from H, substituted or unsubstituted alkyl groups, substituted or 30 unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups;

- R^{17} and R^{21} may be the same or different and are independently selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, $-C(=O)H$, $-C(=O)$ -alkyl groups, $-C(=O)$ -aryl groups, $-C(=O)NH_2$,
 5 $-C(=O)NH$ (alkyl) groups, $-C(=O)NH$ (aryl) groups, $-C(=O)N$ (alkyl)₂ groups, $-C(=O)N$ (aryl)₂ groups, $-C(=O)N$ (alkyl)(aryl) groups, $-C(=O)O$ -alkyl groups, $-C(=O)O$ -aryl groups, substituted or unsubstituted heterocyclalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups,
 10 substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, $-C(=O)$ -heterocyclyl groups, $-C(=O)-O$ -heterocyclyl groups, $-C(=O)NH$ (heterocyclyl) groups, $-C(=O)-N$ (heterocyclyl)₂ groups, $-C(=O)-N$ (alkyl)(heterocyclyl) groups, $-C(=O)-N$ (aryl)(heterocyclyl) groups,
 15 substituted or unsubstituted heterocyclalaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocycloxyalkyl groups;

- R^{18} , R^{23} , R^{24} , and R^{25} may be the same or different and are
 20 independently selected from H, $-NH_2$, $-NH$ (alkyl) groups, $-NH$ (aryl) groups, $-N$ (alkyl)₂ groups, $-N$ (aryl)₂ groups, $-N$ (alkyl)(aryl) groups, $-NH$ (heterocyclyl) groups, $-N$ (heterocyclyl)(alkyl) groups, $-N$ (heterocyclyl)(aryl) groups, $-N$ (heterocyclyl)₂ groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, $-OH$, substituted or unsubstituted alkoxy groups,
 25 substituted or unsubstituted aryloxy groups, substituted or unsubstituted heterocyclyl groups, $-NHOH$, $-N$ (alkyl)OH groups, $-N$ (aryl)OH groups, $-N$ (alkyl)O-alkyl groups, $-N$ (aryl)O-alkyl groups, $-N$ (alkyl)O-aryl groups, or $-N$ (aryl)O-aryl groups;
 and

R^{22} is selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups.

- In the fourth group of compounds, at least one of R^1 , R^2 , R^3 , or R^4 is an $-OR^{15}$ group and R^{15} is selected from substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted diheterocyclylaminoalkyl groups, substituted or unsubstituted (heterocyclyl)(alkyl)aminoalkyl groups, or substituted or unsubstituted (heterocyclyl)(aryl)aminoalkyl groups.

- In the first, second, or third group of compounds Z is preferably an $-NR^{14}$ group, more preferably where R^{14} is H. Preferred compounds of the fourth group include those compounds in which Z is an $-NR^{10}$ group, more preferably where R^{10} is H.

- Y is preferably an $-OR^{10}$ group, an $-NR^{12}R^{13}$ group, or a substituted or unsubstituted alkynyl group, or more preferably is an $-NR^{12}N^{13}$ group in the first, second, third, and fourth groups of compounds. In more preferred compounds of the first group, Y is a $NR^{12}R^{13}$ group and R^{12} is H. In more preferred compounds of the second and third groups, Y is a $NR^{12}R^{13}$ group and one or both of R^{12} and R^{13} are H.

- Other preferred compounds of the first, second, and third groups include those where Y is selected from $-N(CH_3)_2$, $-NH(CH_3)$, $-NH(CH_2CH_3)$, $-N(CH_2CH_3)_2$, $-NH(aryl)$ groups, $-N(aryl)_2$ groups, $-NHNH_2$, $-NHN(CH_3)_2$, $-N(CH_3)NH(CH_3)$, $-NH(CH_2)_mNH_2$ groups, $-NH(CH_2)_mNH(alkyl)$ groups, $-NH(CH_2)_mN(alkyl)_2$ groups, $-N(alkyl)(CH_2)_mNH_2$ groups,

- N(alkyl)(CH₂)_m-NH(alkyl) groups, -N(alkyl)(CH₂)_mN(alkyl)₂ groups,
 -NH(CH₂)_n(heterocyclyl) groups, -N(alkyl)[(CH₂)_n(heterocyclyl)] groups,
 -NH(CH₂)_mOH groups, -NH(CH₂)_mOCH₃ groups, -NHCH₂CH(NH₂)CH(CH₃)₂,
 -NH(2-aminocyclohexyl), -NH(cyclohexyl), -NH(OCH₃), -NH(N-morpholinyl),
 5 -NH(quinuclidyl), especially -NH(quinuclid-3-yl), or groups where R¹² and R¹³ join
 to form a substituted or unsubstituted saturated 5 or 6 membered N-containing ring,
 where m is an integer ranging from 2 to 4 such as 2, 3, or 4 and n is an integer
 ranging from 0 to 3 such as 0, 1, 2, or 3.

- More preferred compounds of the first, second, and third also
 10 include those in which Y is selected from -NH(5-benzimidazolyl),
 -NH(CH₂)₂N(CH₃)₂, -NH(CH₂)₂OH, -NH(CH₂)(4-imidazolyl), -NH(CH₂)(3-
 imidazolyl), -NH(CH₂)(4-pyridyl), -NH(CH₂)(2-pyridyl), -NH(CH₂)(3-pyridyl),
 -NH(CH₂)(2-tetrahydrofuranyl), -NH(CH₂)(4-piperidinyl), -NH(CH₂)(3-
 piperidinyl), -NH(CH₂)₂[2-(N-methylpyrrolidinyl)], -NH(CH₂)₂(2-pyrrolidinyl),
 15 -NH(CH₂)[2-(N-methylpyrrolidinyl)], -NH(CH₂)(2-pyrrolidinyl), -NH(3-
 piperidinyl), or -NH(3-pyrrolidinyl).

- Preferred compounds of the first and second groups include those
 compounds where R¹, R², R³, R⁴, R⁶, R⁷, and R⁸ are all H. Other preferred
 compounds of the first, second, third, and fourth groups include those where R¹ is
 20 selected from H, substituted or unsubstituted alkoxy groups, substituted or
 unsubstituted heterocyclalkoxy groups, substituted or unsubstituted
 heterocycloxy groups, substituted or unsubstituted heterocyclyl groups, substituted
 or unsubstituted alkyl-, heterocyclyl-, or aryl-aminoalkyl groups, substituted or
 unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted
 25 diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl
 groups, substituted or unsubstituted alkyl- or aryl-aminoalkoxy groups, substituted
 or unsubstituted dialkylaminoalkoxy groups, or substituted or unsubstituted
 diarylaminoalkoxy groups. Still other compounds of the first, second, third, and
 fourth groups include those in which R¹ is selected from F, Cl, substituted or
 30 unsubstituted alkoxy groups, substituted or unsubstituted heterocyclalkoxy groups,

- substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted alkyl-, heterocyclyl-, or aryl-aminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or
- 5 unsubstituted alkylaminoalkoxy groups, substituted or unsubstituted arylaminoalkoxy groups, substituted or unsubstituted dialkylaminoalkoxy groups, substituted or unsubstituted diarylaminoalkoxy groups, or substituted or unsubstituted (alkyl)(aryl)aminoalkoxy groups. Particular examples include:
- OCH₃, -OCH₂CH₂(N-morpholinyl), -N-morpholinyl, -N-cis-dialkylmorpholinyl,
- 10 -N-(4-alkyl)piperazinyl, -OCH₂CH₂N(alkyl)₂ groups, -OCH₂CH₂NH(alkyl) groups, -OCH₂CH₂NH₂, -OCH₂CH₂NH(aryl) groups, -OCH₂CH₂N(aryl)₂ groups, -OCH₂CH₂N(alkyl)(aryl) groups, alkoxy groups, -O(4-piperidinyl), -O[4-(1-alkyl)piperidinyl] groups, -O[3-(1-alkyl)piperidinyl] groups, -O[3-quinuclidinyl], -OCH₂(2-pyridyl), -OCH₂(4-pyridyl), -O(3-pyrrolidinyl), or -O[3-(1-
- 15 alkyl)pyrrolidinyl] groups.


- Still other preferred compounds of the first, second, third, and fourth groups include those in which R² is selected from F, Cl, -NO₂, -OCH₃, N-morpholinyl, -N-cis-dialkylmorpholinyl, -N-(4-alkyl)piperazinyl, or -OCH₂(2-pyridyl). Other preferred compounds of the first, second, third, and fourth groups
- 20 include those where R² is selected from H, F, Cl, -NO₂, substituted or unsubstituted heterocyclalkoxy groups, or substituted or unsubstituted heterocyclyl groups. Yet other preferred compounds of the first, second, third, and fourth groups include those where R² is selected from F, Cl, -NO₂, substituted or unsubstituted alkoxy groups, substituted or unsubstituted heterocyclalkoxy groups, substituted or
- 25 unsubstituted heterocyclyl groups, substituted or unsubstituted alkyl-, heterocycl-, and aryl-aminoalkyl groups, substituted or unsubstituted dialkyl- and diaryl-aminoalkyl groups, substituted or unsubstituted alkylaryl-aminoalkyl groups, substituted or unsubstituted alkyl- and aryl-aminoalkoxy groups, substituted or unsubstituted dialkyl- and diaryl-aminoalkoxy groups, or substituted or
- 30 unsubstituted alkylaryl-aminoalkoxy groups.

Still further preferred compounds of the first, second, third, and fourth groups include those where R^6 is an alkyl group having from one to four carbon atoms. In other preferred compounds of the first, second, third, and fourth groups, R^7 is an alkyl group having from one to four carbon atoms. Still further preferred compounds of the four groups are those in which R^6 or R^7 is an $-OR^{19}$ group and R^{19} is an alkyl group, an aryl group, a heterocyclyl group, or a heterocyclylalkyl group.


In still other preferred compounds of the first, second, third, and fourth groups, R^6 or R^7 is a $-OCH_2(CH_2)_q(\text{heterocyclyl})$ group and q is 0, 1, 2, 3, or 4, more preferably where the heterocyclyl group of the $-OCH_2(CH_2)_q(\text{heterocyclyl})$ group is a heterocycle selected from substituted or unsubstituted morpholine, substituted or unsubstituted piperazine, substituted or unsubstituted piperidine, substituted or unsubstituted pyrrolidine, substituted or unsubstituted pyrrole, substituted or unsubstituted imidazole, substituted or unsubstituted pyrazole, substituted or unsubstituted 1,2,3-triazole, substituted or unsubstituted 1,2,4-triazole, substituted or unsubstituted tetrazole, substituted or unsubstituted thiomorpholine, substituted or unsubstituted homopiperazine, substituted or unsubstituted oxazolidin-2-one, substituted or unsubstituted pyrrolidin-2-one, substituted or unsubstituted pyridine, substituted or unsubstituted oxazole, substituted or unsubstituted isoxazole, substituted or unsubstituted thiazole, substituted or unsubstituted isothiazole, substituted or unsubstituted furan, substituted or unsubstituted thiophene, substituted or unsubstituted tetrahydrofuran, substituted or unsubstituted tetrahydrothiophene, substituted or unsubstituted benzimidazole, substituted or unsubstituted benzoxazole, or substituted or unsubstituted benzothiazole.

In groups including heterocyclyl groups, the heterocyclyl may be attached in various ways. For example, in the $-OCH_2(CH_2)_q(\text{heterocyclyl})$ group, the heterocyclyl group may be bonded to a methylene carbon of the $-OCH_2(CH_2)_q$ group of the $-OCH_2(CH_2)_q(\text{heterocyclyl})$ through various ring members. By way of non-limiting example, where q is 1 and the heterocyclyl group is tetrahydrofuran,

the group could be represented by the formula $-\text{OCH}_2\text{CH}_2(\text{tetrahydrofuranyl})$ which corresponds to the following two structures:




III




IV


where structure III represents the group that can be referred to as the $-\text{OCH}_2\text{CH}_2(2\text{-tetrahydrofuranyl})$ group and structure IV represents the group that can be referred to as the $-\text{OCH}_2\text{CH}_2(3\text{-tetrahydrofuranyl})$ group. When the heterocyclyl group is a N-containing heterocycle, such as, but not limited to piperidine, piperazine, morpholine, or pyrrolidine, the heterocycle can be bonded to the methylene carbon through a ring carbon atom or through a nitrogen atom in the ring of the N-containing heterocycle. Both of these are preferred. Where the heterocyclyl group is a piperidine and q is 2 for an $-\text{OCH}_2(\text{CH}_2)_q(\text{heterocyclyl})$ group, the following structures are possible and preferred:




V



VI




VII




VIII

Structure V is an example of a $-\text{O}(\text{CH}_2)_3(\text{N-piperidinyl})$ or $-\text{O}(\text{CH}_2)_3(1\text{-piperidinyl})$ group. Structure VI is an example of a $-\text{O}(\text{CH}_2)_3(2\text{-piperidinyl})$ group. Structure VII is an example of a $-\text{O}(\text{CH}_2)_3(3\text{-piperidinyl})$

group. Structure VIII is an example of a $-\text{O}(\text{CH}_2)_3(4\text{-piperidiny})$ group. Where the heterocyclyl group is a piperazine and q is 1 for an $-\text{OCH}_2(\text{CH}_2)_q(\text{heterocyclyl})$ group, the following structures are possible and preferred:




IX




X


5 Structure IX is an example of a $-\text{O}(\text{CH}_2)_2(2\text{-piperaziny})$ group, and structure X is an example of a $-\text{O}(\text{CH}_2)_2(1\text{-piperaziny})$ or $-\text{O}(\text{CH}_2)_2(\text{N-piperaziny})$ group. Where the heterocyclyl group is a morpholine and q is 1 for an $-\text{OCH}_2(\text{CH}_2)_q(\text{heterocyclyl})$ group, the following structures are possible and preferred:



XI



XII



XIII

10

Structure XI is an example of a $-\text{O}(\text{CH}_2)_2(3\text{-morpholiny})$ group, structure XII is an example of a $-\text{O}(\text{CH}_2)_2(4\text{-morpholiny})$ or $-\text{O}(\text{CH}_2)_2(\text{N-morpholiny})$ group, and structure XIII is an example of a $-\text{O}(\text{CH}_2)_2(2\text{-morpholiny})$ group. It will be observed that where the group is a pyrrolidine, and q is 1, the structures available include $-\text{O}(\text{CH}_2)_2(1\text{-pyrrolidiny})$ or $-\text{O}(\text{CH}_2)_2(\text{N-pyrrolidiny})$, $-\text{O}(\text{CH}_2)_2(2\text{-pyrrolidiny})$, and $-\text{O}(\text{CH}_2)_2(3\text{-pyrrolidiny})$.

15

Yet other preferred compounds of the first, second, third, and fourth groups include those where at least one of R⁵, R⁶, R⁷, and R⁸ is a substituted or unsubstituted heterocyclyl group, more specifically a substituted or unsubstituted heterocyclyl group comprising at least one O or N atom, still more particularly a substituted or unsubstituted heterocyclyl group selected from morpholine, piperazine, piperidine, 1,2,3-triazole, 1,2,4-triazole, tetrazole, pyrrolidine, pyrazole, pyrrole, thiomorpholine, thiomorpholine in which the S atom of the thiomorpholine group is bonded to one or more O atoms, homopiperazine, benzimidazole, oxazolidin-2-one, pyrrolidin-2-one, imidazole, isoxazole, oxazole, isothiazole, thiazole, thiophene, furan, pyran, tetrahydrothiophene, tetrahydrofuran, tetrahydropyran, or pyridine.

Other preferred compounds of the first, second, third, and fourth groups include those in which at least one of R¹⁹ or R²¹ is selected from substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, (alkyl)(aryl)aminoalkyl groups, or substituted or unsubstituted heterocyclylalkyl groups, including: -CH₂(CH₂)_pNH₂ groups, -CH₂(CH₂)_pNH(alkyl) groups, -CH₂(CH₂)_pNH(aryl) groups, -CH₂(CH₂)_pN(alkyl)₂ groups, -CH₂(CH₂)_pN(aryl)₂ groups, -CH₂(CH₂)_pN(alkyl)(aryl) groups, or -CH₂(CH₂)_p(heterocyclyl) groups, wherein p is an integer ranging from 0 to 4 and the heterocyclyl group of the -CH₂(CH₂)_p(heterocyclyl) group is a N-containing heterocycle selected from substituted or unsubstituted morpholine, substituted or unsubstituted pyrrolidine, substituted or unsubstituted piperazine, substituted or unsubstituted piperidine, substituted or unsubstituted pyrrole, substituted or unsubstituted imidazole, substituted or unsubstituted pyrazole, substituted or unsubstituted 1,2,3-triazole, substituted or unsubstituted 1,2,4-triazole, substituted or unsubstituted tetrazole, substituted or unsubstituted thiomorpholine, substituted or unsubstituted homopiperazine, substituted or unsubstituted oxazolidin-2-one, substituted or unsubstituted pyrrolidin-2-one, substituted or unsubstituted pyridine, substituted or unsubstituted oxazole, substituted or unsubstituted isoxazole,

substituted or unsubstituted thiazole, substituted or unsubstituted isothiazole, substituted or unsubstituted benzimidazole, substituted or unsubstituted benzoxazole, or substituted or unsubstituted benzothiazole.

- Still further preferred compounds according to the first, second, third, and fourth groups include those in which R^{25} is selected from substituted or unsubstituted aryl groups, substituted or unsubstituted alkyl groups, $-NH_2$, $-NH(alkyl)$ groups, $-N(alkyl)_2$ groups, $-NH(aryl)$ groups, $-N(aryl)_2$ groups, $-N(alkyl)(aryl)$ groups, $-NH(heterocyclyl)$ groups, $-N(heterocyclyl)(alkyl)$ groups, $-N(heterocyclyl)(aryl)$ groups, $-N(heterocyclyl)_2$ groups, or N-containing heterocycles. In such compounds, the N-containing heterocycles are bonded to the carbonyl carbon of the $-C(=O)-R^{25}$ group through either a nitrogen atom or a carbon atom in the rings of the N-containing heterocycles. In more preferred such compounds that are provided, the N-containing heterocycle of the R^{25} group is selected from substituted or unsubstituted morpholine, substituted or unsubstituted pyrrolidine, substituted or unsubstituted piperazine, substituted or unsubstituted piperidine, substituted or unsubstituted pyrrole, substituted or unsubstituted imidazole, substituted or unsubstituted pyrazole, substituted or unsubstituted 1,2,3-triazole, substituted or unsubstituted 1,2,4-triazole, substituted or unsubstituted tetrazole, substituted or unsubstituted thiomorpholine, substituted or unsubstituted homopiperazine, substituted or unsubstituted oxazolidin-2-one, substituted or unsubstituted pyrrolidin-2-one, substituted or unsubstituted pyridine, substituted or unsubstituted oxazole, substituted or unsubstituted isoxazole, substituted or unsubstituted thiazole, substituted or unsubstituted isothiazole, substituted or unsubstituted benzimidazole, substituted or unsubstituted benzoxazole, or substituted or unsubstituted benzothiazole.

Preferred compounds according to the first, third, and fourth groups of compounds are also those where R^9 is H.


Preferred compounds of the fourth group of compounds include those where R^1 is an $-OR^{15}$ group and R^{15} is selected from substituted or unsubstituted

-77-

heterocyclalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted

- 5 (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocycl groups, substituted or unsubstituted heterocyclaminoalkyl groups, substituted or unsubstituted diheterocyclaminoalkyl groups, substituted or unsubstituted (heterocycl)(alkyl)aminoalkyl groups, or substituted or unsubstituted (heterocycl)(aryl)aminoalkyl groups.

- 10 Other particularly preferred inhibitors of VEGF-RTK are compounds having the structure II, tautomers of the compounds, pharmaceutically acceptable salts of the compounds, and pharmaceutically acceptable salts of the tautomers. Structure II has the following formula:



II

- 15 In compounds having structure II, Y is selected from H, -OH, -OR¹⁰ groups, -SH, -SR¹¹ groups, -NR¹²R¹³ groups, -CN, -C(=O)-R¹⁴ groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted aralkyl groups, substituted or unsubstituted heterocyclalkyl groups, substituted or
- 20 unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or

unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or
 5 unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocycloxyalkyl groups.

In preferred compounds of structure II, Y is selected from H, -OH, -OR⁹ groups, or -NR¹¹R¹² groups. More preferably, Y is a -NR¹¹R¹² group. Still more preferably, Y is a -NR¹¹R¹² group and both R¹¹ and R¹² are hydrogen. In
 10 other preferred compounds having the structure II, Y is selected from -N(CH₃)₂, -NH(CH₃), -NH(CH₂CH₃), -N(CH₂CH₃)₂, -NH(aryl) groups, -N(aryl)₂ groups, -NHNH₂, -NHN(CH₃)₂, -N(CH₃)NH(CH₃), -NH(CH₂)_mNH₂ groups, -NH(CH₂)_mNH(alkyl) groups, -NH(CH₂)_mN(alkyl)₂ groups, -N(alkyl)(CH₂)_mNH₂ groups, -N(alkyl)(CH₂)_mNH(alkyl) groups, -N(alkyl)(CH₂)_mN(alkyl)₂ groups,
 15 -NH(CH₂)_n(heterocyclyl) groups, -N(alkyl)[(CH₂)_n(heterocyclyl)] groups, -NH(CH₂)_mOH groups, -NH(CH₂)_mOCH₃ groups, -NHCH₂CH(NH₂)CH(CH₃)₂, -NH(2-aminocyclohexyl), -NH(cyclohexyl), -NHOCH₃, -NH(N-morpholinyl), -NH(quinuclidyl), especially -NH(quinuclid-3-yl), and groups where R¹¹ and R¹² join to form a substituted or unsubstituted saturated 5 or 6 membered N-containing
 20 ring, where m is 2, 3, or 4 and n is 0, 1, 2, or 3. Still more preferred compounds of this type are those in which Y is selected from -NH(5-benzimidazolyl), -NH(CH₂)₂N(CH₃)₂, -NH(CH₂)₂OH, -NH(CH₂)(4-imidazolyl), -NH(CH₂)(3-imidazolyl), -NH(CH₂)(4-pyridyl), -NH(CH₂)(2-pyridyl), -NH(CH₂)(3-pyridyl), -NH(CH₂)(2-tetrahydrofuranyl), -NH(CH₂)(4-piperidinyl), -NH(CH₂)(3-
 25 piperidinyl), -NH(CH₂)₂[2-(N-methyl-pyrrolidinyl)], -NH(CH₂)₂(2-pyrrolidinyl), -NH(CH₂)₂[2-(N-methylpyrrolidinyl)], -NH(CH₂)(2-pyrrolidinyl), -NH(3-piperidinyl), or -NH(3-pyrrolidinyl).

In compounds of structure II, X¹, X², X³, and X⁴ are selected from C or N and at least one of X¹, X², X³, and X⁴ is N. In some preferred compounds of
 30 structure II, X¹ is N, R⁵ is absent or H, and X², X³, and X⁴ are all C. In other

preferred compounds of structure II, X^2 is N, R^6 is absent or H, and X^1 , X^3 , and X^4 are C. In other preferred compounds of structure II, X^3 is N, R^7 is absent or H, and X^1 , X^2 , and X^4 are all C. In still other preferred compounds of structure II, X^4 is N, R^8 is absent or H, and X^1 , X^2 , and X^3 are all C.

- 5 In compounds of structure II, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 may be the same or different and are independently selected from H, Cl, Br, F, I, $-\text{NO}_2$, $-\text{CN}$, $-\text{OH}$, $-\text{OR}^{15}$ groups, $-\text{NR}^{16}\text{R}^{17}$ groups, $-\text{C}(=\text{O})\text{R}^{18}$ groups, $-\text{SH}$, $-\text{SR}^{19}$ groups, $-\text{S}(=\text{O})\text{R}^{20}$ groups, $\text{S}(=\text{O})_2\text{R}^{21}$ groups, substituted or unsubstituted amidinyl groups, substituted or unsubstituted guanidinyl groups, substituted or unsubstituted primary, secondary, or tertiary alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted alkenyl groups, substituted or unsubstituted alkynyl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocycliloxyalkyl groups. In compounds of structure II, R^5 is absent or is H if X^1 is N, R^6 is absent or is H if X^2 is N, R^7 is absent or is H if X^3 is N, and R^8 is absent or is H if X^4 is N.

- Some preferred compounds have the structure II where at least one of R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , or R^8 is a substituted or unsubstituted heterocyclyl group, and, in more particular embodiments, a substituted or unsubstituted heterocyclyl group selected from morpholine, piperazine, piperidine, 1,2,3-triazole, 1,2,4-triazole, tetrazole, pyrrolidine, pyrazole, pyrrole, thiomorpholine, homopiperazine, benzimidazole, oxazolidin-2-one, pyrrolidin-2-one, imidazole, isoxazole, oxazole, isothiazole, thiazole, thiophene, furan, pyran, tetrahydrothiophene, tetrahydrofuran, tetrahydropyran, and pyridine.

Still other preferred compounds having structure II are those in which R^1 , R^2 , R^3 , and R^4 are H. In some preferred embodiments, R^1 is selected from H, substituted or unsubstituted alkoxy groups, substituted or unsubstituted heterocyclalkoxy groups, substituted or unsubstituted heterocycloxy groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted alkyl-, heterocyclyl-, or aryl-aminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted alkyl- or aryl-aminoalkoxy groups, substituted or unsubstituted dialkylaminoalkoxy groups, or substituted or unsubstituted diarylaminoalkoxy groups. In other preferred compounds, R^1 is selected from F, Cl, substituted or unsubstituted alkoxy groups, substituted or unsubstituted heterocyclalkoxy groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted alkyl-, heterocyclyl-, or aryl-aminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted alkylaminoalkoxy groups, substituted or unsubstituted arylaminoalkoxy groups, substituted or unsubstituted dialkylaminoalkoxy groups, substituted or unsubstituted diarylaminoalkoxy groups, or substituted or unsubstituted (alkyl)(aryl)aminoalkoxy groups. Particular examples include: -OCH₃, -OCH₂CH₂(N-morpholinyl), -N-morpholinyl, -N-cis-dimethylmorpholinyl, -N-(4-alkyl)piperazinyl, -OCH₂CH₂N(alkyl)₂ groups, -OCH₂CH₂NH(alkyl) groups, -OCH₂CH₂NH₂, -OCH₂CH₂NH(aryl) groups, -OCH₂CH₂N(aryl)₂ groups, alkoxy groups, -OCH₂CH₂N(alkyl)(aryl) groups, -O(4-piperidinyl), -O[4-(1-alkyl)piperidinyl] groups, -O[3-(1-alkyl)piperidinyl] groups, -O[3-quinuclidinyl], -OCH₂(2-pyridyl), -OCH₂(4-pyridyl), -O(3-pyrrolidinyl), -O[3-(1-alkyl)pyrrolidinyl], or other -O(heterocyclyl) groups not listed in this paragraph.

In other compounds having the structure II, R^2 is selected from F, Cl, -NO₂, -OCH₃, -N-morpholinyl, -N-cis-dialkylmorpholinyl, -N-(4-alkyl)piperazinyl, or -OCH₂(2-pyridyl). In yet other compounds having the structure II, R^2 is selected from H, F, Cl, -NO₂, substituted or unsubstituted

heterocyclalkoxy groups, or substituted or unsubstituted heterocycl groups. Yet other preferred compounds having the structure II include those where R² is selected from F, Cl, -NO₂, substituted or unsubstituted alkoxy groups, substituted or unsubstituted heterocyclalkoxy groups, substituted or unsubstituted heterocycl groups, substituted or unsubstituted alkyl-, heterocycl-, and aryl-aminoalkyl groups, substituted or unsubstituted dialkyl- and diaryl-aminoalkyl groups, substituted or unsubstituted alkylaryl-aminoalkyl groups, substituted or unsubstituted alkyl- and aryl-aminoalkoxy groups, substituted or unsubstituted dialkyl- and diaryl-aminoalkoxy groups, or substituted or unsubstituted alkylaryl-aminoalkoxy groups.

10 In some preferred compounds having structure II, at least two of X¹, X², X³, and X⁴ are C and the corresponding substituents R⁵, R⁶, R⁷, and R⁸ are hydrogen, and at least one of X¹, X², X³, and X⁴ is N. In yet other preferred compounds having structure II, three of R⁵, R⁶, R⁷, and R⁸ are hydrogen and one of X¹, X², X³, and X⁴ is N. In still other more preferred compounds of structure II, 15 R⁶, R⁷, or both R⁶ and R⁷ are alkyl groups such as those having from one to four carbon atoms. In yet other preferred compounds of structure II, R⁶ or R⁷ is an -OR¹⁴ group and R¹⁴ is an alkyl, aryl, heterocycl, or heterocyclalkyl group. In still further compounds of structure II, R⁶ or R⁷ is a -OCH₂(CH₂)_q(heterocycl) group and q is 0, 1, 2, 3, or 4. In more preferred compounds in which R⁶ or R⁷ is 20 a -OCH₂(CH₂)_q(heterocycl) group, the heterocycl group of the -OCH₂(CH₂)_n(heterocycl) group is a heterocycle selected from substituted or unsubstituted morpholine, substituted or unsubstituted piperazine, substituted or unsubstituted piperidine, substituted or unsubstituted pyrrolidine, substituted or unsubstituted pyrrole, substituted or unsubstituted imidazole, substituted or 25 unsubstituted pyrazole, substituted or unsubstituted 1,2,3-triazole, substituted or unsubstituted 1,2,4-triazole, substituted or unsubstituted tetrazole, substituted or unsubstituted thiomorpholine, substituted or unsubstituted thiomorpholine in which the S atom of the thiomorpholine group is bonded to one or more O atoms, substituted or unsubstituted homopiperazine, substituted or unsubstituted oxazolidin- 30 2-one, substituted or unsubstituted pyrrolidin-2-one, substituted or unsubstituted pyridine, substituted or unsubstituted oxazole, substituted or unsubstituted

- isoxazole, substituted or unsubstituted thiazole, substituted or unsubstituted isothiazole, substituted or unsubstituted furan, substituted or unsubstituted thiophene, substituted or unsubstituted tetrahydrofuran, substituted or unsubstituted tetrahydrothiophene, substituted or unsubstituted benzimidazole, substituted or
- 5 unsubstituted benzoxazole, or substituted or unsubstituted benzothiazole.

- In compounds of structure II, R^9 is selected from H, -OH, substituted or unsubstituted alkoxy groups, substituted or unsubstituted aryloxy groups, -NH₂, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or
- 10 unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, -C(=O)H, -C(=O)-alkyl groups, or -C(=O)-aryl groups. One group of particularly preferred compounds of structure II are those in which R^9 is hydrogen.

- In compounds of structure II, R^{10} is selected from substituted or
- 15 unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, substituted or unsubstituted heterocyclylalkyl groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups,
- 20 -C(=O)N(alkyl)(aryl) groups, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(alkyl)(aryl) groups, -N(aryl)₂ groups, -C(=O)NH(heterocyclyl) groups, -C(=O)N(heterocyclyl)₂ groups, -C(=O)N(alkyl)(heterocyclyl) groups, or -C(=O)N(aryl)(heterocyclyl) groups;

- In compounds of structure II, R^{11} and R^{19} may be the same or
- 25 different and are independently selected from substituted or unsubstituted alkyl groups, or substituted or unsubstituted aryl groups whereas R^{12} is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups.

In compounds of structure II, R^{13} is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, -OH, alkoxy groups, aryloxy groups, -NH₂, substituted or unsubstituted heterocyclylalkyl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted alkylamino groups, substituted or unsubstituted arylamino groups, substituted or unsubstituted dialkylamino groups, substituted or unsubstituted diarylamino groups, substituted or unsubstituted (alkyl)(aryl)amino groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -C(=O)-heterocyclyl groups, -C(=O)-O-heterocyclyl groups, -C(=O)NH(heterocyclyl) groups, -C(=O)-N(heterocyclyl)₂ groups, -C(=O)N(alkyl)(heterocyclyl) groups, -C(=O)-N(aryl)(heterocyclyl) groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, or substituted or unsubstituted heterocyclioxyalkyl groups. R^{12} and R^{13} may join together to form a 5 to 7 membered saturated or unsaturated, substituted or unsubstituted N-containing ring.

In compounds of structure II, R^{14} is selected from H, -OH, alkoxy groups, aryloxy groups, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, -NH(heterocyclyl) groups, -N(heterocyclyl)₂ groups, -N(alkyl)(heterocyclyl) groups, or -N(aryl)(heterocyclyl) groups.

In compounds of structure II, R^{15} is selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or

- unsubstituted heterocyclyl groups, substituted or unsubstituted heterocyclylalkyl groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, substituted or unsubstituted
- 5 aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, substituted or unsubstituted heterocyclylaminoalkyl groups, substituted or unsubstituted
- 10 diheterocyclylaminoalkyl groups, substituted or unsubstituted (heterocyclyl)(alkyl)aminoalkyl groups, substituted or unsubstituted (heterocyclyl)(aryl)aminoalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, substituted or unsubstituted hydroxyalkyl groups, or substituted or unsubstituted
- 15 heterocyclyloxyalkyl groups.

- In compounds of structure II, R¹⁶ is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, or substituted or unsubstituted heterocyclyl groups whereas R¹⁷ is selected from H, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, substituted or unsubstituted heterocyclyl groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl
- 20 groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups, substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted
- 25 arylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (aryl)(alkyl)aminoalkyl groups, substituted or unsubstituted heterocyclylalkyl groups, -C(=O)-heterocyclyl groups, -C(=O)-Oheterocyclyl groups, -C(=O)NH(heterocyclyl) groups, -C(=O)N(heterocyclyl)₂ groups, -C(=O)-N(alkyl)(heterocyclyl) groups, -C(=O)N(aryl)(heterocyclyl) groups, substituted or unsubstituted
- 30

heterocyclylaminoalkyl groups, substituted or unsubstituted hydroxyalkyl groups, substituted or unsubstituted alkoxyalkyl groups, substituted or unsubstituted aryloxyalkyl groups, substituted or unsubstituted heterocyclyloxyalkyl groups, -OH, substituted or unsubstituted alkoxy groups, substituted or unsubstituted aryloxy groups, or -NH₂ groups. R¹⁶ and R¹⁷ may join together to form a 5 to 7 membered saturated or unsaturated, substituted or unsubstituted N-containing ring.

Finally, in compounds of structure II, R¹⁸, R²⁰, and R²¹ may be the same or different and are independently selected from H, -NH₂, -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups, substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, -OH, substituted or unsubstituted alkoxy groups, substituted or unsubstituted aryloxy groups, substituted or unsubstituted heterocyclyl groups, -NHOH, -N(alkyl)OH groups, -N(aryl)OH groups, -N(alkyl)O-alkyl groups, -N(aryl)O-alkyl groups, -N(alkyl)O-aryl groups, or -N(aryl)O-aryl groups.

Compounds having the structure II may include those in which R¹⁸ is selected from substituted or unsubstituted alkyl groups, substituted or unsubstituted aryl groups, -NH₂, -NH(alkyl) groups, -N(alkyl)₂ groups, -NH(aryl) groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups, -NH(heterocyclyl) groups, -N(heterocyclyl)(alkyl) groups, -N(heterocyclyl)(aryl) groups, -N(heterocyclyl)₂ groups, or N-containing heterocycles, and the N-containing heterocycles are bonded to the carbonyl carbon of the -C(=O)-R¹⁸ group through either a nitrogen atom or a carbon atom in the rings of the N-containing heterocycles. In still more preferred compounds in which R¹⁸ is a N-containing heterocycle, the N-containing heterocycle of the R¹⁸ group is selected from substituted or unsubstituted morpholine, substituted or unsubstituted pyrrolidine, substituted or unsubstituted piperazine, substituted or unsubstituted piperidine, substituted or unsubstituted pyrrole, substituted or unsubstituted imidazole, substituted or unsubstituted pyrazole, substituted or unsubstituted 1,2,3-triazole, substituted or unsubstituted 1,2,4-triazole, substituted or unsubstituted tetrazole, substituted or unsubstituted thiomorpholine, substituted or unsubstituted homopiperazine, substituted or


unsubstituted oxazolidin-2-one, substituted or unsubstituted pyrrolidin-2-one, substituted or unsubstituted pyridine, substituted or unsubstituted oxazole, substituted or unsubstituted isoxazole, substituted or unsubstituted thiazole, substituted or unsubstituted isothiazole, substituted or unsubstituted benzimidazole, substituted or unsubstituted benzoxazole, or substituted or unsubstituted benzothiazole.

Other preferred compounds having structure II are provided in which R^{15} or R^{17} is selected from substituted or unsubstituted aminoalkyl groups, substituted or unsubstituted alkylaminoalkyl groups, substituted or unsubstituted arylaminoalkyl groups, substituted or unsubstituted dialkylaminoalkyl groups, substituted or unsubstituted diarylaminoalkyl groups, substituted or unsubstituted (alkyl)(aryl)aminoalkyl groups, or substituted or unsubstituted heterocyclylaminoalkyl groups including: $-\text{CH}_2(\text{CH}_2)_p\text{NH}_2$ groups, $-\text{CH}_2(\text{CH}_2)_p\text{NH}(\text{alkyl})$ groups, $-\text{CH}_2(\text{CH}_2)_p\text{NH}(\text{aryl})$ groups, $-\text{CH}_2(\text{CH}_2)_p\text{N}(\text{alkyl})_2$ groups, $-\text{CH}_2(\text{CH}_2)_p\text{N}(\text{aryl})_2$ groups, $-\text{CH}_2(\text{CH}_2)_p\text{N}(\text{alkyl})(\text{aryl})$ groups, or $-\text{CH}_2(\text{CH}_2)_p(\text{heterocyclyl})$ groups, where p is an integer ranging from 0 to 4 and the heterocyclyl group of the $-\text{CH}_2(\text{CH}_2)_p(\text{heterocyclyl})$ group is a N-containing heterocycle selected from substituted or unsubstituted morpholine, substituted or unsubstituted pyrrolidine, substituted or unsubstituted piperazine, substituted or unsubstituted piperidine, substituted or unsubstituted pyrrole, substituted or unsubstituted imidazole, substituted or unsubstituted pyrazole, substituted or unsubstituted 1,2,3-triazole, substituted or unsubstituted 1,2,4-triazole, substituted or unsubstituted tetrazole, substituted or unsubstituted thiomorpholine, substituted or unsubstituted homopiperazine, substituted or unsubstituted oxazolidin-2-one, substituted or unsubstituted pyrrolidin-2-one, substituted or unsubstituted pyridine, substituted or unsubstituted oxazole, substituted or unsubstituted isoxazole, substituted or unsubstituted thiazole, substituted or unsubstituted isothiazole, substituted or unsubstituted benzimidazole, substituted or unsubstituted benzoxazole, or substituted or unsubstituted benzothiazole.

Compounds of structure I are readily synthesized from simple starting molecules as shown in the following Examples. Compounds of structure I may generally be prepared using benzene substituted with nitrile or carboxylic acid groups in addition to other optional groups.

- 5 Compounds of structure I may be synthesized from simple starting molecules as shown in Schemes 1-4 and exemplified in the Examples. As shown in Scheme 1, compounds of structure I may generally be prepared using aromatic compounds substituted with amines and carboxylic acid groups.


Scheme 1.




- 10 As shown in Scheme 1, a substituted aromatic compound such as a substituted or unsubstituted 2-aminobenzoic acid may be reacted with an acyl halide such as methyl 2-(chlorocarbonyl)acetate to produce an amide that will react with a substituted or unsubstituted 1,2-diaminobenzene. The resulting product is a 4-hydroxy-substituted compound of structure I. One skilled in the art will recognize
- 15 that the procedure set forth in Scheme 1 may be modified to produce various compounds.

- A method for preparing 4-amino substituted compounds of structure I is shown in Scheme 2. As shown in Scheme 2, aromatic compounds substituted with amine and nitrile groups may be used to synthesize 4-amino substituted
- 20 compounds of structure I. A compound such as ethyl 2-cyanoacetate may be


- reacted with ethanol to produce ethyl 3-ethoxy-3-iminopropanoate hydrochloride. Subsequent reaction with a substituted or unsubstituted 1,2-phenylenediamine provides substituted or unsubstituted ethyl 2-benzimidazol-2-ylacetate. Reaction of a substituted or unsubstituted ethyl 2-benzimidazol-2-ylacetate with an aromatic compound having an amine and nitrile group such as substituted or unsubstituted 2-aminobenzonitrile with a base such as lithium bis(trimethylsilyl)amide or a Lewis acid such as tin tetrachloride provides the substituted or unsubstituted 4-amino substituted compound of structure I.

Scheme 2.

- Scheme 3 illustrates a general synthetic route that allows for the synthesis of 4-dialkylamino and 4-alkylamino compounds of structure I. An inspection of Scheme 3 shows that 4-hydroxy substituted compounds of structure I may be converted into the 4-chloro derivative by reaction with phosphorous oxychloride or thionyl chloride. The 4-chloro derivative may then be reacted with an alkylamine or dialkylamine to produce the corresponding 4-alkylamino or 4-dialkylamino derivative. Deprotection affords the final 4-alkylamino or 4-dialkylamino compounds of structure I. Other groups that may be reacted with the 4-chloro derivative in this manner include, but are not limited to, ROH, RSH, and CuCN.


Scheme 3.

As shown in Scheme 4, the synthesis of compounds of structure I having a H, alkyl group, aryl group, or heterocyclyl group in the 4-position may be accomplished using a substituted or unsubstituted 2-benzimidazol-2-ylacetate prepared as shown in Schemes 2 and 3.

Scheme 4.

Heteroaromatic diamines may be used as precursors of compounds of structure II. The synthesis of compounds of structure II where Y = NH₂ is depicted in Scheme 5.

Scheme 5.



A compound such as ethyl cyanoacetate may be condensed with a substituted or unsubstituted heterocycle containing two ortho amino groups such as substituted or unsubstituted 1,2-diaminopyridine to obtain a substituted or unsubstituted 2-imidazolo[5,4-b]pyridin-2-ylacetate, which may subsequently be hydrolyzed in acidic medium to provide a substituted or unsubstituted ethyl 2-imidazolo[5,4-b]pyridin-2-ylacetate. As an alternate route, a substituted or unsubstituted ethyl 2-imidazolo[5,4-b]pyridin-2-ylacetate may be obtained from a compound such as the hydrochloride salt of 3-ethoxy-3-iminopropanoate and a substituted or unsubstituted 1,2-diaminopyridine. Reaction of a substituted or unsubstituted ethyl 2-imidazolo[5,4-b]pyridin-2-ylacetates with an aromatic compound having an amine and nitrile group such as substituted or unsubstituted 2-aminobenzonitrile with a base such as lithium bis(trimethylsilyl)amide provides the substituted or unsubstituted compound of structure II.

The instant invention also provides for compositions which may be prepared by mixing one or more compounds of the instant invention, or pharmaceutically acceptable salts or tautomers thereof, with pharmaceutically acceptable carriers, excipients, binders, diluents or the like, to treat or ameliorate a variety of disorders related to the activity of VEGF-RTK, more particularly angiogenesis associated with cancer. A therapeutically effective dose further refers to that amount of one or more compounds of the instant invention sufficient to result

in amelioration of symptoms of the disorder. The pharmaceutical compositions of the instant invention can be manufactured by methods well known in the art such as conventional granulating, mixing, dissolving, encapsulating, lyophilizing, emulsifying or levigating processes, among others. The compositions can be in the form of, for example, granules, powders, tablets, capsules, syrup, suppositories, injections, emulsions, elixirs, suspensions or solutions. The instant compositions can be formulated for various routes of administration, for example, by oral administration, by transmucosal administration, by rectal administration, or subcutaneous administration as well as intrathecal, intravenous, intramuscular, intraperitoneal, intranasal, intraocular or intraventricular injection. The compound or compounds of the instant invention can also be administered in a local rather than a systemic fashion, such as injection as a sustained release formulation. The following dosage forms are given by way of example and should not be construed as limiting the instant invention.

For oral, buccal, and sublingual administration, powders, suspensions, granules, tablets, pills, capsules, gelcaps, and caplets are acceptable as solid dosage forms. These can be prepared, for example, by mixing one or more compounds of the instant invention, or pharmaceutically acceptable salts or tautomers thereof, with at least one additive or excipient such as a starch or other additive. Suitable additives or excipients are sucrose, lactose, cellulose sugar, mannitol, maltitol, dextran, sorbitol, starch, agar, alginates, chitins, chitosans, pectins, tragacanth gum, gum arabic, gelatins, collagens, casein, albumin, synthetic or semi-synthetic polymers or glycerides, methyl cellulose, hydroxypropylmethyl-cellulose, and/or polyvinylpyrrolidone. Optionally, oral dosage forms can contain other ingredients to aid in administration, such as an inactive diluent, or lubricants such as magnesium stearate, or preservatives such as paraben or sorbic acid, or anti-oxidants such as ascorbic acid, tocopherol or cysteine, a disintegrating agent, binders, thickeners, buffers, sweeteners, flavoring agents or perfuming agents. Additionally, dyestuffs or pigments may be added for identification. Tablets and pills may be further treated with suitable coating materials known in the art.

Liquid dosage forms for oral administration may be in the form of pharmaceutically acceptable emulsions, syrups, elixirs, suspensions, slurries and solutions, which may contain an inactive diluent, such as water. Pharmaceutical formulations may be prepared as liquid suspensions or solutions using a sterile
5 liquid, such as, but not limited to, an oil, water, an alcohol, and combinations of these. Pharmaceutically suitable surfactants, suspending agents, emulsifying agents, may be added for oral or parenteral administration.

As noted above, suspensions may include oils. Such oil include, but are not limited to, peanut oil, sesame oil, cottonseed oil, corn oil and olive oil.
10 Suspension preparation may also contain esters of fatty acids such as ethyl oleate, isopropyl myristate, fatty acid glycerides and acetylated fatty acid glycerides. Suspension formulations may include alcohols, such as, but not limited to, ethanol, isopropyl alcohol, hexadecyl alcohol, glycerol and propylene glycol. Ethers, such as but not limited to, poly(ethyleneglycol), petroleum hydrocarbons such as mineral
15 oil and petrolatum; and water may also be used in suspension formulations.

For nasal administration, the pharmaceutical formulations may be a spray or aerosol containing and appropriate solvents and optionally other compounds such as, but not limited to, stabilizers, antimicrobial agents, antioxidants, pH modifiers, surfactants, bioavailability modifiers and combinations
20 of these. A propellant for an aerosol formulation may include compressed air, nitrogen, carbon dioxide, or a hydrocarbon based low boiling solvent. The compound or compounds of the instant invention are conveniently delivered in the form of an aerosol spray presentation from a nebulizer or the like.

Injectable dosage forms generally include aqueous suspensions or oil
25 suspensions which may be prepared using a suitable dispersant or wetting agent and a suspending agent. Injectable forms may be in solution phase or in the form of a suspension, which is prepared with a solvent or diluent. Acceptable solvents or vehicles include sterilized water, Ringer's solution, or an isotonic aqueous saline solution. Alternatively, sterile oils may be employed as solvents or suspending

agents. Preferably, the oil or fatty acid is non-volatile, including natural or synthetic oils, fatty acids, mono-, di- or tri-glycerides.

For injection, the pharmaceutical formulation may be a powder suitable for reconstitution with an appropriate solution as described above.

- 5 Examples of these include, but are not limited to, freeze dried, rotary dried or spray dried powders, amorphous powders, granules, precipitates, or particulates. For injection, the formulations may optionally contain stabilizers, pH modifiers, surfactants, bioavailability modifiers and combinations of these. The compounds may be formulated for parenteral administration by injection such as by bolus
10 injection or continuous infusion. A unit dosage form for injection may be in ampoules or in multi-dose containers.

- For rectal administration, the pharmaceutical formulations may be in the form of a suppository, an ointment, an enema, a tablet or a cream for release of compound in the intestines, sigmoid flexure and/or rectum. Rectal suppositories are
15 prepared by mixing one or more compounds of the instant invention, or pharmaceutically acceptable salts or tautomers of the compound, with acceptable vehicles, for example, cocoa butter or polyethylene glycol, which is present in a solid phase at normal storing temperatures, and present in a liquid phase at those temperatures suitable to release a drug inside the body, such as in the rectum. Oils
20 may also be employed in the preparation of formulations of the soft gelatin type and suppositories. Water, saline, aqueous dextrose and related sugar solutions, and glycerols may be employed in the preparation of suspension formulations which may also contain suspending agents such as pectins, carbomers, methyl cellulose, hydroxypropyl cellulose or carboxymethyl cellulose, as well as buffers and
25 preservatives.

Besides those representative dosage forms described above, pharmaceutically acceptable excipients and carries are generally known to those skilled in the art and are thus included in the instant invention. Such excipients and

carriers are described, for example, in "Remingtons Pharmaceutical Sciences" Mack Pub. Co., New Jersey (1991), which is incorporated herein by reference.

The formulations of the invention may be designed for to be short-acting, fast-releasing, long-acting, and sustained-releasing as described below.

- 5 Thus, the pharmaceutical formulations may also be formulated for controlled release or for slow release.

The instant compositions may also comprise, for example, micelles or liposomes, or some other encapsulated form, or may be administered in an extended release form to provide a prolonged storage and/or delivery effect.

- 10 Therefore, the pharmaceutical formulations may be compressed into pellets or cylinders and implanted intramuscularly or subcutaneously as depot injections or as implants such as stents. Such implants may employ known inert materials such as silicones and biodegradable polymers.

- Specific dosages may be adjusted depending on conditions of disease,
15 the age, body weight, general health conditions, sex, and diet of the subject, dose intervals, administration routes, excretion rate, and combinations of drugs. Any of the above dosage forms containing effective amounts are well within the bounds of routine experimentation and therefore, well within the scope of the instant invention.

- 20 A therapeutically effective dose may vary depending upon the route of administration and dosage form. The preferred compound or compounds of the instant invention is a formulation that exhibits a high therapeutic index. The therapeutic index is the dose ratio between toxic and therapeutic effects which can be expressed as the ratio between LD_{50} and ED_{50} . The LD_{50} is the dose lethal to
25 50% of the population and the ED_{50} is the dose therapeutically effective in 50% of the population. The LD_{50} and ED_{50} are determined by standard pharmaceutical procedures in animal cell cultures or experimental animals.

“Treating” within the context of the instant invention, means an alleviation of symptoms associated with a disorder or disease, or halt of further progression or worsening of those symptoms, or prevention or prophylaxis of the disease or disorder. For example, within the context of treating patients in need of an inhibitor of VEGF-RTK, successful treatment may include a reduction in the proliferation of capillaries feeding a tumor or diseased tissue, an alleviation of symptoms related to a cancerous growth or tumor, proliferation of capillaries, or diseased tissue, a halting in capillary proliferation, or a halting in the progression of a disease such as cancer or in the growth of cancerous cells. Treatment may also include administering the pharmaceutical formulations of the present invention in combination with other therapies. For example, the compounds and pharmaceutical formulations of the present invention may be administered before, during, or after surgical procedure and/or radiation therapy. The compounds of the invention can also be administered in conjunction with other anti-cancer drugs including those used in antisense and gene therapy.

A method of treating a patient in need of an inhibitor of vascular endothelial growth factor receptor tyrosine kinase includes administering an effective amount of a pharmaceutical formulation according to the invention to a patient in need thereof.

A method for inhibiting tumor growth in a patient includes administering an effective amount of the compound or a pharmaceutically acceptable salt thereof to a patient having a tumor.

A method for inhibiting the proliferation of capillaries in a patient includes administering an effective amount of the compound or a pharmaceutically acceptable salt thereof according to a patient in need.

A method of preparing pharmaceutical formulations includes mixing any of the above-described compounds with a pharmaceutically acceptable carrier and water or an aqueous solution.

The present invention, thus generally described, will be understood more readily by reference to the following examples, which are provided by way of illustration and are not intended to be limiting of the present invention.

EXAMPLES

5 The following abbreviations are used in the Examples:

	ATP:	Adenosine triphosphate
	BSA:	Bovine Serum Albumin
	DMA:	<i>N,N</i> -Dimethylacetamide
	DMF:	<i>N,N</i> -Dimethylformamide
10	dppf:	1,1'-(diphenylphosphino)ferrocene
	DTT:	DL-Dithiothreitol
	EDTA:	Ethylene diamine tetraacetic acid
	EtOAc:	Ethyl acetate
	EtOH:	Ethanol
15	HBTU:	O-Benzotriazol-1-yl- <i>N,N,N',N'</i> -tetramethyluronium hexafluorophosphate
	IC ₅₀ value:	The concentration of an inhibitor that causes a 50 % reduction in a measured activity.
	LiHMDS:	Lithium bis(trimethylsilyl)amide
20	MeOH:	Methanol
	NMP:	<i>N</i> -methylpyrrolidone
	THF:	Tetrahydrofuran

 The compounds were named using Nomenclator (v. 3.0 & v. 5.0)
25 from CmemInnovation Software, Inc. and ACD/Name v. 4.53.

 The various aryl diamine starting materials used to synthesize
benzimidazole acetates may be obtained from commercial sources, prepared by
methods know to one of skill in the art, or prepared by the following general
Methods 1-15.

-97-


Method 1

- 2,4-Difluoronitrobenzene (1.0 eq) was placed in a dry round-bottomed flask equipped with a dry ice condenser charged with acetone and dry ice.
- 5 Ammonia was condensed into the flask and the resulting solution was stirred at reflux for 7 hours. A yellow precipitate formed within 1 hour. After 7 hours, the condenser was removed and the liquid ammonia was allowed to evaporate over several hours. The crude product was purified by flash chromatography on silica gel (85:15 hexanes:ethyl acetate, product at $R_f = 0.32$, contaminant at $R_f = 0.51$);
- 10 GC/MS m/z 156.1 (M^+), R_t 11.16 minutes.


- The resulting 5-fluoro-2-nitrophenylamine (1.0 eq) and an amine (1.1 eq) e.g. N-methyl piperazine, were dissolved in NMP and triethylamine (2.0 eq) was added. The reaction mixture was heated at 100°C for 3 hours. The solution was then cooled to room temperature and diluted with water. The resulting
- 15 precipitate was filtered and dried under vacuum to provide the 2-nitro-diamino product. Alternatively, the same product may be obtained from commercially available 5-chloro-2-nitrophenylamine under identical conditions except heating at 130°C for 1-2 days. In some examples, the displacement on either 5-fluoro-2-nitrophenylamine or 5-chloro-2-nitrophenylamine can be conducted in neat amine (5
- 20 eq) at 100°C or 130 °C, respectively. The product is isolated in an identical manner. LC/MS m/z 237.1 (MH^+), R_t 1.304 minutes.

- The nitroamine (1.0 eq) and 10% Pd/C (0.1 eq) was suspended in anhydrous ethanol at room temperature. The reaction flask was evacuated and subsequently filled with H_2 . The resulting mixture was then stirred under a
- 25 hydrogen atmosphere overnight. The resulting solution was filtered through Celite and concentrated under vacuum to provide the crude product which was used without further purification.


-98-

Method 2

A round-bottom flask was charged with 2,3-difluoro-6-nitrophenylamine (1 eq) and enough NMP to make a viscous slurry. An amine (5 eq), e.g. N-methyl piperazine, was added and the solution was heated at 100°C. After 2 hours, the solution was cooled and poured into water. A bright yellow solid formed which was filtered and dried. The nitroamine was reduced as in Method 1 to provide the crude product which was used without further purification. LC/MS *m/z* 225.1 (MH⁺), *R*_t 0.335 minutes.

Method 3


To a 0.1 M DMF solution of 1,3-difluoro-2-nitrobenzene was added Et₃N (2 eq) followed by an amine (1 eq), e.g. morpholine. The mixture was stirred for 18 hours and then diluted with water and extracted with ethyl acetate. LC/MS *m/z* 227.2 (MH⁺), *R*_t 2.522 minutes. The combined organic layers were dried over MgSO₄, filtered, and concentrated. Ammonia was condensed into a bomb containing the crude product. The bomb was sealed and heated to 100°C (over 400 psi). After 72 hours the bomb was allowed to cool and the ammonia was evaporated to provide a reddish solid. The nitroamine was reduced as in Method 1 to provide the crude product which was used without further purification. LC/MS *m/z* 194.1 (MH⁺), *R*_t 1.199 minutes.

Method 4

To a stirred NMP solution containing NaH (1.3 eq) was added an alcohol (1.0 eq), e.g. 2-methoxyethanol. The resulting mixture was then stirred for 30 minutes. A slurry of 5-fluoro-2-nitrophenylamine in NMP was then added slowly. The mixture was then heated to 100°C. After 2 hours, the reaction mixture was cooled and water was added. The mixture was then filtered and the captured solid was washed with water and purified by silica gel chromatography (1:1 ethyl acetate:hexane). LC/MS *m/z* 213.2 (MH⁺), *R_t* 2.24 minutes. The nitroamine was reduced as in Method 1 to provide the crude product which was used without further purification. LC/MS *m/z* 183.1 (MH⁺), *R_t* 0.984 minutes.

Method 5

Diisopropyl azodicarboxylate (1.1 eq) was added dropwise to a stirred solution of 4-amino-3-nitrophenol (1.0 eq), triphenylphosphine (1.1 eq), and an alcohol, e.g. N-(2-hydroxyethyl)morpholine (1.0 eq), in tetrahydrofuran at 0°C. The mixture was allowed to warm to room temperature and stirred for 18 hours. The solvent was evaporated, and the product was purified by silica gel chromatography (98:2 CH₂Cl₂:methanol) to yield 4-(2-morpholin-4-ylethoxy)-2-nitrophenylamine as a dark reddish-brown oil. LC/MS *m/z* 268.0 (MH⁺), *R_t* 1.01 minutes. The nitroamine was reduced as in Method 1 to give the crude product which was used without further purification. LC/MS *m/z* 238.3 (MH⁺), *R_t* 0.295 minutes.

Method 6

To a flask charged with 4-amino-3-nitrophenol (1 eq), K_2CO_3 (2 eq),
 5 and 2-butanone was added an alkyl dibromide, e.g. 1,3-dibromopropane
 (1.5 eq). The resulting mixture was then heated at 80°C for 18 hours. After
 cooling, the mixture was filtered, concentrated, and diluted with water. The
 solution was then extracted with CH_2Cl_2 (3 x) and the combined organic layers were
 concentrated to give a solid that was then washed with pentane. LCMS m/z 275.1
 10 (MH⁺), R_t 2.74 minutes.

An acetonitrile solution of the bromide prepared above, an amine,
 e.g. pyrrolidine (5 eq), Cs_2CO_3 (2 eq) and Bu_4NI (0.1 eq) was heated at 70°C for
 48 hours. The reaction mixture was cooled, filtered, and concentrated. The
 residue was dissolved in CH_2Cl_2 , washed with water, and concentrated to give the
 15 desired nitroamine, 2-nitro-4-(3-pyrrolidin-1-ylpropoxy)phenylamine. LCMS m/z
 266.2 (MH⁺), R_t 1.51 minutes. The nitroamine was reduced as in Method 1 to
 provide the crude product which was used without further purification.


Method 7

20 To a suspension of 6-chloro-3-nitropyridin-2-amine (1 eq) in
 acetonitrile was added an amine, e.g. morpholine (4 eq). The resulting reaction
 mixture was stirred at 70°C for 5 hours. The solvent was evaporated under
 reduced pressure, and the residue triturated with ether to provide the desired
 compound as a bright yellow powder. LC/MS m/z 225.0 (MH⁺), R_t 1.79 minutes.

-101-


The nitroamine was reduced as in Method 1 to provide the crude product which was used without further purification.

Method 8:



- 5 A phenol (1 equivalent) and 5-chloro-2-nitro aniline (1 equivalent) were dissolved in DMF, and solid K_2CO_3 (2 equivalents) was added in one portion. The reaction mixture was heated at $120^\circ C$ overnight. The reaction mixture was cooled to room temperature, most of the DMF was distilled off, and water was added to the residue to obtain a precipitate. The solid was dried and purified by
- 10 chromatography on silicagel (2-10% MeOH/ CH_2Cl_2) to afford the desired product. The nitroamine was reduced as in method 1 to give the crude product that was used without further purification.

Method 9




- 15 Furthermore, the introduction of substituents on the benzimidazole ring need not be limited to the early stages of the synthesis and may arise after formation of the quinolinone ring. For example, the crude methyl ester shown in the figure above was dissolved in a 1:1 mixture of EtOH and 30% aqueous KOH and stirred overnight at $70^\circ C$. The reaction mixture was then cooled and acidified
- 20 with 1N HCl to give a precipitate. The solid was filtered, washed with water and dried to obtain 2-(4-amino-2-oxo-1,2-dihydroquinolin-3-yl)-1H-benzimidazole-6-carboxylic acid 2-(4-amino-2-oxo-3-hydroquinolyl)benzimidazole-6-carboxylic acid as a brown solid. LC/MS m/z : 321.1 (MH⁺), R_t 2.26 minutes.


A mixture of 2-(4-amino-2-oxo-1,2-dihydroquinolin-3-yl)-1*H*-benzimidazole-6-carboxylic acid (1 eq) the amine (1 eq), EDC (1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, 1.2 eq), HOAT (1-hydroxy-7-azabenzotriazole, 1.2 eq) and triethylamine (2.5 eq) in DMF, was stirred at 23 °C for 20 h. The reaction mixture was partitioned between water and ethyl acetate. The combined organic layers were dried (Na₂SO₄) and concentrated. Water was added and the precipitate thus formed was filtered off and dried to afford the desired product.

The various 2-amino benzoic acid starting materials used to synthesize isatoic anhydrides may be obtained from commercial sources, prepared by methods known to one of skill in the art, or prepared by the following general Methods 10-11. General isatoic anhydride synthesis methods are described in *J. Med. Chem.* 1981, 24 (6), 735 and *J. Heterocycl. Chem.* 1975, 12(3), 565.

Method 10




Compounds 1-3 were made using similar procedures as found in U.S. Patent No. 4,287,341. Compound 3 was reduced using standard hydrogenation conditions of 10% Pd/C in NH₄OH at 50°C over 48 hours. The product was precipitated by neutralizing with glacial acetic acid, filtering, and washing with water and ether. Yields were about 50%. Compound 5 was prepared in a manner similar to that disclosed in U.S. Patent No. 5,716,993.

Method 11


- Iodination of aniline containing compounds: Iodination was done by a similar procedure as in *J. Med. Chem.* 2001, 44, 6, 917-922. The anthranilic ester in EtOH was added to a mixture of silver sulfate (1 equivalent) and I₂ (1 equivalent). The reaction was typically done after 3 hours at room temperature. The reaction was filtered through celite and concentrated. The residue was taken up in EtOAc and washed with aqueous saturated NaHCO₃ (3x), water (3x), brine (1x), dried (MgSO₄), filtered, and concentrated. The crude product (~5 g) was dissolved in MeOH (60-100 ml), NaOH 6N (25 ml), and water (250 ml). The reactions were typically done after heating at 70-80°C for 4 hours. The reaction mixture was extracted with EtOAc (2x), neutralized with aqueous HCl, filtered to collect the solids, and the solid products were washed with water. The products were dried in vacuo.

In various instances, substitutions on the quinolinone ring may also be introduced after coupling as shown in the general methods 12-15.


-104-

Method 12

Conversion of the C-6 or C-7 halides to an acid group was accomplished using procedures in the following references - Koga, H.; et al., *Tet. Let.*, 1995, 36, 1, 87-90 and Fukuyama, T.; et al., *J. Am. Chem. Soc.*, 1994, 116, 3125-3126.

Method 13

Conversion of the C-6 or C-7 halides to a cyano group was accomplished using procedures in the following reference - Anderson, B.A.; et al., *J. Org. Chem.*, 1998, 63, 8224-828.

Method 14


Conversion of the C-6 or C-7 halides to an aryl group was accomplished using standard Suzuki or Stille procedures such as described below.

-105-

Suzuki Method: To a 1 dram (4 ml) vial was added sequentially the quinolone (1 equivalent), boronic acid (1.2-1.5 equivalents), Pd(dppf)Cl₂, ClCH₂ (0.2 equivalents), DMF (0.5 - 1 ml) and TEA (4 equivalents). The reaction was flushed with argon, capped and heated at 85°C for 12 hours. Once done, the reaction is cooled to room temperature, and filtered with a syringe filter disk. The clear solution is then neutralized with TFA (a couple of drops) and injected directly onto a preparative HPLC. The products are lyophilized to dryness.

Stille Method: To a 1 dram (4 ml) vial was added sequentially the quinolone (1 equivalent), tin reagent (1.8 equivalent), Pd(dppf)Cl₂, ClCH₂ (0.2 equivalents), and DMF (0.5 - 1 ml). The reaction was flushed with argon, capped and heated at 60-85°C for 4 hours. Once done, the reaction is cooled to room temperature, and filtered with a syringe filter disk. The clear solution is then neutralized with TFA (a couple of drops) and injected directly onto a preparative HPLC. The products are lyophilized to dryness.

15 Method 15



A dihaloquinolone such as a difluoroquinolone (12-15 mg) was placed in a 1 dram (2 ml) vial. NMP (dry and pre-purged with argon for 5 minutes) was added to the vial (0.5 ml). The amine reagent (40-50 mg) was added next. If the amine was an HCl salt, the reaction was neutralized with TEA (~1.2-1.5 equivalents). The reaction was purged again with argon for about 5 seconds, and immediately capped. The reaction was typically heated in a heating block at 90-95°C for 18 hours. The reaction was followed by HPLC or LCMS. After taking samples for HPLC, the vial was purged with argon again and capped. Some coupling partners took 24 or 48 hours to reach completion. Less nucleophilic

-106-

amines like pyrrole required the addition of a strong base to reach completion. In these cases, cesium carbonate (2 equivalents based on the amine used) was added to the reaction. Once done, the reaction was cooled to room temperature, and filtered with a syringe filter disk. The clear solution was then neutralized with TFA (a couple of drops) and injected directly onto a preparative HPLC. The products were lyophilized to dryness.

Example 1

Ethyl 2-benzimidazol-2-ylacetate

A solution of 1,2-phenylenediamine (1.0 eq) and ethyl 3-ethoxy-3-
iminopropanoate hydrochloride (1.3 eq) in ethanol was stirred at 90°C overnight.
The reaction was cooled to room temperature and the solvent was removed *in vacuo*. Water and CH₂Cl₂ were added to the residue. The organic layer was separated, dried over Na₂SO₄ and the solvent removed. The solid recovered was used without purification. LC/MS *m/z* 205.2 (MH⁺), R_t 1.44 minutes.

5-(4-Methylpiperazinyl)-2-nitrobenzenecarbonitrile

5-Fluoro-2-nitrobenzenecarbonitrile (1.02 eq) and N-methylpiperazine (1.0 eq) were dissolved in NMP. Triethylamine (2.1 eq) was added, and the resulting solution heated at 100°C for 1 hour. The solution was cooled to room temperature and poured into H₂O. A precipitate formed which was filtered to yield the desired product as a green solid. LC/MS *m/z* 247.3 (MH⁺), R_t 1.46 minutes.

2-Amino-5-(4-methylpiperazinyl)benzenecarbonitrile

5-(4-Methylpiperazinyl)-2-nitrobenzenecarbonitrile (1.0 eq) was dissolved in EtOAc. The flask was purged with nitrogen, and 10% Pd/C (0.1 eq) was added. The flask was evacuated and purged with H₂ three times. The resulting mixture was stirred for three days at room temperature. The mixture was filtered through Celite and the filter pad was washed with EtOAc. The solvent was removed *in vacuo* to give a yellow solid which was purified by silica gel

-107-

chromatography (5:1:95 MeOH:Et₃N:EtOAc) to give the desired product as a yellow solid. LC/MS *m/z* 217.3 (MH⁺), R_t 0.95 minutes.

Method A

4-Amino-3-benzimidazol-2-yl-6-(4-methylpiperazinyl)hydroquinolin-2-one

- 5 Ethyl 2-benzimidazol-2-ylacetate (1.1 eq) and 2-amino-5-(4-methylpiperazinyl)benzenecarbonitrile (1.0 eq) were dissolved in 1,2-dichloroethane, and then SnCl₄ (11 eq) was added. The mixture was heated at reflux overnight. Upon cooling, the mixture was concentrated *in vacuo*. NaOH (3 M) was added to the solid, and the mixture heated at 80°C for 0.5 hours. The solid
10 was filtered and washed sequentially with H₂O, CH₂Cl₂, and acetone. LC/MS indicated that the product was present in the acetone layer and the solid. These fractions were combined and purified by silica gel chromatography (5-10% MeOH in CH₂Cl₂ with 1% Et₃N) to give the desired product. LC/MS *m/z* 375.4 (MH⁺), R_t 1.65 minutes.

15

Example 2

6-Amino-2-(2-morpholin-4-ylethoxy)benzenecarbonitrile

- 4-(Hydroxyethyl)morpholine (1.02 eq) was added to NaH (1.2 eq) in NMP. After 10 minutes, 6-amino-2-fluorobenzenecarbonitrile (1.0 eq) was added in NMP. The resulting mixture was heated at 100°C for 1 hour. The mixture was
20 then cooled and poured into H₂O. The aqueous layer was extracted with EtOAc. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo* to yield a brown gum. The crude material was purified by silica gel chromatography (5:1:95 MeOH:Et₃N:EtOAc) to give the desired product. LC/MS *m/z* 248.3 (MH⁺), R_t 1.26 minutes.

- 25 **4-Amino-3-benzimidazol-2-yl-5-(2-morpholin-4-ylethoxy)hydroquinolin-2-one**
The title compound was synthesized as described in Example 1, Method A using 6-amino-2-(2-morpholin-4-ylethoxy)benzenecarbonitrile. LC/MS *m/z* 406.4 (MH⁺), R_t 1.67 minutes.

Example 3**4-(2-Morpholin-4-ylethoxy)-2-nitrophenylamine**

Diisopropyl azodicarboxylate (1.1 eq) was added dropwise to a stirred solution of 4-amino-3-nitrophenol (1.0 eq), triphenylphosphine (1.1 eq), and
5 N-(2-hydroxyethyl)morpholine (1.0 eq), in THF at 0°C. The mixture was allowed to warm to room temperature and left to stir for 18 hours. The solvent was evaporated and the product was purified by silica gel chromatography (98:2 CH₂Cl₂:MeOH) to yield a dark reddish-brown oil. LC/MS *m/z* 268.0 (MH⁺), R_t 1.01 minutes.

10 4-(2-Morpholin-4-ylethoxy)benzene-1,2-diamine

To a solution 4-(2-morpholin-4-ylethoxy)-2-nitrophenylamine (1.0 eq) in EtOH was added Pd/C (0.1 eq). The reaction vessel was repeatedly purged with nitrogen, then stirred under a hydrogen atmosphere (1 atm) for 18 hours. The product was filtered through a Celite plug, and the plug washed with EtOH. The
15 diamine was used without purification. LC/MS *m/z* 238.3 (MH⁺), R_t 0.295 minutes.

Ethyl 2-[5-(2-morpholin-4-ylethoxy)benzimidazol-2-yl]acetate

The title compound was synthesized as described in Example 1 using 4-(2-morpholin-4-ylethoxy)benzene-1,2-diamine. The organic layer was
20 concentrated and the residue was purified by silica gel chromatography (10:1:2 CH₂Cl₂:MeOH:EtOAc) to yield a dark reddish brown oil. LC/MS *m/z* 334.4 (MH⁺) R_t 1.08 minutes.

4-Amino-3-[5-(2-morpholin-4-ylethoxy)benzimidazol-2-yl]-6-nitrohydroquinolin-2-one

25 The title compound was synthesized as described in Example 1, Method A using ethyl 2-[5-(2-morpholin-4-ylethoxy)benzimidazol-2-yl]acetate and 5-nitroanthranilonitrile. The crude product was purified by silica gel chromatography (5-10% MeOH in CH₂Cl₂ with 1% Et₃N) to give the desired product. LC/MS *m/z* 451.2 (MH⁺), R_t 1.89 minutes.

Example 4**4-Amino-5-(2-morpholin-4-ylethoxy)-3-[5-(2-morpholin-4-ylethoxy)-benzimidazol-2-yl]hydroquinolin-2-one**

- The title compound was synthesized as described in Example 1,
5 Method A using ethyl 2-[5-(2-morpholin-4-ylethoxy)benzimidazol-2-yl]acetate and 6-amino-2-(2-morpholin-4-ylethoxy)benzenecarbonitrile. LC/MS *m/z* 535.4 (MH⁺), *R_t* 1.44 minutes.

Example 5**2-[(Ethoxycarbonyl)methyl]benzimidazole-5-carboxylic acid**

- 10 The title compound was synthesized as described in Example 1 using 3,4-diaminobenzoic acid. The crude material was purified by silica gel chromatography (5:95 MeOH:CH₂Cl₂) to afford the desired product as a white to off-white solid. LC/MS *m/z* 249.1 (MH⁺), *R_t* 1.35 minutes.

Ethyl 2-[5-(N,N-dimethylcarbamoyl)benzimidazol-2-yl]acetate

- 15 2-[(Ethoxycarbonyl)methyl]benzimidazole-5-carboxylic acid (1.0 eq) was dissolved in THF. HBTU (1.1 eq) and diisopropylethylamine (2.0 eq) were added, followed by dimethylamine (2.0 M in THF, 1.1 eq). The reaction was stirred at room temperature overnight then concentrated and the resulting residue was purified by silica gel chromatography (5:95 MeOH:CH₂Cl₂) to afford the
20 desired compound. LC/MS *m/z* 276.2 (MH⁺), *R_t* 1.18 minutes.

[2-(4-amino-2-oxo(3-hydroquinolyl))benzimidazol-5-yl]-N,N-dimethylcarboxamide.

- The title compound was synthesized as described in Example 1,
Method A using ethyl 2-[5-(N,N-dimethylcarbamoyl)benzimidazol-2-yl]acetate and
25 anthranilonitrile. The resulting solid was collected by filtration and washed with water followed by acetone to afford the desired product as a white solid. LC/MS *m/z* 348.3 (MH⁺), *R_t* 1.87 minutes.

Example 6

4-Amino-3-[5-(morpholin-4-ylcarbonyl)benzimidazol-2-yl]hydroquinolin-2-one.

2-[(Ethoxycarbonyl)methyl]benzimidazole-5-carboxylic acid (1.0 eq) was dissolved in THF. HBTU (1.1 eq) and diisopropylethylamine (2.0 eq) were added, followed by morpholine (1.1 eq). The reaction was stirred at room temperature for 3 days then concentrated and purified by silica gel chromatography (5-10% methanol/dichloromethane). The product-containing fractions were concentrated and dissolved in anhydrous 1,2-dichloroethane. Anthranilonitrile (1.0 eq) was added followed by SnCl_4 (5.0 eq) and the reaction was heated at 90°C overnight. The reaction mixture was concentrated and the resulting residue was re-dissolved in NaOH (2 M) and heated at 90°C for 4 hours. After cooling to room temperature, the resulting solid was collected and washed with water followed by acetone to afford the desired product. LC/MS m/z 390.2 (MH+), R_t 1.95 minutes.

Example 7

4-Bromobenzene-1,2-diamine

A solution of 4-bromo-2-nitroaniline (1.0 eq) and SnCl_2 (2.2 eq) in EtOH was heated at reflux for 3 hours. After this time, the solution was poured onto ice, brought to pH 10 with 2M NaOH and extracted with Et₂O. The combined organic layers were dried over MgSO_4 and concentrated. The resulting brown oil was purified by silica gel chromatography (0-50% EtOAc:hexanes) to provide a light yellow solid. LC/MS m/z 187.1 (MH+), R_t 1.33 minutes.

2-Nitro-4-(2-thienyl)phenylamine

4-Bromo-2-nitroaniline (1.0 eq) and Na_2CO_3 (2.0 eq) were dissolved in DMF/H₂O (5:1) at room temperature. Nitrogen was bubbled through the reaction mixture for 5 minutes and $\text{PdCl}_2(\text{dppf})_2$ (0.1 eq) was added. After stirring at 23°C for approximately 10 minutes, 2-thiopeneboronic acid (1.1 eq) in DMF was added and the reaction was heated at 90°C for 12 hours. After this time, the solution was concentrated and partitioned between EtOAc and H₂O. The layers were separated and the aqueous layer was extracted with EtOAc. The combined

-111-

organic layers were dried over MgSO_4 and concentrated under reduced pressure. The resulting black residue was purified by silica gel chromatography (0-20% EtOAc:hexanes) to yield an orange solid. LC/MS m/z 221.1 (MH^+), R_t 2.67 minutes.

5 Ethyl 2-[5-(2-thienyl)benzimidazol-2-yl]acetate

2-Nitro-4-(2-thienyl)phenylamine (1.0 eq) and 10% Pd/C (0.1 eq) were suspended in anhydrous EtOH at room temperature. The reaction flask was evacuated and subsequently filled with H_2 . The resulting mixture was allowed to stir under a hydrogen atmosphere for 3 hours. Ethyl 3-ethoxy-3-iminopropanoate hydrochloride (2.0 eq) was then added and the resulting mixture was heated at reflux for 12 hours. After this time, the solution was filtered through a plug of Celite, concentrated, dissolved in 50 ml of 2N HCl and washed with CH_2Cl_2 . The aqueous layer was brought to pH 12 with concentrated $\text{NH}_4\text{OH}(\text{aq})$ and extracted with CH_2Cl_2 . The combined organic layers were dried with MgSO_4 and concentrated to yield a brown oil which was purified by silica gel chromatography (5:95 MeOH: CH_2Cl_2) to provide a yellow solid. LC/MS m/z 287.1 (MH^+), R_t 1.98 minutes.

4-Amino-3-[5-(2-thienyl)benzimidazol-2-yl]hydroquinolin-2-one

The title compound was synthesized as described in Example 1, Method A using ethyl 2-[5-(2-thienyl)benzimidazol-2-yl]acetate and anthranilonitrile. LC/MS m/z 359.2 (MH^+), R_t 2.68 minutes.

Example 8

5-Fluoro-2-nitrophenylamine

2,4-Difluoronitrobenzene (1.0 eq) was placed in a dry round-bottomed flask equipped with a dry ice condenser charged with acetone/dry ice. Ammonia was condensed into the flask and the resulting solution was stirred at reflux for 7 hours. A yellow precipitate was formed within 1 hours. After 7 hours, the condenser was removed and the liquid ammonia was allowed to evaporate over several hours. The crude product was purified by flash chromatography on silica

-112-

gel (85:15 hexanes:EtOAc, product at $R_f = 0.32$, contaminant at $R_f = 0.51$).
GC/MS m/z 156.1 (M+), R_t 11.16 minutes.

2-Nitro-5-[1-(1,2,4-triazolyl)]phenylamine

- 5 5-Fluoro-2-nitrophenylamine (1.0 eq), 1H-1,2,4-triazole (3.0 eq) and NaH (3.0 eq) in NMP were heated at 100°C for 1 hour. The solution was cooled to room temperature and slowly poured onto ice water. The resulting precipitate was filtered and dried under vacuum to yield the desired product. The resulting solid was recrystallized from EtOH to afford pure product as a bright yellow solid. LC/MS m/z 206.2 (MH+), R_t 1.88 minutes.

10 **Ethyl 2-{5-[1-(1,2,4-triazolyl)]benzimidazol-2-yl}acetate**

The title compound was synthesized as described in Example 7 using 2-nitro-5-[1-(1,2,4-triazolyl)]phenylamine. LC/MS m/z 272.1 (MH+), R_t 1.19 minutes.

15 **4-Amino-3-{5-[1-(1,2,4-triazolyl)]benzimidazol-2-yl}hydroquinolin-2-one**

The title compound was synthesized as described in Example 1, Method A using ethyl 2-{5-[1-(1,2,4-triazolyl)]benzimidazol-2-yl}acetate and anthranilonitrile. The crude solid was collected and purified by silica gel chromatography (92:7:1 CH₂Cl₂:MeOH:Et₃N). LC/MS m/z 344.3 (MH+), R_t 2.01 minutes.

20

Example 9**Method B**

- N*-(4-Chloro-2-cyanophenyl)-2-(5-morpholin-4-ylbenzimidazol-2-yl)acetamideLiHMDS (2.5 eq) was added to ethyl 2-[5-(2-morpholin-4-ylethoxy)benzimidazol-2-yl]acetate (1.0 eq) in THF at -78°C. After 1 hour, 2-amino-5-chlorobenzenecarbonitrile (0.82 eq) in THF was added. The reaction was allowed to warm to 23°C and stirred overnight. The resulting mixture was quenched with NH₄Cl (aqueous saturated solution) and extracted with EtOAc. The combined organic layers were washed with H₂O and brine, dried over Na₂SO₄,
- 25

-113-

filtered and concentrated *in vacuo* to yield a brown solid. The crude material was purified by silica gel chromatography (5:1 EtOAc:hexane) to give the desired product. LC/MS *m/z* 396.1 (MH⁺), R_f 1.79 minutes.

4-Amino-6-chloro-3-(5-morpholin-4-ylbenzimidazol-2-yl)hydroquinolin-2-one

5 N-(4-chloro-2-cyanophenyl)-2-(5-morpholin-4-ylbenzimidazol-2-yl)acetamide (1.0 eq) was heated in NaOMe (0.5 M in MeOH, 18 eq) at 70°C for 2 hours. The resulting mixture was cooled, and the resulting solid was filtered and washed with water to give the desired product. LC/MS *m/z* 396.4 (MH⁺), R_f 2.13 minutes.

10

Example 10

2-Nitro-5-piperidylphenylamine

The title compound was synthesized as described in Example 8 using piperidine (3.0 eq, excess acts as base in place of NaH). The desired product was obtained as a yellow, crystalline solid. LC/MS *m/z* 222.2 (MH⁺), R_f 2.53
15 minutes.

Ethyl 2-(5-piperidylbenzimidazol-2-yl)acetate

The title compound was synthesized as described in Example 7 using 2-nitro-5-piperidylphenylamine. The desired product was obtained as a yellow oil. LC/MS *m/z* 288.3 (MH⁺), R_f 1.31 minutes.

20 **4-amino-3-(5-piperidylbenzimidazol-2-yl)hydroquinolin-2-one**

The title compound was synthesized as described in Example 9, Method B using ethyl 2-(5-piperidylbenzimidazol-2-yl)acetate and anthranilonitrile. The acyclic amide was used crude in the NaOMe cyclization step. The desired product was obtained following purification by silica gel chromatography
25 (96.5:3.0:0.5 CH₂Cl₂:MeOH:Et₃N, R_f 0.2). LC/MS *m/z* 360.4 (MH⁺), R_f 1.83 minutes.

Example 11**[1-(3-Amino-4-nitrophenyl)pyrrolidin-3-yl]dimethylamine**

The title compound was synthesized as described in Example 8 using 3-(dimethylamino)pyrrolidine (3.0 eq, excess amine was used as the base in place of NaH). LC/MS *m/z* 251.3 (MH⁺), R_t 1.25 minutes.

Ethyl 2-{5-[3-(dimethylamino)pyrrolidinyl]benzimidazol-2-yl}acetate

The title compound was synthesized as described in Example 7 using [1-(3-amino-4-nitrophenyl)pyrrolidin-3-yl]dimethylamine. The desired product was obtained as a yellow oil. LC/MS *m/z* 317.4 (MH⁺), R_t 1.36 minutes.

10 **2-{5-[3-(Dimethylamino)pyrrolidinyl]benzimidazol-2-yl}-N-(4-chloro-2-cyanophenyl)acetamide**

The title compound was synthesized as described in Example 9, Method B using ethyl 2-{5-[3-(dimethylamino)pyrrolidinyl]benzimidazol-2-yl}acetate. LC/MS *m/z* 423.4 (MH⁺), R_t 1.67 minutes.

15 **4-Amino-3-{5-[3-(dimethylamino)pyrrolidinyl]benzimidazol-2-yl}-6-chlorohydroquinolin-2-one**

The title compound was synthesized as described in Example 9, Method B using 2-{5-[3-(dimethylamino)pyrrolidinyl]benzimidazol-2-yl}-N-(4-chloro-2-cyanophenyl)acetamide. LC/MS *m/z* 423.4 (MH⁺), R_t 1.71 minutes.

20

Example 12**Ethyl 2-[5-(dimethylamino)benzimidazol-2-yl]acetate**

The title compound was synthesized as described in Example 7 using (3-amino-4-nitrophenyl)dimethylamine. The resulting tan film was purified by silica gel chromatography (5:1:94 MeOH:Et₃N:CH₂Cl₂) to give the desired product.

25 LC/MS 248.3 *m/z* (MH⁺), R_t 1.24 minutes.

-115-

2-[5-(Dimethylamino)benzimidazol-2-yl]-N-(2-cyanophenyl)acetamide

The title compound was synthesized as described in Example 9, Method B using ethyl 2-[5-(dimethylamino)benzimidazol-2-yl]acetate and anthranilonitrile. LC/MS m/z 320.2 (MH⁺), R_t 1.68 minutes.

5 4-Amino-3-[5-(dimethylamino)benzimidazol-2-yl]hydroquinolin-2-one

The title compound was synthesized as described in Example 9, Method B using 2-[5-(dimethylamino)benzimidazol-2-yl]-N-(2-cyanophenyl)acetamide. LC/MS m/z 320.2 (MH⁺), R_t 1.72 minutes.

Example 13**10 Ethyl 2-(5-cyanobenzimidazol-2-yl)acetate**

The title compound was synthesized as described in Example 7 using 4-amino-3-nitro-benzonitrile. LC/MS m/z 230.2 (MH⁺), R_t 1.29 minutes.

2-(4-Amino-2-oxo-3-hydroquinolyl)benzimidazole-5-carbonitrile

- The title compound was synthesized as described in Example 9, Method B using ethyl 2-(5-cyanobenzimidazol-2-yl)acetate and anthranilonitrile (no acyclic amide observed so the NaOMe step was not needed). LC/MS m/z 302.3 (MH⁺), R_t 2.62 minutes.

Example 14**2-(4-Amino-2-oxo-3-hydroquinolyl)benzimidazole-5-carboxamidine**

- 20 2-(4-Amino-2-oxo-3-hydroquinolyl)benzimidazole-5-carbonitrile (1.0 eq) in EtOH was placed into a glass bomb, cooled to 0°C and HCl (g) was bubbled through for 15 minutes. The bomb was then sealed, brought to room temperature and stirred overnight. The solvent was removed *in vacuo*. The residue was dissolved in EtOH in a glass bomb and cooled to 0°C. NH₃ (g) was bubbled
25 through for 15 minutes and the bomb was sealed and heated to 80°C for 5 hours.

The solvent was removed *in vacuo* and the crude product was purified by reversed-phase HPLC. LC/MS *m/z* 319.2 (MH⁺), *R_t* 1.70 minutes.

Example 15

4-Amino-3-[5-(2-morpholin-4-ylethoxy)-benzimidazol-2-yl]hydroquinolin-2-one

5 The title compound was synthesized as described in Example 9, Method B using anthranilonitrile. The crude acyclic amide was used without purification in the NaOMe cyclization step. The crude final product was purified by reversed-phase HPLC (DMSO/5% TFA). LC/MS *m/z* 406.4 (MH⁺), *R_t* 1.56 minutes.

Example 16

10

5-Morpholin-4-yl-2-nitrophenylamine

The title compound was synthesized as described in Example 8 using morpholine (3.0 eq, excess amine was used as the base in place of NaH). LC/MS *m/z* 224.1 (MH⁺), *R_t* 1.89 minutes.

15 Ethyl 2-(5-morpholin-4-ylbenzimidazol-2-yl)acetate

The title compound was synthesized as described in Example 7 using 5-morpholin-4-yl-2-nitrophenylamine. The crude yellow oil was purified by silica gel chromatography (89.5:10:0.5 CH₂Cl₂:MeOH:Et₃N) to yield the desired product as a yellow solid. LC/MS *m/z* 290.3 (MH⁺), *R_t* 1.31 minutes.

20 Method C

4-Hydroxy-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-benzylhydroquinolin-2-one

To a solution of ethyl 2-(5-morpholin-4-ylbenzimidazol-2-yl)acetate (1.0 eq) in anhydrous THF at -78°C under an atmosphere of nitrogen was added LiHMDS (1 M in THF, 3.1 eq) and the solution was stirred for 1 hour. A solution
25 of 1-benzylbenzo[d]1,3-oxazaperhydroine-2,4-dione (1.05 eq) in anhydrous THF was then added dropwise and the resulting solution was allowed to warm to 0°C

-117-

over 1 hour. The resulting mixture was quenched with a saturated aqueous solution of ammonium chloride and the organic layer was separated. The aqueous layer was extracted with CH₂Cl₂ (4 times). The combined organic layers were dried over Na₂SO₄, concentrated *in vacuo*, and the crude material was dissolved in toluene and
5 heated at reflux for 16 hours. The toluene was removed *in vacuo* and the crude material was used without further purification. The product was obtained as a white solid. LC/MS *m/z* 453.1 (MH⁺), R_t 2.91 minutes.

4-Hydroxy-3-(5-morpholin-4-ylbenzimidazol-2-yl)hydroquinolin-2-one

Crude 4-hydroxy-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-
10 benzylhydroquinolin-2-one (1.0 eq) was dissolved in trifluoromethanesulfonic acid and heated at 40°C for 16 hours. The resulting solution was diluted with water and neutralized with 6 N NaOH (aq), whereupon a yellow precipitate formed. The crude solid was isolated by centrifugation and purified by reversed-phase HPLC to produce the desired product as a bright yellow solid. LC/MS *m/z* 363.3 (MH⁺), R_t
15 1.77 minutes.

Example 17

N-[1-(3-Amino-4-nitrophenyl)pyrrolidin-3-yl](*tert*-butoxy)carboxamide

The title compound was synthesized as described in Example 8 using
3-(*tert*-butoxycarbonylamino)pyrrolidine (1.01 eq) with diisopropylethylamine (2.0
20 eq) as the base (in place of NaH). The product was obtained as an orange, crystalline solid. LC/MS *m/z* 323.3 (MH⁺), R_t 2.53 minutes.

Ethyl 2-(5-{3-[(*tert*-butoxy)carbonylamino]pyrrolidinyl}benzimidazol-2-yl)acetate

The title compound was synthesized as described in Example 7 using
25 *N*-[1-(3-amino-4-nitrophenyl)pyrrolidin-3-yl](*tert*-butoxy)carboxamide. The product was obtained as a yellow oil. LC/MS *m/z* 323.3 (MH⁺), R_t 2.53 minutes.

3-[5-(3-aminopyrrolidinyl)benzimidazol-2-yl]-4-hydroxyhydroquinolin-2-one

The title compound was synthesized following the procedure described in Example 16, Method C using ethyl 2-(5-{3-[(*tert*-butoxy)carbonylamino]pyrrolidinyl}benzimidazol-2-yl)acetate. The product was
5 obtained as a yellow solid following cleavage of the benzyl group (see procedure in Example 15). LC/MS *m/e* 362.3 (MH⁺), *R_t* 1.55 minutes.

Example 18**(3-Amino-4-nitrophenyl)[2-(dimethylamino)ethyl]methylamine**

The title compound was synthesized as described in Example 8 using
10 1,1,4-trimethylethylenediamine (1.01 eq) with diisopropylethylamine (2.0 eq) as the base (in place of NaH). The product was obtained as a bright yellow, crystalline solid. LC/MS *m/z* 239.3 (MH⁺), *R_t* 1.29 minutes.

Ethyl 2-(5-{[2-(dimethylamino)ethyl]methylamino}benzimidazol-2-yl)acetate

The title compound was synthesized as described in Example 7 using
15 (3-amino-4-nitrophenyl)[2-(dimethylamino)ethyl]methylamine. The desired product was obtained as a yellow oil. LC/MS *m/z* 305.2 (MH⁺), *R_t* 1.17 minutes.

3-(5-{[2-(Dimethylamino)ethyl]methylamino}benzimidazol-2-yl)-4-hydroxy-1-benzylhydroquinolin-2-one

The title compound was synthesized as described in Example 16,
20 Method C using ethyl 2-(5-{[2-(dimethylamino)ethyl]methylamino}benzimidazol-2-yl)acetate. The product was obtained as a pale yellow solid. LC/MS *m/z* 468.4 (MH⁺), *R_t* 2.26 minutes.

3-(5-{[2-(Dimethylamino)ethyl]methylamino}benzimidazol-2-yl)-4-hydroxyhydroquinolin-2-one

The title compound was synthesized as described in Example 16,
25 Method C using 3-(5-{[2-(dimethylamino)ethyl]methylamino}benzimidazol-2-yl)-4-hydroxy-1-benzylhydroquinolin-2-one. The crude material was purified by

reversed-phase HPLC to yield the product as a yellow solid. LC/MS m/z 378.4 (MH⁺), R_t 1.99 minutes.

Example 19

Method D

5 4-Chloro-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-benzylhydroquinolin-2-one

A solution of 4-hydroxy-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-benzylhydroquinolin-2-one (1.0 eq) and POCl₃ in a dry, round-bottomed flask was heated at 80°C for 2 hours. The excess POCl₃ was removed *in vacuo* and the crude material was quenched with water. The crude product was collected by filtration
10 and purified by silica gel chromatography (1:9 MeOH:CH₂Cl₂). 4-Chloro-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-benzylhydroquinolin-2-one was isolated as a red solid. LC/MS m/z 471.4 (MH⁺), R_t 2.35 minutes.

4-[(2-Methoxyethyl)amino]-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-benzylhydroquinolin-2-one

15 A solution of 4-chloro-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-benzylhydroquinolin-2-one (1.0 eq) and EtOH was treated with 2-methoxyethylamine (10 eq) at room temperature. The resulting solution was heated at reflux for 16 hours and then the solvent was removed *in vacuo*. The crude solid was sonicated in water, filtered, sonicated in hexanes, and filtered again. The crude
20 product was used without further purification. LC/MS m/z 510.4 (MH⁺), R_t 2.20 minutes.

4-[(2-Methoxyethyl)amino]-3-(5-morpholin-4-ylbenzimidazol-2-yl)hydroquinolin-2-one

25 4-[(2-methoxyethyl)amino]-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-benzylhydroquinolin-2-one was debenzylated using the procedure described in Example 16 to produce the title compound. LC/MS m/z 420.2 (MH⁺), R_t 1.57 minutes. 4-[(2-hydroxyethyl)amino]-3-(5-morpholin-4-ylbenzimidazol-2-yl)hydroquinolin-2-one was produced as a side product (see below).

4-[(2-hydroxyethyl)amino]-3-(5-morpholin-4-ylbenzimidazol-2-yl)hydroquinolin-2-one

The title compound was obtained as a side-product of the debenzilation of 4-[(2-methoxyethyl)amino]-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-benzylhydroquinolin-2-one using the procedure described in Example 16 and was isolated by reversed-phase HPLC as a yellow solid. LC/MS *m/z* 406.2 (MH⁺), R_t 1.39 minutes.

Example 20

4-(Methoxyamino)-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-benzylhydroquinolin-2-one

The title compound was synthesized as described in Example 19, Method D using *O*-methylhydroxylamine. The product was used without purification.

4-(Methoxyamino)-3-(5-morpholin-4-ylbenzimidazol-2-yl)hydroquinolin-2-one

The title compound was obtained as a yellow solid after debenzilation of 4-(methoxyamino)-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-benzylhydroquinolin-2-one using the procedure described in Example 16. LC/MS *m/z* 392.2 (MH⁺), R_t 1.82 minutes.

Example 21

***tert*-Butyl-3-{[3-(5-morpholin-4-ylbenzimidazol-2-yl)-2-oxo-1-benzyl-4-hydroquinolyl]amino}piperidinecarboxylate**

The title compound was synthesized as described in Example 19, Method D using 1-*tert*-butoxycarbonyl-3-aminopiperidine. The product was used without purification.

3-(5-Morpholin-4-ylbenzimidazol-2-yl)-4-(3-piperidylamino)hydroquinolin-2-one

The product was obtained as a yellow solid after debenzilation of *tert*-butyl-3-{[3-(5-morpholin-4-ylbenzimidazol-2-yl)-2-oxo-1-benzyl-4-hydroquinolyl]amino}piperidinecarboxylate using the procedure described in

-121-

Example 16. The t-butoxycarbonyl group is removed under the reaction conditions. LC/MS m/z 445.4 (MH⁺), R_t 1.73 minutes.

Example 22

5 *tert*-Butyl-3-([3-(5-morpholin-4-ylbenzimidazol-2-yl)-2-oxo-1-benzyl-4-hydroquinolyl]amino)methyl)piperidinecarboxylate

The title compound was synthesized as described in Example 19, Method D using 1-*tert*-butoxycarbonyl-3-aminomethylpiperidine. The product was used without purification.

10 3-(5-Morpholin-4-ylbenzimidazol-2-yl)-4-[(3-piperidylmethyl)amino]-hydroquinolin-2-one

The title compound was obtained as a yellow solid after debenzylation of *tert*-butyl-3-([3-(5-morpholin-4-ylbenzimidazol-2-yl)-2-oxo-1-benzyl-4-hydroquinolyl]amino)methyl)piperidinecarboxylate using the procedure described in Example 16. LC/MS m/z 459.6 (MH⁺), R_t 1.71 minutes.

15

Example 23

4-{[2-(Dimethylamino)ethyl]amino}-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-benzylhydroquinolin-2-one

20 The title compound was synthesized as described in Example 19, Method D using 1,1-dimethylethylenediamine. The product was used without purification.

4-{[2-(Dimethylamino)ethyl]amino}-3-(5-morpholin-4-ylbenzimidazol-2-yl)hydroquinolin-2-one

25 The title compound was obtained as a yellow solid after debenzylation of 4-{[2-(dimethylamino)ethyl]amino}-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-benzylhydroquinolin-2-one using the procedure described in Example 16. LC/MS m/z 433.4 (MH⁺), R_t 1.55 minutes.

Example 24**3-(5-Morpholin-4-ylbenzimidazol-2-yl)-4-[(oxolan-2-ylmethyl)amino]-1-benzylhydroquinolin-2-one**

The title compound was synthesized as described in Example 19,
5 Method D using 2-aminomethyltetrahydrofuran. The product was used without purification.

3-(5-Morpholin-4-ylbenzimidazol-2-yl)-4-[(oxolan-2-ylmethyl)amino]-hydroquinolin-2-one

The title compound was obtained as a yellow solid after
10 debenzylation of 3-(5-morpholin-4-ylbenzimidazol-2-yl)-4-[(oxolan-2-ylmethyl)amino]-1-benzylhydroquinolin-2-one using the procedure described in Example 16. LC/MS *m/z* 446.5 (MH⁺), R_t 2.19 minutes.

Example 25**4-{[2-(Methylamino)ethyl]amino}-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-benzylhydroquinolin-2-one**
15

The title compound was synthesized as described in Example 19,
Method D using 1-*tert*-butoxycarbonyl-1-methylethylenediamine. The product was used without purification.

4-{[2-(Methylamino)ethyl]amino}-3-(5-morpholin-4-ylbenzimidazol-2-yl)hydroquinolin-2-one
20

The title compound was obtained as a yellow solid after
debenzylation of 4-{[2-(methylamino)ethyl]amino}-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-benzylhydroquinolin-2-one using the procedure described in Example 16. The *t*-butoxycarbonyl group is removed under the reaction conditions.
25 LC/MS *m/z* 419.4 (MH⁺), R_t 1.50 minutes.

Example 26

***tert*-Butyl-3-{[3-(5-morpholin-4-ylbenzimidazol-2-yl)-2-oxo-1-benzyl-4-hydroquinolyl]amino}pyrrolidinecarboxylate**

The title compound was synthesized as described in Example 19,

- 5 Method D using 1-*tert*-butoxycarbonyl-3-aminopyrrolidine. The product was used without purification.

3-(5-Morpholin-4-ylbenzimidazol-2-yl)-4-(pyrrolidin-3-ylamino)hydroquinolin-2-one

The title compound was obtained as a yellow solid after

- 10 debenzylation of *tert*-butyl-3-{[3-(5-morpholin-4-ylbenzimidazol-2-yl)-2-oxo-1-benzyl-4-hydroquinolyl]amino}pyrrolidinecarboxylate using the procedure described in Example 16. LC/MS *m/z* 431.4 (MH⁺), R_t 1.50 minutes.

Example 27

- 15 **4-(((2*S*)-2-Amino-4-methylpentyl)amino)-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-benzylhydroquinolin-2-one**

The title compound was synthesized as described in Example 19,

Method D using (2*S*)-2-*tert*-butoxycarbonylamino-4-methylpentylamine. The product was used without purification.

- 20 **4-(((2*S*)-2-Amino-4-methylpentyl)amino)-3-(5-morpholin-4-ylbenzimidazol-2-yl)hydroquinolin-2-one**

The title compound was obtained as a yellow solid after

debenzylation of 4-(((2*S*)-2-amino-4-methylpentyl)amino)-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-benzylhydroquinolin-2-one using the procedure described in Example 16. LC/MS *m/z* 461.4 (MH⁺), R_t 1.78 minutes.

Example 28

t-Butoxycarbonyl protected 4-(((2S)-2-amino-3-methylbutyl)amino)-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-benzylhydroquinolin-2-one

- The title compound was synthesized as described in Example 19,
5 Method D using (2S)-2-*tert*-butoxycarbonylamino-3-methylbutylamine. The product was used without purification.

4-(((2S)-2-Amino-3-methylbutyl)amino)-3-(5-morpholin-4-ylbenzimidazol-2-yl)hydroquinolin-2-one

- The title compound was obtained as a yellow solid after
10 debenzylation of 4-(((2S)-2-amino-3-methylbutyl)amino)-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-benzylhydroquinolin-2-one using the procedure described in Example 16. The t-butoxycarbonyl group is removed under the reaction conditions. LC/MS *m/z* 447.5 (MH⁺), R_t 2.96 minutes.

Example 29

- 15 **4-Amino-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-benzylhydroquinolin-2-one**

The title compound was synthesized as described in Example 19,
Method D using ammonia in a sealed glass tube. The product was used without purification.

4-Amino-3-(5-morpholin-4-ylbenzimidazol-2-yl)hydroquinolin-2-one

- 20 The title compound was obtained as a bright yellow solid after debenzylation of 4-amino-3-(5-morpholin-4-ylbenzimidazol-2-yl)-1-benzylhydroquinolin-2-one using the procedure described in Example 16 and purification by reversed-phase HPLC. LC/MS *m/z* 362.3 (MH⁺), R_t 1.61 minutes.

Example 30

- 25 **3-Benzimidazol-2-yl-4-hydroxy-1-benzylhydroquinolin-2-one**

The title compound was synthesized as described in Example 16,
Method C using ethyl 2-benzimidazol-2-ylacetate. The product was obtained as a

-125-

white solid and used without further purification. LC/MS m/z 368.4 (MH⁺), R_t 2.99 minutes.

3-(Benzimidazol-2-yl)-4-chloro-1-benzylhydroquinolin-2-one

- The title compound was synthesized as described in Example 19,
5 Method D using 3-benzimidazol-2-yl-4-hydroxy-1-benzylhydroquinolin-2-one. The crude product was used without purification.

Example 31

3-Benzimidazol-2-yl-4-(methylamino)hydroquinolin-2-one

- The benzylated title compound was synthesized as described in
10 Example 19, Method D using methylamine and 3-(benzimidazol-2-yl)-4-chloro-1-benzylhydroquinolin-2-one. The product was obtained after debenzylation as a yellow solid using the procedure described in Example 16. LC/MS m/z 291.3 (MH⁺), R_t 1.64 minutes.

Example 32

- 15 **3-Benzimidazol-2-yl-4-(ethylamino)hydroquinolin-2-one**

- The benzylated title compound was synthesized as described in
Example 19, Method D using ethylamine and 3-(benzimidazol-2-yl)-4-chloro-1-benzylhydroquinolin-2-one. The title compound was obtained after debenzylation as a yellow solid using the procedure described in Example 16. LC/MS m/z 305.3
20 (MH⁺), R_t 2.01 minutes.

Example 33

3-Benzimidazol-2-yl-4-[(oxolan-2-ylmethyl)amino]hydroquinolin-2-one

- The benzylated title compound was synthesized as described in
Example 19, Method D using 2-aminomethyltetrahydrofuran and 3-(benzimidazol-
25 2-yl)-4-chloro-1-benzylhydroquinolin-2-one. The title compound was obtained after debenzylation as a yellow solid using the procedure described in Example 16.
LC/MS m/z 361.2 (MH⁺), R_t 1.74 minutes.

Example 34**3-Benzimidazol-2-yl-4-[(4-piperidylmethyl)amino]hydroquinolin-2-one**

The protected title compound was synthesized as described in Example 19, Method D using 1-*tert*-butoxycarbonyl-4-aminomethylpiperidine and
5 3-(benzimidazol-2-yl)-4-chloro-1-benzylhydroquinolin-2-one. The title compound was obtained after deprotection and debenylation as a yellow solid using the procedure described in Example 16. LC/MS *m/z* 374.3 (MH⁺), *R_t* 1.29 minutes.

Example 35**3-Benzimidazol-2-yl-4-[(4-fluorophenyl)amino]hydroquinolin-2-one**

10 The benzylation title compound was synthesized as described in Example 19, Method D using 4-fluoroaniline and 3-(benzimidazol-2-yl)-4-chloro-1-benzylhydroquinolin-2-one. The title compound was obtained after debenylation as a yellow solid using the procedure described in Example 16. LC/MS *m/z* 371.2 (MH⁺), *R_t* 1.92 minutes.

Example 36**3-Benzimidazol-2-yl-4-(methoxyamino)hydroquinolin-2-one**

15 The benzylation title compound was synthesized as described in Example 19, Method D using *O*-methylhydroxylamine and 3-(benzimidazol-2-yl)-4-chloro-1-benzylhydroquinolin-2-one. The title compound was obtained after
20 debenylation as a yellow solid using the procedure described in Example 16. LC/MS *m/z* 307.3 (MH⁺), *R_t* 1.77 minutes.

Example 37**3-Benzimidazol-2-yl-4-(benzimidazol-6-ylamino)hydroquinolin-2-one**

The benzylation title compound was synthesized as described in
25 Example 19, Method D using 5-aminobenzimidazole and 3-(benzimidazol-2-yl)-4-chloro-1-benzylhydroquinolin-2-one. The title compound was obtained after debenylation as a yellow solid using the procedure described in Example 16. LC/MS *m/z* 393.4 (MH⁺), *R_t* 1.41 minutes.

-127-

Example 38**3-Benzimidazol-2-yl-4-(phenylamino)hydroquinolin-2-one**

The benzylated title compound was synthesized as described in Example 19, Method D using aniline and 3-(benzimidazol-2-yl)-4-chloro-1-benzylhydroquinolin-2-one. The title compound was obtained after debenzylation as a yellow solid using the procedure described in Example 16. LC/MS m/z 353.4 (MH⁺), R_t 2.38 minutes.

Example 39**3-Benzimidazol-2-yl-4-(quinuclidin-3-ylamino)hydroquinolin-2-one**

The benzylated title compound was synthesized as described in Example 19, Method D using 3-aminoquinuclidine and 3-(benzimidazol-2-yl)-4-chloro-1-benzylhydroquinolin-2-one. The title compound was obtained after debenzylation as a yellow solid using the procedure described in Example 16. LC/MS m/z 386.4 (MH⁺), R_t 1.82 minutes.

Example 40**3-Benzimidazol-2-yl-4-[(imidazol-5-ylmethyl)amino]hydroquinolin-2-one**

The benzylated title compound was synthesized as described in Example 19, Method D using 4-aminomethyl-1H-imidazole and 3-(benzimidazol-2-yl)-4-chloro-1-benzylhydroquinolin-2-one. The title compound was obtained after debenzylation as a yellow solid using the procedure described in Example 16. LC/MS m/z 357.4 (MH⁺), R_t 1.34 minutes.

Example 41**3-Benzimidazol-2-yl-4-(morpholin-4-ylamino)hydroquinolin-2-one**

The benzylated title compound was synthesized as described in Example 19, Method D using 4-aminomorpholine and 3-(benzimidazol-2-yl)-4-chloro-1-benzylhydroquinolin-2-one. The title compound was obtained after debenzylation as a yellow solid using the procedure described in Example 16. LC/MS *m/z* 362.4 (MH⁺), R_t 1.42 minutes.

Example 42**3-Benzimidazol-2-yl-4-hydrazinohydroquinolin-2-one**

The benzylated title compound was synthesized as described in Example 19, Method D using hydrazine and 3-(benzimidazol-2-yl)-4-chloro-1-benzylhydroquinolin-2-one. The title compound was obtained as a yellow solid after debenzylation using the procedure described in Example 16. LC/MS *m/z* 292.3 (MH⁺), R_t 1.19 minutes.

Example 43**3-Benzimidazol-2-yl-2-oxohydroquinoline-4-carbonitrile**

3-Benzimidazol-2-yl-4-chloro-1-benzylhydroquinolin-2-one (1 eq) was dissolved in DMA, and CuCN (10 eq) was added in one portion. The reaction mixture was stirred at 90°C overnight. The resulting mixture was allowed to cool to room temperature, water was added, and the orange precipitate was removed by filtration. The solid was treated with a solution of hydrated FeCl₃ at 70°C for 1 hour. The suspension was centrifuged and the solution removed. The remaining solid was washed with 6N HCl (2 times), sat. Na₂CO₃ (2 times), water (2 times) and lyophilized. The resulting powder was dissolved in 1 mL of triflic acid and heated at 60°C overnight. The resulting mixture was cooled to 0°C and water was slowly added. Saturated LiOH was added dropwise to the suspension to a pH of 8, then the solid was filtered and washed with water (3 times). Purification by reversed-phase HPLC afforded the desired product. LC/MS *m/z* 287.1 (MH⁺), R_t 1.89 minutes.

Example 44**Ethyl 2-(5,6-dimethylbenzimidazol-2-yl)acetate**

The title compound was synthesized as described in Example 1 using 4,5-dimethylbenzene-1,2-diamine. The crude yellow oil was purified first by silica gel chromatography (96.5:3.0:0.5, CH₂Cl₂:MeOH:Et₃N), and then by
5 recrystallization from toluene to yield the title compound as a pale, yellow solid. LC/MS *m/z* 233.1 (MH⁺), R_t 1.73 minutes.

3-(5,6-Dimethylbenzimidazol-2-yl)-4-hydroxy-1-benzylhydroquinolin-2-one

The title compound was synthesized as described in Example 16,
10 Method C using ethyl 2-(5,6-dimethylbenzimidazol-2-yl)acetate. The crude material was purified by silica gel chromatography (98.5:1.5, CH₂Cl₂:MeOH) to yield the title compound as a yellow solid. LC/MS *m/z* 396.2 (MH⁺), R_t 3.60 minutes.

3-(5,6-Dimethylbenzimidazol-2-yl)-4-chloro-1-benzylhydroquinolin-2-one

15 The title compound was synthesized as described in Example 19, Method D using 3-(5,6-dimethylbenzimidazol-2-yl)-4-hydroxy-1-benzylhydroquinolin-2-one. The title compound was obtained as an orange-yellow solid. LC/MS *m/z* 414.2 (MH⁺), R_t 2.47 minutes.

tert-Butyl 3-{[3-(5,6-dimethylbenzimidazol-2-yl)-2-oxo-1-benzyl-4-hydroquinolyl]amino}piperidinecarboxylate
20

The title compound was synthesized as described in Example 19, Method D using 1-*tert*-butoxycarbonyl-3-aminopiperidine. The crude material was purified by silica gel chromatography (99:1 CH₂Cl₂:MeOH) to yield the title compound as a yellow solid. LC/MS *m/z* 578.5 (MH⁺), R_t 3.05 minutes.

3-(5,6-Dimethylbenzimidazol-2-yl)-4-(3-piperidylamino)hydroquinolin-2-one
25

tert-Butyl 3-{[3-(5,6-dimethylbenzimidazol-2-yl)-2-oxo-1-benzyl-4-hydroquinolyl]amino}piperidine-carboxylate was debenzylated as described in Example 16. The crude material was purified by reversed-phase HPLC to yield the title compound as a light yellow solid. LC/MS *m/z* 388.4 (MH⁺), R_t 1.61 minutes.

Example 45**3H-Imidazo[4,5-b]pyridin-2-ylacetonitrile**

5 Ethyl cyanoacetate (1.5 eq) and 2,3-diaminopyridine (1 eq) were heated at 185°C for 30 minutes. The reaction mixture was cooled to room temperature and the black solid was triturated with ether. The desired product was thus obtained as a dark brown powder. LC/MS m/z 159.1 (MH⁺), R_t 0.44 minutes.

Ethyl 3H-imidazo[4,5-b]pyridin-2-ylacetate

10 3H-Imidazo[4,5-b]pyridin-2-ylacetonitrile was suspended in EtOH, and gaseous HCl was bubbled through for 3 hours. The suspension initially seemed to dissolve, but a precipitate started forming almost immediately. The reaction mixture was cooled to 0°C and a cold saturated NaHCO₃ solution was carefully added. Solid NaHCO₃ was also added to bring the pH to a value of 7.6. The
15 aqueous phase was then extracted with EtOAc, and the organic extracts were dried (Na₂SO₄). After evaporation of the solvent under reduced pressure, the residue was purified by chromatography on silicagel (10% MeOH in CH₂Cl₂ with 1% Et₃N) providing the desired product as a light brown solid. LC/MS m/z 206.1 (MH⁺), R_t 0.97 minutes.

20 4-Amino-3-(3H-imidazo[4,5-b]pyridin-2-yl)quinolin-2(1H)-one

LiHMDS (3.0 eq) was added to ethyl 3H-imidazo[4,5-b]pyridin-2-ylacetate (1.0 eq) in THF at -78°C. After 20 minutes, a solution of 2-aminobenzene carbonitrile (1.1 eq) in THF was added. The resulting mixture was allowed to warm to room temperature, stirred for 3 hours, and then refluxed
25 overnight. The mixture was cooled to 0°C and quenched with an aqueous saturated NH₄Cl solution. A precipitate formed, was filtered off, and was washed repeatedly with ether to yield the desired compound as a light brown solid. LC/MS m/z 278.2 (MH⁺), R_t 1.82 minutes.

Example 46**6-Morpholin-4-yl-3-nitropyridin-2-amine**

Morpholine (4 eq) was added to a suspension of 6-chloro-3-nitropyridin-2-amine (1 eq) in CH₃CN, and the reaction mixture was stirred at 70°C for 5 hours. The solvent was evaporated under reduced pressure, and the residue was triturated with ether to afford the desired compound as a bright yellow powder. LC/MS *m/z* 225.0 (MH⁺), R_t 1.79 minutes.

Ethyl (5-morpholin-4-yl-3H-imidazo[4,5-b]pyridin-2-yl)acetate

To a solution 6-chloro-3-nitropyridin-2-amine (1.0 eq) in EtOH was added Pd/C (0.1 eq). The reaction vessel was repeatedly purged with hydrogen and then stirred under a hydrogen atmosphere (1 atm) for 18 hours. Ethyl 3-ethoxy-3-iminopropanoate hydrochloride (2.0 eq) was added in one portion, and the reaction mixture was refluxed overnight. The reaction mixture was cooled to room temperature, filtered through a celite plug, and the plug was washed with EtOH. After evaporation of the solvent under reduced pressure, the residue was purified by silica gel chromatography (5% MeOH in CH₂Cl₂ with 1% Et₃N) providing the desired product as a brown solid. LC/MS *m/z* 291.3 (MH⁺), R_t 1.71 minutes.

4-Amino-3-(5-morpholin-4-yl-3H-imidazo[4,5-b]pyridin-2-yl)quinolin-2(1H)-one

The title compound was synthesized as described in Example 45, Method E using ethyl 2-(5-morpholin-4-ylimidazolo[5,4-b]pyridin-2-yl)acetate and 2-aminobenzenecarbonitrile, with a modified workup procedure.. After quenching with a saturated aqueous ammonium chloride solution, the two phases were separated and the aqueous phase extracted with EtOAc. Upon standing, a solid formed and precipitated out of the organic extracts. The precipitate, a dark brown solid, was filtered off and dried. Purification by reverse phase chromatography afforded the desired product as a reddish solid. LC/MS *m/z* 363.2 (MH⁺), R_t 2.20 minutes.

Example 47

4-Amino-5-[(2R,6S)-2,6-dimethylmorpholin-4-yl]-3-(3H-imidazo[4,5-b]pyridin-2-yl)quinolin-2(1H)-one

LiHMDS (3.0 eq) was added to ethyl 3H-imidazo[4,5-b]pyridin-2-ylacetate (1.0 eq) in THF at -78°C. After 20 minutes, a solution of 2-amino-6-[(2R,6S)-2,6-dimethylmorpholin-4-yl]benzonitrile (1.1 eq) in THF was added. The resulting mixture was allowed to warm to room temperature, stirred for 2 hours and then heated to 60°C overnight. The mixture was cooled to 0°C and quenched with an aqueous saturated NH₄Cl solution. The aqueous phase was extracted with CH₂Cl₂ (5 times) and the organic extracts were collected, dried (Na₂SO₄), and concentrated. The crude product was purified by HPLC. LC/MS *m/z* 391.2 (MH⁺), R_t 2.35 minutes.

Example 48

Ethyl {5-[3-(dimethylamino)pyrrolidin-1-yl]-3H-imidazo[4,5-b]pyridin-2-yl}acetate

6-chloro-3-nitro-2-aminopyridine (1.0 eq) and 3-(dimethylamino)pyrrolidine (1.1 eq) were dissolved in CH₃CN and diisopropylethylamine (2.0 eq) was added. The reaction mixture was heated at 70°C overnight. The solution was cooled to room temperature, and the solvent was evaporated. The residue was triturated with ether and water and dried under vacuum (LC/MS *m/z* 252.2 (MH⁺), R_t 1.09 minutes). The isolated product (1.0 eq) and 10% Pd/C (0.1 eq) were suspended in anhydrous EtOH at room temperature. The reaction flask was evacuated and subsequently filled with H₂. The resulting mixture was allowed to stir under a hydrogen atmosphere overnight. Ethyl 3-ethoxy-3-iminopropanoate hydrochloride (2.0 eq) was then added and the resulting mixture was heated at reflux overnight. The solution was then filtered through Celite and evaporated under reduced pressure. The residue was suspended in CH₂Cl₂ and concentrated NH₄OH was added until a pH of 11 was achieved. The NH₄Cl thus formed was filtered off. The two phases were separated, and the organic phase was dried (Na₂SO₄). Evaporation of the solvent and trituration of the

-133-

residue with ether gave a light green powder. LC/MS m/z 318.1 (MH⁺), R_t 1.11 minutes.

4-Amino-3-{5-[3-(dimethylamino)pyrrolidin-1-yl]-3H-imidazo[4,5-b]pyridin-2-yl}quinolin-2(1H)-one

- 5 LiHMDS (3.5 eq) was added to ethyl {5-[3-(dimethylamino)pyrrolidin-1-yl]-3H-imidazo[4,5-b]pyridin-2-yl}acetate (1.0 eq) in THF at -40 °C. After 10 minutes, a solution of 2-aminobenzecarbonitrile (1.1 eq) in THF was added. The resulting mixture was allowed to warm to room temperature, stirred for 1 h and then heated to 60 °C overnight. The mixture was
- 10 cooled to room temperature and quenched with NH₄Cl (aq, saturated). The aqueous phase was extracted with CH₂Cl₂ (5 times). The product crashes out of the organic solution during the extractions. Evaporation of the solvent under reduced pressure afforded a brown solid that was triturated repeatedly with MeOH and acetone to obtain a yellow greenish powder. LC/MS m/z 390.2 (MH⁺), R_t 1.48 minutes.

15

Example 49

2-(4-Ethylpiperazinyl)-6-nitrobenzenecarbonitrile

- 2,6-Dinitrobenzenecarbonitrile (1.0 eq) and ethylpiperazine (3.6 eq) were dissolved in DMF. The resulting solution heated at 90°C for 2 hours. The solution was cooled to room temperature and poured into H₂O. A precipitate
- 20 formed which was filtered to yield the desired product as a brown solid. LC/MS m/z 260.1 (MH⁺), R_t 1.69 minutes.

6-Amino-2-(4-ethylpiperazinyl)benzenecarbonitrile

- 2-(4-Ethylpiperazinyl)-6-nitrobenzenecarbonitrile (1.0 eq) was dissolved in EtOH and EtOAc. The flask was purged with N₂, and 10% Pd/C (0.1
- 25 eq) was added. The flask was evacuated and purged with H₂ three times. The resulting mixture was stirred overnight at room temperature. The mixture was filtered through Celite, and the filter pad was washed with EtOAc. The solvent was

-134-

removed *in vacuo* to provide the desired product as a yellow solid. LC/MS *m/z* 231.2 (MH⁺), R_t 1.42 minutes.

4-Amino-3-(1H-benzimidazol-2-yl)-5-(4-ethylpiperazin-1-yl)quinolin-2(1H)-one

t-BuLi (3.1 eq) was added to ethyl 2-benzimidazol-2-ylacetate (1.0 eq) and 6-amino-2-(4-ethylpiperazinyl)benzenecarbonitrile (1.0 eq) in THF at 0°C. The reaction was stirred overnight. The resulting mixture was quenched with NH₄Cl (aq, saturated) and extracted with EtOAc. The combined organic layers were washed with H₂O and brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo* to yield a brown solid. The crude material was triturated with CH₂Cl₂ and MeOH to provide a tan solid. LC/MS *m/z* 389.1 (MH⁺), R_t 1.80 minutes.

Examples 50-154

The 2-aminobenzonitriles or isatoic anhydride starting materials used to synthesize Examples 50-154 are readily recognizable by one skilled in the art. They were either commercially available or synthesized following the examples shown above (e.g. Examples 1, 2, and 49). The anhydride 6-chloro-1-(phenylmethyl)-2H-3,1-benzoxazine-2,4(1H)-dione was synthesized following the general isatoic anhydride synthesis methods described in *J. Med. Chem.* **1981**, 24 (6), 735 and *J. Heterocycl. Chem.* **1975**, 12(3), 565.

The benzimidazole acetates were formed by reacting aryl diamines with ethyl 3-ethoxy-3-iminopropanoate hydrochloride as shown in Example 1. The requisite diamines used in the syntheses are also readily recognizable by one skilled in the art and may be synthesized following Methods 1-9. The isatoic anhydrides were coupled with the benzimidazole acetates using methods C and D. The 2-aminobenzonitriles were coupled with the benzimidazole acetates using method B, the coupling method of example 49, or the general procedure set forth below.

Method E

LiHMDS (3-4 eq) was added to the benzimidazole acetate (1.0 eq) in THF (at a constant temperature ranging from -78°C to 0°C). After 20 minutes, a solution of the 2-aminobenzonitrile (1.1 eq) in THF was then added. The resulting mixture was allowed to warm to room temperature, stirred for 1-3 hours and was then heated to approx. 40°C - 65°C (1 hour to 12 hours). The mixture was cooled to 0°C and quenched with NH_4Cl (aq, saturated). The aqueous phase was extracted with CH_2Cl_2 or EtOAc, and the organic extracts were collected, dried (Na_2SO_4), and filtered. Evaporation of the solvent under reduced pressure and purification of the residue by silica gel chromatography or HPLC provided the 4-amino quinolinone products.

Example	Name	LC/MS m/z (MH+)
50	4-amino-3-{5-[(3S)-3-(dimethylamino)pyrrolidin-1-yl]-1H-benzimidazol-2-yl}quinolin-2(1H)-one	389.4
51	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-chloroquinolin-2(1H)-one	420
52	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-chloroquinolin-2(1H)-one	420
53	3-(1H-benzimidazol-2-yl)-4-[(3R)-3-(dimethylamino)pyrrolidin-1-yl]quinolin-2(1H)-one	374.2
54	3-(1H-benzimidazol-2-yl)-6-chloro-4-[(3R)-3-(dimethylamino)pyrrolidin-1-yl]quinolin-2(1H)-one	408.1
55	4-amino-3-[5-(4-ethylpiperazin-1-yl)-1H-benzimidazol-2-yl]-1-methylquinolin-2(1H)-one	403.2
56	4-amino-3-(6-piperazin-1-yl)-1H-benzimidazol-2-ylquinolin-2(1H)-one	361.2
57	4-amino-3-[6-(pyridin-4-ylmethyl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	368.2
58	4-amino-3-{5-[(3R,5S)-3,5-dimethylpiperazin-1-yl]-1H-benzimidazol-2-yl}quinolin-2(1H)-one	389.4

59	4-amino-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	375.2
60	4-amino-3-(6-methyl-5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one	376
61	4-amino-3-{5-[(1-methylpiperidin-3-yl)oxy]-1H-benzimidazol-2-yl}quinolin-2(1H)-one	390.1
62	4-amino-3-{5-[(2R,6S)-2,6-dimethylmorpholin-4-yl]-6-fluoro-1H-benzimidazol-2-yl}quinolin-2(1H)-one	408.2
63	4-amino-3-{5-[(1-methylpyrrolidin-3-yl)oxy]-1H-benzimidazol-2-yl}quinolin-2(1H)-one	376.2
64	4-amino-3-[5-(4-methyl-1,4-diazepan-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	389.2
65	4-amino-3-{5-[(3R)-3-(dimethylamino)pyrrolidin-1-yl]-1H-benzimidazol-2-yl}quinolin-2(1H)-one	389.2
66	4-amino-6-chloro-3-{5-[(3R)-3-(dimethylamino)pyrrolidin-1-yl]-1H-benzimidazol-2-yl}quinolin-2(1H)-one	423
67	ethyl {4-[2-(4-amino-2-oxo-1,2-dihydroquinolin-3-yl)-1H-benzimidazol-6-yl]piperazin-1-yl}acetate	447.2
68	4-amino-3-{6-[methyl(1-methylpiperidin-4-yl)amino]-1H-benzimidazol-2-yl}quinolin-2(1H)-one	403.1
69	3-[6-(4-acetylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-aminoquinolin-2(1H)-one	403.3
70	4-amino-3-[6-(1,4'-bipiperidin-1'-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	443.3
71	2-(4-amino-2-oxo-1,2-dihydroquinolin-3-yl)-1H-benzimidazole-6-carboxylic acid	321.2
72	4-amino-5-(methyloxy)-3-[6-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	405.3
73	4-amino-3-{6-[4-(1-methylethyl)piperazin-1-yl]-1H-benzimidazol-2-yl}quinolin-2(1H)-one	403.3
74	{4-[2-(4-amino-2-oxo-1,2-dihydroquinolin-3-yl)-1H-benzimidazol-6-yl]piperazin-1-yl}acetic acid	419.2
75	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)quinolin-2(1H)-one	386.1
76	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)quinolin-2(1H)-one	386.1
77	4-amino-3-[5-(4-ethylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	389.1

78	4-amino-3-(5-((2S,5S)-2- [(dimethylamino)methyl]-5- methylmorpholin-4-yl)-1H-benzimidazol-2- yl)quinolin-2(1H)-one	433.3
79	4-amino-6-chloro-3-[5-(4-methylpiperazin- 1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)- one	409.2
80	4-amino-6-chloro-3-{5-[(3S)-3- (dimethylamino)pyrrolidin-1-yl]-1H- benzimidazol-2-yl}quinolin-2(1H)-one	423.1
81	4-amino-5,6-dichloro-3-{5-[(3S)-3- (dimethylamino)pyrrolidin-1-yl]-1H- benzimidazol-2-yl}quinolin-2(1H)-one	457.2
82	4-amino-5,6-dichloro-3-[5-(4- methylpiperazin-1-yl)-1H-benzimidazol-2- yl]quinolin-2(1H)-one	443.2
83	4-amino-3-(1H-benzimidazol-2-yl)-6- [(pyridin-2-ylmethyl)oxy]quinolin-2(1H)-one	384.2
84	4-amino-3-(1H-benzimidazol-2-yl)-6- [(2R,6S)-2,6-dimethylmorpholin-4- yl]quinolin-2(1H)-one	390.1
85	4-amino-3-(1H-benzimidazol-2-yl)-6- morpholin-4-ylquinolin-2(1H)-one	362.2
86	4-amino-3-(1H-benzimidazol-2-yl)-5-[(1- methylpiperidin-3-yl)oxy]quinolin-2(1H)-one	390.2
87	4-amino-3-(1H-benzimidazol-2-yl)-5- [(pyridin-2-ylmethyl)oxy]quinolin-2(1H)-one	384.1
88	4-amino-3-(5-morpholin-4-yl-1H- benzimidazol-2-yl)-5-[(pyridin-4- ylmethyl)oxy]quinolin-2(1H)-one	469.2
89	4-amino-3-(1H-benzimidazol-2-yl)-5- (methyloxy)quinolin-2(1H)-one	307.1
90	4-amino-3-(5-methyl-1H-benzimidazol-2- yl)-5-(methyloxy)quinolin-2(1H)-one	321.1
91	4-amino-3-{5-[(2R,6S)-2,6- dimethylmorpholin-4-yl]-1H-benzimidazol- 2-yl}-5-(methyloxy)quinolin-2(1H)-one	420.2
92	4-amino-3-(1H-benzimidazol-2-yl)-5- morpholin-4-ylquinolin-2(1H)-one	362.2
93	4-amino-3-(1H-benzimidazol-2-yl)-5- [(2R,6S)-2,6-dimethylmorpholin-4- yl]quinolin-2(1H)-one	390.2
94	4-amino-3-(1H-benzimidazol-2-yl)-5-(4- methylpiperazin-1-yl)quinolin-2(1H)-one	375.1
95	4-amino-5,6-dichloro-3-(5-morpholin-4-yl- 1H-benzimidazol-2-yl)quinolin-2(1H)-one	430

96	3-{5-[(2-morpholin-4-ylethyl)oxy]-1H-benzimidazol-2-yl}quinolin-2(1H)-one	391.3
97	4-amino-3-{5-[(3-pyrrolidin-1-ylpropyl)oxy]-1H-benzimidazol-2-yl}quinolin-2(1H)-one	404
98	4-amino-3-{5-[(3-morpholin-4-ylpropyl)oxy]-1H-benzimidazol-2-yl}quinolin-2(1H)-one	420.4
99	4-amino-6-fluoro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one	380
100	4-amino-3-{5-[3-(dimethylamino)pyrrolidin-1-yl]-1H-benzimidazol-2-yl}-6-fluoroquinolin-2(1H)-one	407
101	4-amino-3-(1H-benzimidazol-2-yl)-6-fluoroquinolin-2(1H)-one	295
102	4-amino-3-(6-fluoro-5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one	380
103	4-amino-3-{5-[(tetrahydrofuran-2-ylmethyl)oxy]-1H-benzimidazol-2-yl}quinolin-2(1H)-one	377
104	4-amino-6-fluoro-3-(6-fluoro-5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one	398
105	4-amino-3-[6-fluoro-5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	393
106	4-amino-3-(5-{[2-(methyloxy)ethyl]oxy}-1H-benzimidazol-2-yl)quinolin-2(1H)-one	351
107	4-amino-3-[4,6-difluoro-5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	411
108	4-amino-3-{5-[3-(dimethylamino)pyrrolidin-1-yl]-1H-benzimidazol-2-yl}-5-fluoroquinolin-2(1H)-one	407.1
109	4-amino-5-fluoro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	393.1
110	4-amino-5-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	409.1
111	4-amino-3-{5-[3-(dimethylamino)pyrrolidin-1-yl]-6-fluoro-1H-benzimidazol-2-yl}quinolin-2(1H)-one	407.1
112	4-amino-5-chloro-3-{5-[3-(dimethylamino)pyrrolidin-1-yl]-1H-benzimidazol-2-yl}quinolin-2(1H)-one	423.1
113	4-amino-6-chloro-3-{5-[3-(dimethylamino)pyrrolidin-1-yl]-6-fluoro-1H-benzimidazol-2-yl}quinolin-2(1H)-one	441

114	4-amino-5-[(2R,6S)-2,6-dimethylmorpholin-4-yl]-3-(3H-imidazo[4,5-b]pyridin-2-yl)quinolin-2(1H)-one	391.2
115	4-amino-3-(6-thiomorpholin-4-yl)-1H-benzimidazol-2-yl)quinolin-2(1H)-one	378.4
116	4-amino-3-[5-(4-cyclohexylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	443.1
117	4-amino-3-[6-[3-(diethylamino)pyrrolidin-1-yl]-1H-benzimidazol-2-yl]quinolin-2(1H)-one	417.1
118	4-amino-3-[6-(4-pyridin-2-ylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	438.3
119	4-amino-3-[5-(4-methylpiperazin-1-yl)-3H-imidazo[4,5-b]pyridin-2-yl]quinolin-2(1H)-one	376.3
120	4-amino-6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-imidazo[4,5-b]pyridin-2-yl]quinolin-2(1H)-one	410.2
121	2-(4-amino-2-oxo-1,2-dihydroquinolin-3-yl)-N-methyl-N-(1-methylpiperidin-4-yl)-1H-benzimidazole-5-carboxamide	431.3
122	4-amino-3-(5-[[4-(1-methylethyl)piperazin-1-yl]carbonyl]-1H-benzimidazol-2-yl)quinolin-2(1H)-one	431.3
123	4-amino-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-6-nitroquinolin-2(1H)-one	420.2
124	4-amino-3-[5-(1,4'-bipiperidin-1'-ylcarbonyl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	471.1
125	4-amino-3-[5-[(4-methylpiperazin-1-yl)carbonyl]-1H-benzimidazol-2-yl]quinolin-2(1H)-one	403.3
126	4-amino-3-[5-(1-oxidothiomorpholin-4-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	394.5
127	3-[5-[(4-acetylpiperazin-1-yl)carbonyl]-1H-benzimidazol-2-yl]-4-aminoquinolin-2(1H)-one	431.3
128	4-amino-3-(5-[[3-(3-(dimethylamino)pyrrolidin-1-yl)carbonyl]-1H-benzimidazol-2-yl]quinolin-2(1H)-one	417.4

129	4-amino-3-(5-[(3S)-3-(dimethylamino)pyrrolidin-1-yl]carbonyl)-1H-benzimidazol-2-yl)quinolin-2(1H)-one	417.4
130	4-amino-3-(5-[[4-(dimethylamino)piperidin-1-yl]carbonyl]-1H-benzimidazol-2-yl)quinolin-2(1H)-one	431.4
131	methyl 2-(4-amino-5-fluoro-2-oxo-1,2-dihydroquinolin-3-yl)-1H-benzimidazole-6-carboxylate	353.2
132	4-amino-3-[5-(1,3'-bipyrrolidin-1'-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	415.5
133	4-amino-3-[5-(pyridin-3-yloxy)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	370.2
134	4-amino-5,6-bis(methyloxy)-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	435.5
135	2-(4-amino-2-oxo-1,2-dihydroquinolin-3-yl)-N-[2-(dimethylamino)ethyl]-N-methyl-1H-benzimidazole-5-carboxamide	405.3
136	2-(4-amino-2-oxo-1,2-dihydroquinolin-3-yl)-N-methyl-N-(1-methylpyrrolidin-3-yl)-1H-benzimidazole-5-carboxamide	417.2
137	4-amino-3-{5-[(5-methyl-2,5-diazabicyclo[2.2.1]hept-2-yl)carbonyl]-1H-benzimidazol-2-yl}quinolin-2(1H)-one	415.2
138	4-amino-3-{5-[(4-cyclohexylpiperazin-1-yl)carbonyl]-1H-benzimidazol-2-yl}quinolin-2(1H)-one	471.6
139	4-amino-3-{5-[(2-piperidin-1-ylethyl)amino]-1H-benzimidazol-2-yl}quinolin-2(1H)-one	403.2
140	ethyl 4-[[2-(4-amino-2-oxo-1,2-dihydroquinolin-3-yl)-1H-benzimidazol-5-yl]amino]piperidine-1-carboxylate	447.3
141	4-amino-3-[5-[(5R)-5-[(methyloxy)methyl]pyrrolidin-3-yl]amino]-1H-benzimidazol-2-yl]quinolin-2(1H)-one	405.2
142	4-amino-3-{5-[(pyridin-2-ylmethyl)amino]-1H-benzimidazol-2-yl}quinolin-2(1H)-one	383.3
143	4-amino-3-[5-(piperidin-3-ylamino)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	375.2
144	4-amino-5-fluoro-3-{5-[(pyridin-2-ylmethyl)amino]-1H-benzimidazol-2-yl}quinolin-2(1H)-one	401.3

-141-

145	ethyl 4-[[2-(4-amino-5-fluoro-2-oxo-1,2-dihydroquinolin-3-yl)-1H-benzimidazol-5-yl]amino]piperidine-1-carboxylate	465.5
146	4-amino-5-fluoro-3-[5-(piperidin-3-ylamino)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	393.3
147	4-amino-3-(1H-benzimidazol-2-yl)-6-bromoquinolin-2(1H)-one	357.1
148	4-amino-3-(1H-benzimidazol-2-yl)-7-bromoquinolin-2(1H)-one	357.1
149	4-amino-3-(5-bromo-1H-benzimidazol-2-yl)quinolin-2(1H)-one	357.1
150	N,N-dimethyl-2-(2-oxo-1,2-dihydroquinolin-3-yl)-1H-benzimidazole-5-carboxamide	333.1
151	4-amino-3-(5-thien-2-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one	359.2
152	2-(4-amino-2-oxo-1,2-dihydroquinolin-3-yl)-N,N-dimethyl-1H-benzimidazole-5-sulfonamide	384.1
153	4-amino-6-iodo-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	501.1
154	4-amino-3-(5-{2-[(dimethylamino)methyl]-morpholin-4-yl}-1H-benzimidazol-2-yl)quinolin-2(1H)-one	419.2

Examples 155-270

Examples 155 to 270 shown in the following Table were synthesized using the methods described above such as Methods 1-15 and those set forth in the Schemes and other Examples or modified as apparent to one of reasonable skill in the art using commercially available materials.

Example	Name	LC/MS m/z (MH+)
155	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-chloro-6-iodoquinolin-2(1H)-one	547
156	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-nitroquinolin-2(1H)-one	431
157	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-methylquinolin-2(1H)-one	401
158	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6,7-difluoroquinolin-2(1H)-one	422
159	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-chloroquinolin-2(1H)-one	421
160	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-bromoquinolin-2(1H)-one	465
161	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-2-oxo-1,2-dihydroquinoline-6-carbonitrile	411
162	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoroquinolin-2(1H)-one	404
163	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6,7-bis(methoxy)quinolin-2(1H)-one	447
164	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6,7-dichloroquinolin-2(1H)-one	455
165	1-[4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-2-oxo-1,2-dihydroquinolin-7-yl]piperidine-4-carboxamide	531
166	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-7-[(3-hydroxypropyl)amino]quinolin-2(1H)-one	478
167	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-(dimethylamino)-6-fluoroquinolin-2(1H)-one	448
168	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-5-fluoroquinolin-2(1H)-one	404

169	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-(4-nitrophenyl)quinolin-2(1H)-one	508
170	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-{[2-(dimethylamino)ethyl]amino}-6-fluoroquinolin-2(1H)-one	491
171	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-7-(1H-imidazol-1-yl)quinolin-2(1H)-one	471
172	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-[4-(methyloxy)phenyl]quinolin-2(1H)-one	493
173	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-7-morpholin-4-ylquinolin-2(1H)-one	490
174	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-6,7-difluoro-3-(3H-imidazo[4,5-b]pyridin-2-yl)quinolin-2(1H)-one	423
175	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-(3-nitrophenyl)quinolin-2(1H)-one	508
176	1-[4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-2-oxo-1,2-dihydroquinolin-7-yl]piperidine-3-carboxamide	531
177	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-5-methylquinolin-2(1H)-one	401
178	6-(3-acetylphenyl)-4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(3H-imidazo[4,5-b]pyridin-2-yl)quinolin-2(1H)-one	506
179	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-5-chloroquinolin-2(1H)-one	421
180	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-6-fluoro-3-(3H-imidazo[4,5-b]pyridin-2-yl)-7-morpholin-4-ylquinolin-2(1H)-one	491
181	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-(cyclopropylamino)-6-fluoroquinolin-2(1H)-one	460
182	N-{3-[4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(3H-imidazo[4,5-b]pyridin-2-yl)-2-oxo-1,2-dihydroquinolin-6-yl]phenyl}acetamide	521

183	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-7-(4-methylpiperazin-1-yl)quinolin-2(1H)-one	503
184	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-6-fluoro-7-(1H-imidazol-1-yl)-3-(3H-imidazo[4,5-b]pyridin-2-yl)quinolin-2(1H)-one	472
185	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-7-[(2-pyridin-2-ylethyl)amino]quinolin-2(1H)-one	525
186	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-7-piperidin-1-ylquinolin-2(1H)-one	488
187	6-chloro-3-(3H-imidazo[4,5-b]pyridin-2-yl)quinolin-2(1H)-one	298
188	ethyl 1-[4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-2-oxo-1,2-dihydroquinolin-7-yl]piperidine-4-carboxylate	560
189	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-(1-benzothien-2-yl)quinolin-2(1H)-one	519
190	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-7-pyrrolidin-1-ylquinolin-2(1H)-one	474
191	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(3H-imidazo[4,5-b]pyridin-2-yl)-6-[2-(trifluoromethyl)phenyl]quinolin-2(1H)-one	532
192	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(3H-imidazo[4,5-b]pyridin-2-yl)-6-[2-(methyloxy)phenyl]quinolin-2(1H)-one	494
193	ethyl 1-[4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-2-oxo-1,2-dihydroquinolin-7-yl]piperidine-3-carboxylate	560
194	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-(4-ethylphenyl)quinolin-2(1H)-one	491
195	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-7-[(2-methylpropyl)amino]quinolin-2(1H)-one	476
196	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-5-methylquinolin-2(1H)-one	401
197	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-6-(2,4-dichlorophenyl)-3-(3H-imidazo[4,5-b]pyridin-2-yl)quinolin-2(1H)-one	532

198	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-[3-(trifluoromethyl)phenyl]quinolin-2(1H)-one	531
199	3-(1H-benzimidazol-2-yl)-4-(dimethylamino)quinolin-2(1H)-one	305
200	4-hydroxy-3-(1H-imidazo[4,5-f]quinolin-2-yl)quinolin-2(1H)-one	329
201	4-hydroxy-3-(1H-imidazo[4,5-b]pyridin-2-yl)quinolin-2(1H)-one	279
202	4-[4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-5-fluoro-2-oxo-1,2-dihydroquinolin-6-yl]benzoic acid	525
203	4-[4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-5-fluoro-2-oxo-1,2-dihydroquinolin-6-yl]benzamide	524
204	N-{3-[4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-5-fluoro-2-oxo-1,2-dihydroquinolin-6-yl]phenyl}acetamide	538
205	3-[4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-5-fluoro-2-oxo-1,2-dihydroquinolin-6-yl]benzoic acid	525
206	4-[4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-fluoro-2-oxo-1,2-dihydroquinolin-6-yl]benzoic acid	525
207	N-{3-[4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-fluoro-2-oxo-1,2-dihydroquinolin-6-yl]phenyl}acetamide	538
208	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-chloro-6-(2-methylphenyl)quinolin-2(1H)-one	511
209	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-2-oxo-1,2-dihydroquinoline-7-carbonitrile	411
210	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-(methyloxy)quinolin-2(1H)-one	417
211	4-[4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-2-oxo-1,2-dihydroquinolin-7-yl]benzamide	506
212	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-7-(methyloxy)quinolin-2(1H)-one	434
213	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-chloro-7-(dimethylamino)quinolin-2(1H)-one	464

214	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-(dimethylamino)-6-iodoquinolin-2(1H)-one	555
215	3-[4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-(1H-imidazol-1-yl)-2-oxo-1,2-dihydroquinolin-6-yl]benzoic acid	573
216	4-[4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-2-oxo-7-piperidin-1-yl-1,2-dihydroquinolin-6-yl]benzoic acid	590
217	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-(methyloxy)-6-[4-(methylsulfonyl)phenyl]quinolin-2(1H)-one	571
218	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-8-methylquinolin-2(1H)-one	401
219	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6,7-difluoroquinolin-2(1H)-one	422
220	3-(1H-benzimidazol-2-yl)-6-methyl-4-(piperidin-3-ylamino)quinolin-2(1H)-one	374
221	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-[2-(methyloxy)phenyl]quinolin-2(1H)-one	493
222	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-[3-(methyloxy)phenyl]quinolin-2(1H)-one	493
223	3-(1H-benzimidazol-2-yl)-6,7-difluoro-4-(piperidin-4-ylamino)quinolin-2(1H)-one	396
224	3-(1H-benzimidazol-2-yl)-6,7-difluoro-4-(pyrrolidin-3-ylamino)quinolin-2(1H)-one	382
225	3-(1H-benzimidazol-2-yl)-6-chloro-4-[(3-morpholin-4-ylpropyl)amino]quinolin-2(1H)-one	439
226	6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-(piperidin-4-ylamino)quinolin-2(1H)-one	480
227	6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[(piperidin-2-ylmethyl)amino]quinolin-2(1H)-one	494
228	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one	506

229	6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-(piperidin-3-ylamino)quinolin-2(1H)-one	480
230	6-chloro-4-[[2-(dimethylamino)ethyl]amino]-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one	468
231	4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one	506
232	6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[(piperidin-3-ylmethyl)amino]quinolin-2(1H)-one	494
233	6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[(piperidin-4-ylmethyl)amino]quinolin-2(1H)-one	494
234	4-[[1(1R,2R)-2-aminocyclohexyl]amino]-6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one	494
235	4-[(4-aminocyclohexyl)amino]-6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one	494
236	4-[[2(2S)-2-amino-3-methylbutyl]amino]-6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one	482
237	4-[[4-(aminomethyl)phenyl]methyl]amino]-6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one	516
238	6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[(pyrrolidin-2-ylmethyl)amino]quinolin-2(1H)-one	480
239	4-[[1(1R)-1-(aminomethyl)propyl]amino]-6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one	468
240	4-[[1(1S)-2-amino-1-(phenylmethyl)ethyl]amino]-6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one	530
241	6-chloro-4-[[3-(4-methylpiperazin-1-yl)propyl]amino]-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one	537
242	6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[[1-(phenylmethyl)piperidin-4-yl]amino]quinolin-2(1H)-one	570
243	6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[(3-morpholin-4-ylpropyl)amino]quinolin-2(1H)-one	524

244	6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[(2-piperidin-1-ylethyl)amino]quinolin-2(1H)-one	508
245	6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[(pyridin-3-ylmethyl)amino]quinolin-2(1H)-one	488
246	6-chloro-4-[[3-(1H-imidazol-1-yl)propyl]amino]-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one	505
247	6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[(pyridin-4-ylmethyl)amino]quinolin-2(1H)-one	488
248	6-chloro-4-[[2-(methylamino)ethyl]amino]-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one	454
249	6-chloro-4-[[[(2-methyl-1-piperidin-4-yl-1H-benzimidazol-5-yl)methyl]amino]-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one	624
250	6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[(2-pyrrolidin-1-ylethyl)amino]quinolin-2(1H)-one	494
251	6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-(pyrrolidin-3-ylamino)quinolin-2(1H)-one	466
252	4-[[[(1R,2R)-2-aminocyclohexyl]amino]-6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	507
253	4-[(4-aminocyclohexyl)amino]-6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	507
254	4-[[[4-(aminomethyl)phenyl]methyl]amino]-6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	529
255	6-chloro-4-[[2-(methylamino)ethyl]amino]-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	467
256	6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-[[3-(4-methylpiperazin-1-yl)propyl]amino]quinolin-2(1H)-one	550
257	6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-[[1-(phenylmethyl)piperidin-4-yl]amino]quinolin-2(1H)-one	583

258	6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-[(2-pyrrolidin-1-ylethyl)amino]quinolin-2(1H)-one	507
259	6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-(pyrrolidin-3-ylamino)quinolin-2(1H)-one	479
260	6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-(piperidin-4-ylamino)quinolin-2(1H)-one	493
261	6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[(2-piperidin-2-ylethyl)amino]quinolin-2(1H)-one	508
262	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-7-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one	506
263	7-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-(piperidin-3-ylamino)quinolin-2(1H)-one	480
264	6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-[(piperidin-2-ylmethyl)amino]quinolin-2(1H)-one	507
265	6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-[[{(2S)-pyrrolidin-2-ylmethyl}amino]quinolin-2(1H)-one	493
266	6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-[[{(2R)-pyrrolidin-2-ylmethyl}amino]quinolin-2(1H)-one	493
267	6-chloro-4-([{(2S)-1-ethylpyrrolidin-2-yl}methyl)amino)-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	521
268	6-chloro-4-([{(2R)-1-ethylpyrrolidin-2-yl}methyl)amino)-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one	521
269	4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-[4-(methyloxy)phenyl]quinolin-2(1H)-one	493
270	6-(3-aminophenyl)-4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)quinolin-2(1H)-one	478

-150-

Assay Procedures

In vitro kinase assays for receptor tyrosine kinases

The kinase activity of various protein tyrosine kinases can be measured by providing ATP and a suitable peptide or protein tyrosine-containing substrate, and assaying the transfer of phosphate moiety to the tyrosine residue. Recombinant proteins corresponding to the cytoplasmic domains of the flt-1 (VEGFR1), KDR (VEGFR2), and bFGF receptors were expressed in Sf9 insect cells using a Baculovirus expression system (InVitrogen) and purified via Glu antibody interaction (for Glu-epitope tagged constructs) or by Metal Ion Chromatography (for His₆ tagged constructs). For each assay, test compounds were serially diluted in DMSO then mixed with an appropriate kinase reaction buffer plus ATP. Kinase protein and an appropriate biotinylated peptide substrate were added to give a final volume of 100 μ L, reactions were incubated for 1-2 hours at room temperature and stopped by the addition of 50 μ L of 45 mM EDTA, 50 mM Hepes pH 7.5. Stopped reaction mix (75 μ L) was transferred to a streptavidin coated microtiter plate (Boehringer Mannheim) and incubated for 1 hour. Phosphorylated peptide product was measured with the DELFIA time-resolved fluorescence system (Wallac), using a Eu-labeled anti-phosphotyrosine antibody PT66 with the modification that the DELFIA assay buffer was supplemented with 1 mM MgCl₂ for the antibody dilution. Time resolved fluorescence was read on a Wallac 1232 DELFIA fluorometer. The concentration of each compound for 50% inhibition (IC₅₀) was calculated by non-linear regression using XL Fit data analysis software.

Flt-1, KDR, and bFGFR kinases were assayed in 50 mM Hepes pH 7.0, 2 mM MgCl₂, 10 mM MnCl₂, 1 mM NaF, 1 mM DTT, 1 mg/ml BSA, 2 μ M ATP, and 0.42 μ M biotin-GGGGQDGKDYIVLPI-NH₂. Flt-1, KDR, and bFGFR kinases were added at 0.1 μ g/mL, 0.05 μ g/mL, or 0.1 μ g/mL respectively.

Each of the following compounds was synthesized and assayed using the procedures described above:

- 3-{5-[2-(ethylamino)ethoxy]-1H-benzimidazol-2-yl}-4-hydroxy-2(1H)-quinolinone;
3-[5-(4-aminophenoxy)-1H-benzimidazol-2-yl]-4-hydroxy-2(1H)-quinolinone; 3-{6-
5 [[2-(dimethylamino)ethyl](methylamino)-1H-benzimidazol-2-yl]-4-hydroxy-2(1H)-
quinolinone; 4-hydroxy-3-[5-(4-morpholinyl)-1H-benzimidazol-2-yl]-2(1H)-
quinolinone; 3-[5-(3-amino-1-pyrrolidinyl)-1H-benzimidazol-2-yl]-4-hydroxy-2(1H)-
quinolinone; N,N-dimethyl-2-(2-oxo-1,2-dihydro-3-quinolinyl)-1H-benzimidazole-5-
carboxamide; 3-{5-[2-(4-morpholinyl)ethoxy]-1H-benzimidazol-2-yl}-2(1H)-
10 quinolinone; 3-{5-[3-(dimethylamino)-1-pyrrolidinyl]-1H-benzimidazol-2-yl}-2(1H)-
quinolinone; 3-(1H-benzimidazol-2-yl)-2-oxo-1,2-dihydro-4-quinolinecarbonitrile;
4-amino-3-{5-[2-(4-morpholinyl)ethoxy]-1H-benzimidazol-2-yl}-2(1H)-quinolinone;
4-amino-3-[6-(4-morpholinyl)-1H-benzimidazol-2-yl]-2(1H)-quinolinone; 4-amino-
3-[6-(3-amino-1-pyrrolidinyl)-1H-benzimidazol-2-yl]-2(1H)-quinolinone; 2-(4-
15 amino-2-oxo-1,2-dihydro-3-quinolinyl)-1H-benzimidazole-5-carbonitrile; 2-(4-
amino-2-oxo-1,2-dihydro-3-quinolinyl)-N,N-dimethyl-1H-benzimidazole-5-
carboxamide; 4-amino-3-{5-[3-(dimethylamino)-1-pyrrolidinyl]-1H-benzimidazol-2-
yl}-2(1H)-quinolinone; 2-(4-amino-2-oxo-1,2-dihydro-3-quinolinyl)-1H-
benzimidazole-6-carboximidamide; 4-amino-3-[5-(4-morpholinylcarbonyl)-1H-
20 benzimidazol-2-yl]-2(1H)-quinolinone; 4-amino-3-[5-(1H-1,2,4-triazol-1-yl)-1H-
benzimidazol-2-yl]-2(1H)-quinolinone; 4-amino-3-[5-(dimethylamino)-1H-
benzimidazol-2-yl]-2(1H)-quinolinone; 4-amino-3-[5-(1-piperidinyl)-1H-
benzimidazol-2-yl]-2(1H)-quinolinone; 4-amino-3-[5-(2-thienyl)-1H-benzimidazol-2-
yl]-2(1H)-quinolinone; 4-amino-3-{5-[3-(1-pyrrolidinyl)propoxy]-1H-benzimidazol-
25 2-yl}-2(1H)-quinolinone; 4-amino-3-{5-[3-(4-morpholinyl)propoxy]-1H-
benzimidazol-2-yl}-2(1H)-quinolinone; 4-amino-3-[5-(3,5-dimethyl-1-piperazinyl)-
1H-benzimidazol-2-yl]-2(1H)-quinolinone; 4-amino-3-[5-(2,6-dimethyl-4-
morpholinyl)-1H-benzimidazol-2-yl]-2(1H)-quinolinone; 4-amino-3-[5-(4-methyl-1-
piperazinyl)-1H-benzimidazol-2-yl]-2(1H)-quinolinone; 4-amino-3-(1H-
30 benzimidazol-2-yl)-6-[hydroxy(oxido)amino]-2(1H)-quinolinone; 4-amino-3-(1H-
benzimidazol-2-yl)-5-[2-(4-morpholinyl)ethoxy]-2(1H)-quinolinone; 4-amino-3-(1H-

- benzimidazol-2-yl)-6-(4-methyl-1-piperazinyl)-2(1H)-quinolinone; 4-amino-3-(1H-benzimidazol-2-yl)-5-[(1-methyl-3-piperidinyl)oxy]-2(1H)-quinolinone; 4-amino-6-chloro-3-[5-(4-morpholinyl)-1H-benzimidazol-2-yl]-2(1H)-quinolinone; 4-amino-6-chloro-3-{5-[3-(dimethylamino)-1-pyrrolidinyl]-1H-benzimidazol-2-yl}-2(1H)-quinolinone; 4-amino-6-[hydroxy(oxido)amino]-3-{5-[2-(4-morpholinyl)ethoxy]-1H-benzimidazol-2-yl}-2(1H)-quinolinone; 4-amino-5-[2-(4-morpholinyl)ethoxy]-3-{5-[2-(4-morpholinyl)ethoxy]-1H-benzimidazol-2-yl}-2(1H)-quinolinone; 4-amino-3-(1H-benzimidazol-2-yl)-6-(2-pyridinylmethoxy)-2(1H)-quinolinone; 4-amino-6-fluoro-3-[5-(4-morpholinyl)-1H-benzimidazol-2-yl]-2(1H)-quinolinone; 4-amino-3-{5-[3-(dimethylamino)-1-pyrrolidinyl]-1H-benzimidazol-2-yl}-6-fluoro-2(1H)-quinolinone; 3-(1H-benzimidazol-2-yl)-4-[(tetrahydro-2-furanylmethyl)amino]-2(1H)-quinolinone; 3-(1H-benzimidazol-2-yl)-4-(methylamino)-2(1H)-quinolinone; 3-(1H-benzimidazol-2-yl)-4-(ethylamino)-2(1H)-quinolinone; 3-(1H-benzimidazol-2-yl)-4-{[2-(1-methyl-2-pyrrolidinyl)ethyl]amino}-2(1H)-quinolinone; 3-(1H-benzimidazol-2-yl)-4-[(4-piperidinylmethyl)amino]-2(1H)-quinolinone; 3-(1H-benzimidazol-2-yl)-4-(4-fluoroanilino)-2(1H)-quinolinone; 4-(1-azabicyclo[2.2.2]oct-3-ylamino)-3-(1H-benzimidazol-2-yl)-2(1H)-quinolinone; 3-(1H-benzimidazol-2-yl)-4-(1H-benzimidazol-6-ylamino)-2(1H)-quinolinone; 4-anilino-3-(1H-benzimidazol-2-yl)-2(1H)-quinolinone; 3-(1H-benzimidazol-2-yl)-4-(methoxyamino)-2(1H)-quinolinone; 3-(1H-benzimidazol-2-yl)-4-[(1H-imidazol-5-ylmethyl)amino]-2(1H)-quinolinone; 3-(1H-benzimidazol-2-yl)-4-(4-morpholinylamino)-2(1H)-quinolinone; 3-(1H-benzimidazol-2-yl)-4-hydrazino-2(1H)-quinolinone; 4-(1-azabicyclo[2.2.2]oct-3-ylamino)-3-(1H-benzimidazol-2-yl)-2(1H)-quinolinone; 4-(1-azabicyclo[2.2.2]oct-3-ylamino)-3-(1H-benzimidazol-2-yl)-2(1H)-quinolinone; 4-[(2-methoxyethyl)amino]-3-[6-(4-morpholinyl)-1H-benzimidazol-2-yl]-2(1H)-quinolinone; 4-[(2-hydroxyethyl)amino]-3-[5-(4-morpholinyl)-1H-benzimidazol-2-yl]-2(1H)-quinolinone; 4-(methoxyamino)-3-[5-(4-morpholinyl)-1H-benzimidazol-2-yl]-2(1H)-quinolinone; 3-[5-(4-morpholinyl)-1H-benzimidazol-2-yl]-4-(3-piperidinylamino)-2(1H)-quinolinone; 3-[5-(4-morpholinyl)-1H-benzimidazol-2-yl]-4-[(3-piperidinylmethyl)amino]-2(1H)-quinolinone; 4-{[2-(dimethylamino)ethyl]amino}-3-[5-(4-morpholinyl)-1H-benzimidazol-2-yl]-2(1H)-

- quinolinone; 3-[5-(4-morpholinyl)-1H-benzimidazol-2-yl]-4-[(tetrahydro-2-furanylmethyl)amino]-2(1H)-quinolinone; 4-{[2-(methylamino)ethyl]amino}-3-[5-(4-morpholinyl)-1H-benzimidazol-2-yl]-2(1H)-quinolinone; 3-[5-(4-morpholinyl)-1H-benzimidazol-2-yl]-4-(3-pyrrolidinylamino)-2(1H)-quinolinone; 4-[(2-amino-4-methylpentyl)amino]-3-[5-(4-morpholinyl)-1H-benzimidazol-2-yl]-2(1H)-quinolinone; 4-[(2-amino-3-methylbutyl)amino]-3-[5-(4-morpholinyl)-1H-benzimidazol-2-yl]-2(1H)-quinolinone; 3-(5,6-dimethyl-1H-benzimidazol-2-yl)-4-(3-piperidinylamino)-2(1H)-quinolinone; 4-[(2-aminocyclohexyl)amino]-3-[5-(4-morpholinyl)-1H-benzimidazol-2-yl]-2(1H)-quinolinone; 4-[(2-aminocyclohexyl)amino]-3-[5-(4-morpholinyl)-1H-benzimidazol-2-yl]-2(1H)-quinolinone; 3-(1H-benzimidazol-2-yl)-4-hydroxybenzo[g]quinolin-2(1H)-one; 4-amino-3-(3H-imidazo[4,5-b]pyridin-2-yl)quinolin-2(1H)-one; 4-amino-3-(5-morpholin-4-yl-3H-imidazo[4,5-b]pyridin-2-yl)quinolin-2(1H)-one; 4-amino-5-[(2R,6S)-2,6-dimethylmorpholin-4-yl]-3-(3H-imidazo[4,5-b]pyridin-2-yl)quinolin-2(1H)-one; 4-amino-3-{5-[3-(dimethylamino)pyrrolidin-1-yl]-3H-imidazo[4,5-b]pyridin-2-yl}quinolin-2(1H)-one; 4-amino-3-{5-[(3S)-3-(dimethylamino)pyrrolidin-1-yl]-1H-benzimidazol-2-yl}quinolin-2(1H)-one; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-chloroquinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-chloroquinolin-2(1H)-one; 3-(1H-benzimidazol-2-yl)-4-[(3R)-3-(dimethylamino)pyrrolidin-1-yl]quinolin-2(1H)-one; 3-(1H-benzimidazol-2-yl)-6-chloro-4-[(3R)-3-(dimethylamino)pyrrolidin-1-yl]quinolin-2(1H)-one; 4-amino-3-[5-(4-ethylpiperazin-1-yl)-1H-benzimidazol-2-yl]-1-methylquinolin-2(1H)-one; 4-amino-3-(6-piperazin-1-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 4-amino-3-[6-(pyridin-4-ylmethyl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-amino-3-{5-[(3R,5S)-3,5-dimethylpiperazin-1-yl]-1H-benzimidazol-2-yl}quinolin-2(1H)-one; 4-amino-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-amino-3-(6-methyl-5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 4-amino-3-{5-[(1-methylpiperidin-3-yl)oxy]-1H-benzimidazol-2-yl}quinolin-2(1H)-one; 4-amino-3-{5-[(2R,6S)-2,6-dimethylmorpholin-4-yl]-6-fluoro-1H-benzimidazol-2-yl}quinolin-2(1H)-one; 4-amino-3-{5-[(1-methylpyrrolidin-3-

- yl)oxy]-1H-benzimidazol-2-yl}quinolin-2(1H)-one; 4-amino-3-[5-(4-methyl-1,4-diazepan-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-amino-3-{5-[(3R)-3-(dimethylamino)pyrrolidin-1-yl]-1H-benzimidazol-2-yl}quinolin-2(1H)-one; 4-amino-6-chloro-3-{5-[(3R)-3-(dimethylamino)pyrrolidin-1-yl]-1H-benzimidazol-2-yl}quinolin-2(1H)-one; ethyl {4-[2-(4-amino-2-oxo-1,2-dihydroquinolin-3-yl)-1H-benzimidazol-6-yl]piperazin-1-yl}acetate; 4-amino-3-{6-[methyl(1-methylpiperidin-4-yl)amino]-1H-benzimidazol-2-yl}quinolin-2(1H)-one; 3-[6-(4-acetylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-aminoquinolin-2(1H)-one; 4-amino-3-[6-(1,4'-bipiperidin-1'-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 2-(4-amino-2-oxo-1,2-dihydroquinolin-3-yl)-1H-benzimidazole-6-carboxylic acid; 4-amino-5-(methyloxy)-3-[6-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-amino-3-{6-[4-(1-methylethyl)piperazin-1-yl]-1H-benzimidazol-2-yl}quinolin-2(1H)-one; {4-[2-(4-amino-2-oxo-1,2-dihydroquinolin-3-yl)-1H-benzimidazol-6-yl]piperazin-1-yl}acetic acid; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)quinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)quinolin-2(1H)-one; 4-amino-3-[5-(4-ethylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-amino-3-(5-{(2S,5S)-2-[(dimethylamino)methyl]-5-methylmorpholin-4-yl}-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 4-amino-6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-amino-6-chloro-3-{5-[(3S)-3-(dimethylamino)pyrrolidin-1-yl]-1H-benzimidazol-2-yl}quinolin-2(1H)-one; 4-amino-5,6-dichloro-3-{5-[(3S)-3-(dimethylamino)pyrrolidin-1-yl]-1H-benzimidazol-2-yl}quinolin-2(1H)-one; 4-amino-5,6-dichloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-amino-3-(1H-benzimidazol-2-yl)-6-[(pyridin-2-ylmethyl)oxy]quinolin-2(1H)-one; 4-amino-3-(1H-benzimidazol-2-yl)-6-[(2R,6S)-2,6-dimethylmorpholin-4-yl]quinolin-2(1H)-one; 4-amino-3-(1H-benzimidazol-2-yl)-6-morpholin-4-ylquinolin-2(1H)-one; 4-amino-3-(1H-benzimidazol-2-yl)-5-[(1-methylpiperidin-3-yl)oxy]quinolin-2(1H)-one; 4-amino-3-(1H-benzimidazol-2-yl)-5-[(pyridin-2-ylmethyl)oxy]quinolin-2(1H)-one; 4-amino-3-(5-morpholin-4-yl)-1H-benzimidazol-2-yl)-5-[(pyridin-4-ylmethyl)oxy]quinolin-2(1H)-one; 4-amino-3-(1H-benzimidazol-2-yl)-5-(methyloxy)quinolin-2(1H)-one; 4-amino-3-(5-methyl-1H-benzimidazol-2-yl)-

- 5-(methyloxy)quinolin-2(1H)-one; 4-amino-3-{5-[(2R,6S)-2,6-dimethylmorpholin-4-yl]-1H-benzimidazol-2-yl}-5-(methyloxy)quinolin-2(1H)-one; 4-amino-3-(1H-benzimidazol-2-yl)-5-morpholin-4-ylquinolin-2(1H)-one; 4-amino-3-(1H-benzimidazol-2-yl)-5-[(2R,6S)-2,6-dimethylmorpholin-4-yl]quinolin-2(1H)-one; 4-amino-3-(1H-benzimidazol-2-yl)-5-(4-methylpiperazin-1-yl)quinolin-2(1H)-one; 4-amino-5,6-dichloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 3-{5-[(2-morpholin-4-ylethyl)oxy]-1H-benzimidazol-2-yl}quinolin-2(1H)-one; 4-amino-3-{5-[(3-pyrrolidin-1-ylpropyl)oxy]-1H-benzimidazol-2-yl}quinolin-2(1H)-one; 4-amino-3-{5-[(3-morpholin-4-ylpropyl)oxy]-1H-benzimidazol-2-yl}quinolin-2(1H)-one; 4-amino-6-fluoro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 4-amino-3-{5-[3-(dimethylamino)pyrrolidin-1-yl]-1H-benzimidazol-2-yl}-6-fluoroquinolin-2(1H)-one; 4-amino-3-(1H-benzimidazol-2-yl)-6-fluoroquinolin-2(1H)-one; 4-amino-3-(6-fluoro-5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 4-amino-3-{5-[(tetrahydrofuran-2-ylmethyl)oxy]-1H-benzimidazol-2-yl}quinolin-2(1H)-one; 4-amino-6-fluoro-3-(6-fluoro-5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 4-amino-3-[6-fluoro-5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-amino-3-(5-{[2-(methyloxy)ethyl]oxy}-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 4-amino-3-[4,6-difluoro-5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-amino-3-{5-[3-(dimethylamino)pyrrolidin-1-yl]-1H-benzimidazol-2-yl}-5-fluoroquinolin-2(1H)-one; 4-amino-5-fluoro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-amino-5-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-amino-3-{5-[3-(dimethylamino)pyrrolidin-1-yl]-6-fluoro-1H-benzimidazol-2-yl}quinolin-2(1H)-one; 4-amino-5-chloro-3-{5-[3-(dimethylamino)pyrrolidin-1-yl]-1H-benzimidazol-2-yl}quinolin-2(1H)-one; 4-amino-6-chloro-3-{5-[3-(dimethylamino)pyrrolidin-1-yl]-6-fluoro-1H-benzimidazol-2-yl}quinolin-2(1H)-one; 4-amino-5-[(2R,6S)-2,6-dimethylmorpholin-4-yl]-3-(3H-imidazo[4,5-b]pyridin-2-yl)quinolin-2(1H)-one; 4-amino-3-(6-thiomorpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 4-amino-3-[5-(4-cyclohexylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-amino-3-{6-[3-(diethylamino)pyrrolidin-1-yl]-1H-benzimidazol-2-yl}quinolin-2(1H)-

- one; 4-amino-3-[6-(4-pyridin-2-ylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-amino-3-[5-(4-methylpiperazin-1-yl)-3H-imidazo[4,5-b]pyridin-2-yl]quinolin-2(1H)-one; 4-amino-6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-imidazo[4,5-b]pyridin-2-yl]quinolin-2(1H)-one; 2-(4-amino-2-oxo-1,2-dihydroquinolin-3-yl)-N-methyl-N-(1-methylpiperidin-4-yl)-1H-benzimidazole-5-carboxamide; 4-amino-3-(5-{[4-(1-methylethyl)piperazin-1-yl]carbonyl}-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 4-amino-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-6-nitroquinolin-2(1H)-one; 4-amino-3-[5-(1,4'-bipiperidin-1'-ylcarbonyl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-amino-3-{5-[(4-methylpiperazin-1-yl)carbonyl]-1H-benzimidazol-2-yl}quinolin-2(1H)-one; 4-amino-3-[5-(1-oxidothiomorpholin-4-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 3-{5-[(4-acetylpiperazin-1-yl)carbonyl]-1H-benzimidazol-2-yl}-4-aminoquinolin-2(1H)-one; 4-amino-3-(5-{[(3R)-3-(dimethylamino)pyrrolidin-1-yl]carbonyl}-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 4-amino-3-(5-{[(3S)-3-(dimethylamino)pyrrolidin-1-yl]carbonyl}-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 4-amino-3-(5-{[4-(dimethylamino)piperidin-1-yl]carbonyl}-1H-benzimidazol-2-yl)quinolin-2(1H)-one; methyl 2-(4-amino-5-fluoro-2-oxo-1,2-dihydroquinolin-3-yl)-1H-benzimidazole-6-carboxylate; 4-amino-3-[5-(1,3'-bipyrrolidin-1'-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-amino-3-[5-(pyridin-3-yloxy)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-amino-5,6-bis(methyloxy)-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 2-(4-amino-2-oxo-1,2-dihydroquinolin-3-yl)-N-[2-(dimethylamino)ethyl]-N-methyl-1H-benzimidazole-5-carboxamide; 2-(4-amino-2-oxo-1,2-dihydroquinolin-3-yl)-N-methyl-N-(1-methylpyrrolidin-3-yl)-1H-benzimidazole-5-carboxamide; 4-amino-3-{5-[(5-methyl-2,5-diazabicyclo[2.2.1]hept-2-yl)carbonyl]-1H-benzimidazol-2-yl}quinolin-2(1H)-one; 4-amino-3-{5-[(4-cyclohexylpiperazin-1-yl)carbonyl]-1H-benzimidazol-2-yl}quinolin-2(1H)-one; 4-amino-3-{5-[(2-piperidin-1-ylethyl)amino]-1H-benzimidazol-2-yl}quinolin-2(1H)-one; ethyl 4-{[2-(4-amino-2-oxo-1,2-dihydroquinolin-3-yl)-1H-benzimidazol-5-yl]amino}piperidine-1-carboxylate; 4-amino-3-[5-({(5R)-5-[(methyloxy)methyl]pyrrolidin-3-yl}amino)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-amino-3-{5-[(pyridin-2-ylmethyl)amino]-1H-

- benzimidazol-2-yl}quinolin-2(1H)-one; 4-amino-3-[5-(piperidin-3-ylamino)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-amino-5-fluoro-3-{5-[(pyridin-2-ylmethyl)amino]-1H-benzimidazol-2-yl}quinolin-2(1H)-one; ethyl 4-{[2-(4-amino-5-fluoro-2-oxo-1,2-dihydroquinolin-3-yl)-1H-benzimidazol-5-yl]amino}piperidine-1-carboxylate; 4-amino-5-fluoro-3-[5-(piperidin-3-ylamino)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-amino-3-(1H-benzimidazol-2-yl)-6-bromoquinolin-2(1H)-one; 4-amino-3-(1H-benzimidazol-2-yl)-7-bromoquinolin-2(1H)-one; 4-amino-3-(5-bromo-1H-benzimidazol-2-yl)quinolin-2(1H)-one; N,N-dimethyl-2-(2-oxo-1,2-dihydroquinolin-3-yl)-1H-benzimidazole-5-carboxamide; 4-amino-3-(5-thien-2-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 2-(4-amino-2-oxo-1,2-dihydroquinolin-3-yl)-N,N-dimethyl-1H-benzimidazole-5-sulfonamide; 4-amino-6-iodo-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-amino-3-(5-{2-[(dimethylamino)methyl]morpholin-4-yl}-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-chloro-6-iodoquinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-nitroquinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-methylquinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6,7-difluoroquinolin-2(1H)-one; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-chloroquinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-bromoquinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-2-oxo-1,2-dihydroquinoline-6-carbonitrile; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoroquinolin-2(1H)-one; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6,7-bis(methyloxy)quinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6,7-dichloroquinolin-2(1H)-one; 1-[4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-2-oxo-1,2-dihydroquinolin-7-yl]piperidine-4-carboxamide; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-7-[(3-hydroxypropyl)amino]quinolin-2(1H)-one; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-(dimethylamino)-6-fluoroquinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-

- ylamino]-3-(1H-benzimidazol-2-yl)-5-fluoroquinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-(4-nitrophenyl)quinolin-2(1H)-one; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-{[2-(dimethylamino)ethyl]amino}-6-fluoroquinolin-2(1H)-one;
- 5 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-7-(1H-imidazol-1-yl)quinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-[4-(methyloxy)phenyl]quinolin-2(1H)-one; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-7-morpholin-4-ylquinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-6,7-difluoro-3-
- 10 (3H-imidazo[4,5-b]pyridin-2-yl)quinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-(3-nitrophenyl)quinolin-2(1H)-one; 1-[4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-2-oxo-1,2-dihydroquinolin-7-yl]piperidine-3-carboxamide; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-5-methylquinolin-2(1H)-one; 6-(3-
- 15 acetylphenyl)-4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(3H-imidazo[4,5-b]pyridin-2-yl)quinolin-2(1H)-one; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-5-chloroquinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-6-fluoro-3-(3H-imidazo[4,5-b]pyridin-2-yl)-7-morpholin-4-ylquinolin-2(1H)-one; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-
- 20 (1H-benzimidazol-2-yl)-7-(cyclopropylamino)-6-fluoroquinolin-2(1H)-one; N-{3-[4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(3H-imidazo[4,5-b]pyridin-2-yl)-2-oxo-1,2-dihydroquinolin-6-yl]phenyl}acetamide; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-7-(4-methylpiperazin-1-yl)quinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-6-fluoro-7-(1H-imidazol-1-
- 25 yl)-3-(3H-imidazo[4,5-b]pyridin-2-yl)quinolin-2(1H)-one; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-7-[(2-pyridin-2-ylethyl)amino]quinolin-2(1H)-one; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-7-piperidin-1-ylquinolin-2(1H)-one; 6-chloro-3-(3H-imidazo[4,5-b]pyridin-2-yl)quinolin-2(1H)-one; ethyl 1-[4-[(3S)-1-
- 30 azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-2-oxo-1,2-dihydroquinolin-7-yl]piperidine-4-carboxylate; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-

- ylamino]-3-(1H-benzimidazol-2-yl)-6-(1-benzothien-2-yl)quinolin-2(1H)-one; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-7-pyrrolidin-1-ylquinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(3H-imidazo[4,5-b]pyridin-2-yl)-6-[2-(trifluoromethyl)phenyl]quinolin-2(1H)-one;
- 5 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(3H-imidazo[4,5-b]pyridin-2-yl)-6-[2-(methyloxy)phenyl]quinolin-2(1H)-one; ethyl 1-[4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-2-oxo-1,2-dihydroquinolin-7-yl]piperidine-3-carboxylate; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-(4-ethylphenyl)quinolin-2(1H)-one; 4-[(3S)-1-
- 10 azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-7-[(2-methylpropyl)amino]quinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-5-methylquinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-6-(2,4-dichlorophenyl)-3-(3H-imidazo[4,5-b]pyridin-2-yl)quinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-
- 15 (1H-benzimidazol-2-yl)-6-[3-(trifluoromethyl)phenyl]quinolin-2(1H)-one; 3-(1H-benzimidazol-2-yl)-4-(dimethylamino)quinolin-2(1H)-one; 4-hydroxy-3-(1H-imidazo[4,5-f]quinolin-2-yl)quinolin-2(1H)-one; 4-hydroxy-3-(1H-imidazo[4,5-b]pyridin-2-yl)quinolin-2(1H)-one; 4-[4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-5-fluoro-2-oxo-1,2-dihydroquinolin-6-yl]benzoic acid; 4-[4-
- 20 [(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-5-fluoro-2-oxo-1,2-dihydroquinolin-6-yl]benzamide; N-{3-[4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-5-fluoro-2-oxo-1,2-dihydroquinolin-6-yl]phenyl}acetamide; 3-[4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-5-fluoro-2-oxo-1,2-dihydroquinolin-6-yl]benzoic acid; 4-[4-
- 25 [(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-fluoro-2-oxo-1,2-dihydroquinolin-6-yl]benzoic acid; N-{3-[4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-fluoro-2-oxo-1,2-dihydroquinolin-6-yl]phenyl}acetamide; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-chloro-6-(2-methylphenyl)quinolin-2(1H)-one; 4-[(3R)-1-
- 30 azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-2-oxo-1,2-dihydroquinoline-7-carbonitrile; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-

- benzimidazol-2-yl)-7-(methyloxy)quinolin-2(1H)-one; 4-[4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-2-oxo-1,2-dihydroquinolin-7-yl]benzamide; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-fluoro-7-(methyloxy)quinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-chloro-7-(dimethylamino)quinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-(dimethylamino)-6-iodoquinolin-2(1H)-one; 3-[4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-(1H-imidazol-1-yl)-2-oxo-1,2-dihydroquinolin-6-yl]benzoic acid; 4-[4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-2-oxo-7-piperidin-1-yl-1,2-dihydroquinolin-6-yl]benzoic acid; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-7-(methyloxy)-6-[4-(methylsulfonyl)phenyl]quinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-8-methylquinolin-2(1H)-one; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6,7-difluoroquinolin-2(1H)-one; 3-(1H-benzimidazol-2-yl)-6-methyl-4-(piperidin-3-ylamino)quinolin-2(1H)-one; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-[2-(methyloxy)phenyl]quinolin-2(1H)-one; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-[3-(methyloxy)phenyl]quinolin-2(1H)-one; 3-(1H-benzimidazol-2-yl)-6,7-difluoro-4-(piperidin-4-ylamino)quinolin-2(1H)-one; 3-(1H-benzimidazol-2-yl)-6,7-difluoro-4-(pyrrolidin-3-ylamino)quinolin-2(1H)-one; 3-(1H-benzimidazol-2-yl)-6-chloro-4-[(3-morpholin-4-ylpropyl)amino]quinolin-2(1H)-one; 6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-(piperidin-4-ylamino)quinolin-2(1H)-one; 6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[(piperidin-2-ylmethyl)amino]quinolin-2(1H)-one; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-(piperidin-3-ylamino)quinolin-2(1H)-one; 6-chloro-4-{[2-(dimethylamino)ethyl]amino}-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 4-[(3R)-1-azabicyclo[2.2.2]oct-3-ylamino]-6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[(piperidin-3-ylmethyl)amino]quinolin-2(1H)-one; 6-chloro-3-

(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[(piperidin-4-ylmethyl)amino]quinolin-2(1H)-one; 4-[[[(1R,2R)-2-aminocyclohexyl]amino]-6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 4-[(4-aminocyclohexyl)amino]-6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 4-[[[(2S)-2-amino-3-methylbutyl]amino]-6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 4-({[4-(aminomethyl)phenyl]methyl}amino)-6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[(pyrrolidin-2-ylmethyl)amino]quinolin-2(1H)-one; 4-[[[(1R)-1-(aminomethyl)propyl]amino]-6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 4-[[[(1S)-2-amino-1-(phenylmethyl)ethyl]amino]-6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 6-chloro-4-[[3-(4-methylpiperazin-1-yl)propyl]amino]-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[[1-(phenylmethyl)piperidin-4-yl]amino]quinolin-2(1H)-one; 6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[(3-morpholin-4-ylpropyl)amino]quinolin-2(1H)-one; 6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[(2-piperidin-1-ylethyl)amino]quinolin-2(1H)-one; 6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[(pyridin-3-ylmethyl)amino]quinolin-2(1H)-one; 6-chloro-4-[[3-(1H-imidazol-1-yl)propyl]amino]-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[(pyridin-4-ylmethyl)amino]quinolin-2(1H)-one; 6-chloro-4-[[2-(methylamino)ethyl]amino]-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 6-chloro-4-[[2-methyl-1-piperidin-4-yl-1H-benzimidazol-5-yl)methyl]amino]-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)quinolin-2(1H)-one; 6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-[(2-pyrrolidin-1-ylethyl)amino]quinolin-2(1H)-one; 6-chloro-3-(5-morpholin-4-yl-1H-benzimidazol-2-yl)-4-(pyrrolidin-3-ylamino)quinolin-2(1H)-one; 4-[[[(1R,2R)-2-aminocyclohexyl]amino]-6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-[(4-aminocyclohexyl)amino]-6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-({[4-(aminomethyl)phenyl]methyl}amino)-6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 6-chloro-4-[[2-

(methylamino)ethyl]amino}-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-{[3-(4-methylpiperazin-1-yl)propyl]amino}quinolin-2(1H)-one; 6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-{[1-(phenylmethyl)piperidin-4-yl]amino}quinolin-2(1H)-one; 6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-[(2-pyrrolidin-1-ylethyl)amino]quinolin-2(1H)-one; 6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-(pyrrolidin-3-ylamino)quinolin-2(1H)-one; 6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-(piperidin-4-ylamino)quinolin-2(1H)-one; 6-chloro-3-(5-morpholin-4-yl)-1H-benzimidazol-2-yl]-4-[(2-piperidin-2-ylethyl)amino]quinolin-2(1H)-one; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-7-chloro-3-(5-morpholin-4-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 7-chloro-3-(5-morpholin-4-yl)-1H-benzimidazol-2-yl]-4-(piperidin-3-ylamino)quinolin-2(1H)-one; 6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-[(piperidin-2-ylmethyl)amino]quinolin-2(1H)-one; 6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-{[(2S)-pyrrolidin-2-ylmethyl]amino}quinolin-2(1H)-one; 6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-{[(2R)-pyrrolidin-2-ylmethyl]amino}quinolin-2(1H)-one; 6-chloro-4-({[(2S)-1-ethylpyrrolidin-2-yl]methyl}amino)-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 6-chloro-4-({[(2R)-1-ethylpyrrolidin-2-yl]methyl}amino)-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one; 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-[4-(methyloxy)phenyl]quinolin-2(1H)-one; and 6-(3-aminophenyl)-4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)quinolin-2(1H)-one.

Each of the above compounds displayed an IC₅₀ value of less than 10 μM with respect to VEGFR1, VEGFR2, and bFGF.

It should be understood that the organic compounds according to the invention may exhibit the phenomenon of tautomerism. As the chemical structures within this specification can only represent one of the possible tautomeric forms, it

-163-

should be understood that the invention encompasses any tautomeric form of the drawn structure.


It is understood that the invention is not limited to the embodiments set forth herein for illustration, but embraces all such forms thereof as come within
5 the scope of the following claims.

-164-

CLAIMS

What is claimed is:

- 1 1. A compound having the structure I, a tautomer of the
 2 compound, a pharmaceutically acceptable salt of the compound, or a
 3 pharmaceutically acceptable salt of the tautomer



I

4

5

wherein,

- 6 Y is selected from the group consisting of $-OR^{10}$ groups, $-C(=O)-R^{11}$
 7 groups, $-NR^{12}R^{13}$ groups, substituted and unsubstituted alkynyl
 8 groups, substituted and unsubstituted heterocyclalkyl groups,
 9 substituted and unsubstituted alkylaminoalkyl groups, substituted and
 10 unsubstituted dialkylaminoalkyl groups, substituted and unsubstituted
 11 arylaminoalkyl groups, substituted and unsubstituted
 12 diarylaminoalkyl groups, substituted and unsubstituted
 13 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
 14 heterocyclaminoalkyl groups, substituted and unsubstituted
 15 saturated heterocycl groups, substituted and unsubstituted
 16 heterocycloxyalkyl groups, substituted and unsubstituted
 17 hydroxyalkyl groups, and substituted and unsubstituted aryloxyalkyl
 18 groups;

- 19 Z is selected from the group consisting of O, S, and NR^{14} groups;
- 20 R^1 , R^2 , R^3 , and R^4 may be the same or different and are
21 independently selected from the group consisting of H, Cl, Br, F, I,
22 -CN, -NO₂, -OH, -OR¹⁵ groups, -NR¹⁶R¹⁷ groups, substituted and
23 unsubstituted amidinyl groups, substituted and unsubstituted
24 guanidinyll groups, substituted and unsubstituted primary, secondary,
25 and tertiary alkyl groups, substituted and unsubstituted aryl groups,
26 substituted and unsubstituted alkenyl groups, substituted and
27 unsubstituted alkynyl groups, substituted and unsubstituted
28 heterocyclyl groups, substituted and unsubstituted aminoalkyl groups,
29 substituted and unsubstituted alkylaminoalkyl groups, substituted and
30 unsubstituted dialkylaminoalkyl groups, substituted and unsubstituted
31 arylaminoalkyl groups, substituted and unsubstituted
32 diarylaminoalkyl groups, substituted and unsubstituted
33 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
34 heterocyclylalkyl groups, substituted and unsubstituted
35 diheterocyclylaminoalkyl groups, substituted and unsubstituted
36 (heterocyclyl)(alkyl)aminoalkyl groups, substituted and unsubstituted
37 (heterocyclyl)(aryl)aminoalkyl groups, and -C(=O)R¹⁸ groups;
- 38 R^5 , R^6 , R^7 , and R^8 may be the same or different and are
39 independently selected from the group consisting of H, Cl, Br, F, I,
40 -NO₂, -OH, -OR¹⁹ groups, -NR²⁰R²¹ groups, -SH, -SR²² groups,
41 -S(=O)R²³ groups, -S(=O)₂R²⁴ groups, -CN, substituted and
42 unsubstituted amidinyl groups, substituted and unsubstituted
43 guanidinyll groups, substituted and unsubstituted primary, secondary,
44 and tertiary alkyl groups, substituted and unsubstituted aryl groups,
45 substituted and unsubstituted alkenyl groups, substituted and
46 unsubstituted alkynyl groups, substituted and unsubstituted
47 heterocyclyl groups, substituted and unsubstituted heterocyclylalkyl

48 groups, $-C(=O)R^{25}$ groups, substituted and unsubstituted aminoalkyl
49 groups, substituted and unsubstituted alkylaminoalkyl groups,
50 substituted and unsubstituted dialkylaminoalkyl groups, substituted
51 and unsubstituted arylaminoalkyl groups, substituted and
52 unsubstituted diarylaminoalkyl groups, substituted and unsubstituted
53 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
54 heterocyclylaminoalkyl groups, substituted and unsubstituted
55 diheterocyclylaminoalkyl groups, substituted and unsubstituted
56 (heterocyclyl)(alkyl)aminoalkyl groups, substituted and unsubstituted
57 (heterocyclyl)(aryl)aminoalkyl groups, substituted and unsubstituted
58 hydroxyalkyl groups, substituted and unsubstituted alkoxyalkyl
59 groups, substituted and unsubstituted aryloxyalkyl groups, and
60 substituted and unsubstituted heterocycloxyalkyl groups;

61 R^9 and R^{14} may be the same or different and are independently
62 selected from the group consisting of H, -OH, substituted and
63 unsubstituted alkoxy groups, substituted and unsubstituted aryloxy
64 groups, $-NH_2$, substituted and unsubstituted alkylamino groups,
65 substituted and unsubstituted arylamino groups, substituted and
66 unsubstituted dialkylamino groups, substituted and unsubstituted
67 diarylamino groups, substituted and unsubstituted (alkyl)(aryl)amino
68 groups, substituted and unsubstituted alkyl groups, substituted and
69 unsubstituted aryl groups, $-C(=O)H$, $-C(=O)$ -alkyl groups, and
70 $-C(=O)$ -aryl groups;

71 R^{10} is selected from the group consisting of substituted and
72 unsubstituted aryl groups, substituted and unsubstituted heterocyclyl
73 groups, $-C(=O)H$, $-C(=O)$ -alkyl groups, $-C(=O)$ -aryl groups,
74 $-C(=O)O$ -alkyl groups, $-C(=O)O$ -aryl groups, $-C(=O)NH_2$,
75 $-C(=O)NH$ (alkyl) groups, $-C(=O)NH$ (aryl) groups,
76 $-C(=O)N$ (alkyl)₂ groups, $-C(=O)N$ (aryl)₂ groups,

- 77 -C(=O)N(alkyl)(aryl) groups, -NH₂, -NH(alkyl) groups, -NH(aryl)
78 groups, -N(alkyl)₂ groups, -N(alkyl)(aryl) groups, -N(aryl)₂ groups,
79 -NH(heterocyclyl) groups, -N(heterocyclyl)₂ groups,
80 -N(alkyl)(heterocyclyl) groups, -N(aryl)(heterocyclyl),
81 -C(=O)NH(heterocyclyl) groups, -C(=O)N(heterocyclyl)₂ groups,
82 -C(=O)N(alkyl)(heterocyclyl) groups, -C(=O)N(aryl)(heterocyclyl)
83 groups, and substituted and unsubstituted heterocyclalkyl groups;
- 84 R¹¹ is selected from the group consisting of H, -NH₂, -NH(alkyl)
85 groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(aryl)₂ groups,
86 -N(alkyl)(aryl) groups, -NH(heterocyclyl) groups, -N(heterocyclyl)₂
87 groups, -N(alkyl)(heterocyclyl) groups, -N(aryl)(heterocyclyl)
88 groups, -O-alkyl groups, O-aryl groups, heterocyclloxyalkyl
89 groups, and substituted and unsubstituted aryl groups;
- 90 R¹² is selected from the group consisting of H, substituted and
91 unsubstituted alkyl groups, substituted and unsubstituted aryl groups,
92 and substituted and unsubstituted heterocyclyl groups;
- 93 R¹³ is selected from the group consisting of substituted and
94 unsubstituted alkyl groups, substituted and unsubstituted aryl groups,
95 substituted and unsubstituted heterocyclyl groups, -OH, alkoxy
96 groups, aryloxy groups, -NH₂, substituted and unsubstituted
97 heterocyclalkyl groups, substituted and unsubstituted aminoalkyl
98 groups, substituted and unsubstituted alkylaminoalkyl groups,
99 substituted and unsubstituted dialkylaminoalkyl groups, substituted
100 and unsubstituted arylaminoalkyl groups, substituted and
101 unsubstituted diarylaminoalkyl groups, substituted and unsubstituted
102 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
103 alkylamino groups, substituted and unsubstituted arylamino groups,
104 substituted and unsubstituted dialkylamino groups, substituted and

105 unsubstituted diarylamino groups, substituted and unsubstituted
106 (alkyl)(aryl)amino groups, -C(=O)H, -C(=O)-alkyl groups,
107 -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups,
108 -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups,
109 -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups,
110 -C(=O)N(alkyl)(aryl) groups, -C(=O)-heterocyclyl groups,
111 -C(=O)-O-heterocyclyl groups, -C(=O)NH(heterocyclyl) groups,
112 -C(=O)-N(heterocyclyl)₂ groups, -C(=O)N(aryl)(heterocyclyl)
113 groups, substituted and unsubstituted heterocyclylaminoalkyl groups,
114 substituted and unsubstituted hydroxyalkyl groups, substituted and
115 unsubstituted alkoxyalkyl groups, substituted and unsubstituted
116 aryloxyalkyl groups, substituted and unsubstituted
117 heterocyclyoxyalkyl groups, and -C(=O)-N(alkyl)(heterocyclyl)
118 groups;

119 R¹⁵ and R¹⁹ may be the same or different and are independently
120 selected from the group consisting of substituted and unsubstituted
121 alkyl groups, substituted and unsubstituted aryl groups, substituted
122 and unsubstituted heterocyclyl groups, substituted and unsubstituted
123 heterocyclylalkyl groups, -C(=O)H, -C(=O)-alkyl groups,
124 -C(=O)-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups,
125 -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂
126 groups, -C(=O)N(alkyl)(aryl) groups, -NH(heterocyclyl) groups,
127 -N(heterocyclyl)₂ groups, -N(alkyl)(heterocyclyl) groups,
128 -N(aryl)(heterocyclyl) groups, substituted and unsubstituted
129 aminoalkyl groups, substituted and unsubstituted alkylaminoalkyl
130 groups, substituted and unsubstituted dialkylaminoalkyl groups,
131 substituted and unsubstituted arylaminoalkyl groups, substituted and
132 unsubstituted diarylaminoalkyl groups, substituted and unsubstituted
133 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
134 heterocyclylaminoalkyl, substituted and unsubstituted

- 135 diheterocyclylaminoalkyl, substituted and unsubstituted
136 (heterocyclyl)(alkyl)aminoalkyl, substituted and unsubstituted
137 (heterocyclyl)(aryl)aminoalkyl, substituted and unsubstituted
138 alkoxyalkyl groups, substituted and unsubstituted hydroxyalkyl
139 groups, substituted and unsubstituted aryloxyalkyl groups, and
140 substituted and unsubstituted heterocycloxyalkyl groups;
- 141 R^{16} and R^{20} may be the same or different and are independently
142 selected from the group consisting of H, substituted and unsubstituted
143 alkyl groups, substituted and unsubstituted aryl groups, and
144 substituted and unsubstituted heterocyclyl groups;
- 145 R^{17} and R^{21} may be the same or different and are independently
146 selected from the group consisting of H, substituted and unsubstituted
147 alkyl groups, substituted and unsubstituted aryl groups, substituted
148 and unsubstituted heterocyclyl groups, $-C(=O)H$, $-C(=O)$ -alkyl
149 groups, $-C(=O)$ -aryl groups, $-C(=O)NH_2$, $-C(=O)NH(alkyl)$
150 groups, $-C(=O)NH(aryl)$ groups, $-C(=O)N(alkyl)_2$ groups,
151 $-C(=O)N(aryl)_2$ groups, $-C(=O)N(alkyl)(aryl)$ groups,
152 $-C(=O)O$ -alkyl groups, $-C(=O)O$ -aryl groups, substituted and
153 unsubstituted heterocyclylalkyl groups, substituted and unsubstituted
154 aminoalkyl groups, substituted and unsubstituted alkylaminoalkyl
155 groups, substituted and unsubstituted dialkylaminoalkyl groups,
156 substituted and unsubstituted arylaminoalkyl groups, substituted and
157 unsubstituted diarylaminoalkyl groups, substituted and unsubstituted
158 (alkyl)(aryl)aminoalkyl groups, $-C(=O)$ -heterocyclyl groups,
159 $-C(=O)$ -O-heterocyclyl groups, $-C(=O)NH(heterocyclyl)$ groups,
160 $-C(=O)N(heterocyclyl)_2$ groups, $-C(=O)N(aryl)(heterocyclyl)$
161 groups, substituted and unsubstituted heterocyclylaminoalkyl groups,
162 substituted and unsubstituted diheterocyclylaminoalkyl groups,
163 substituted and unsubstituted (heterocyclyl)(alkyl)aminoalkyl groups,

-170-

164 substituted and unsubstituted (heterocyclyl)(aryl)aminoalkyl groups,
165 substituted and unsubstituted hydroxyalkyl groups, substituted and
166 unsubstituted alkoxyalkyl groups, substituted and unsubstituted
167 aryloxyalkyl groups, substituted and unsubstituted
168 heterocyclyloxyalkyl groups, and $-C(=O)-N(\text{alkyl})(\text{heterocyclyl})$
169 groups;

170 R^{18} , R^{23} , R^{24} , and R^{25} may be the same or different and are
171 independently selected from the group consisting of H, $-NH_2$,
172 $-NH(\text{alkyl})$ groups, $-NH(\text{aryl})$ groups, $-N(\text{alkyl})_2$ groups, $-N(\text{aryl})_2$
173 groups, $-N(\text{alkyl})(\text{aryl})$ groups, $-NH(\text{heterocyclyl})$ groups,
174 $-N(\text{heterocyclyl})(\text{alkyl})$ groups, $-N(\text{heterocyclyl})(\text{aryl})$ groups,
175 $-N(\text{heterocyclyl})_2$ groups, substituted and unsubstituted alkyl groups,
176 substituted and unsubstituted aryl groups, $-OH$, substituted and
177 unsubstituted alkoxy groups, substituted and unsubstituted
178 heterocyclyl groups, substituted and unsubstituted aryloxy groups,
179 heterocyclyloxy groups, $-NHOH$, $-N(\text{alkyl})OH$ groups, $-N(\text{aryl})OH$
180 groups, $-N(\text{alkyl})O\text{-alkyl}$ groups, $-N(\text{aryl})O\text{-alkyl}$ groups,
181 $-N(\text{alkyl})O\text{-aryl}$ groups, and $-N(\text{aryl})O\text{-aryl}$ groups; and

182 R^{22} is selected from the group consisting of substituted and
183 unsubstituted alkyl groups, substituted and unsubstituted aryl groups,
184 and substituted and unsubstituted heterocyclyl groups.

1 2. The compound according to claim 1, wherein Y is selected
2 from the group consisting of $-OR^{10}$ groups, $-NR^{12}R^{13}$ groups, and substituted and
3 unsubstituted alkynyl groups.

1 3. The compound according to claim 1, wherein Z is an $-NR^{14}$
2 group.

-171-


1 4. The compound according to claim 1, wherein R¹ is selected
2 from the group consisting of -H, substituted and unsubstituted alkoxy groups,
3 substituted and unsubstituted heterocyclalkoxy groups, substituted and
4 unsubstituted heterocycloxy groups, and substituted and unsubstituted heterocycl
5 groups.

1 5. The compound according to claim 1, wherein R² is selected
2 from the group consisting of H, F, Cl, -NO₂, substituted and unsubstituted
3 heterocyclalkoxy groups, and substituted and unsubstituted heterocycl groups.

1 6. The compound according to claim 1, wherein R⁶ or R⁷ is an
2 alkyl group.

1 7. The compound according to claim 1, wherein R⁶ or R⁷ is an
2 -OR¹⁹ group and R¹⁹ is an alkyl group, an aryl group, a heterocycl group, or a
3 heterocyclalkyl group.

1 8. A compound having the structure I, a tautomer of the
2 compound, a pharmaceutically acceptable salt of the compound, or a
3 pharmaceutically acceptable salt of the tautomer



I

4

5

wherein,

-172-

6 Y is selected from the group consisting of $-OR^{10}$ groups, $-C(=O)-R^{11}$
7 groups, $-NR^{12}R^{13}$ groups, substituted and unsubstituted alkynyl
8 groups, substituted and unsubstituted heterocyclalkyl groups,
9 substituted and unsubstituted alkylaminoalkyl groups, substituted and
10 unsubstituted dialkylaminoalkyl groups, substituted and unsubstituted
11 arylaminoalkyl groups, substituted and unsubstituted
12 diarylaminoalkyl groups, substituted and unsubstituted
13 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
14 heterocyclaminoalkyl groups, substituted and unsubstituted
15 saturated heterocycl groups, substituted and unsubstituted
16 heterocycloxyalkyl groups, substituted and unsubstituted
17 hydroxyalkyl groups, and substituted and unsubstituted aryloxyalkyl
18 groups;

19 Z is selected from the group consisting of O, S, and NR^{14} groups;

20 R^1 , R^2 , R^3 , and R^4 may be the same or different and are
21 independently selected from the group consisting of H, Cl, Br, F, I,
22 $-CN$, $-NO_2$, $-OH$, $-OR^{15}$ groups, $-NR^{16}R^{17}$ groups, substituted and
23 unsubstituted amidinyl groups, substituted and unsubstituted
24 guanidinyl groups, substituted and unsubstituted primary, secondary,
25 and tertiary alkyl groups, substituted and unsubstituted aryl groups,
26 substituted and unsubstituted alkenyl groups, substituted and
27 unsubstituted alkynyl groups, substituted and unsubstituted
28 heterocycl groups, substituted and unsubstituted aminoalkyl groups,
29 substituted and unsubstituted alkylaminoalkyl groups, substituted and
30 unsubstituted dialkylaminoalkyl groups, substituted and unsubstituted
31 arylaminoalkyl groups, substituted and unsubstituted
32 diarylaminoalkyl groups, substituted and unsubstituted
33 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
34 heterocyclalkyl groups, and $-C(=O)R^{18}$ groups;

35 R^5 , R^6 , R^7 , and R^8 may be the same or different and are
36 independently selected from the group consisting of H, Cl, Br, F, I,
37 $-NO_2$, $-OH$, $-OR^{19}$ groups, $-NR^{20}R^{21}$ groups, $-SH$, $-SR^{22}$ groups,
38 $-S(=O)R^{23}$ groups, $-S(=O)_2R^{24}$ groups, $-CN$, substituted and
39 unsubstituted amidinyl groups, substituted and unsubstituted
40 guanidinyl groups, substituted and unsubstituted primary, secondary,
41 and tertiary alkyl groups, substituted and unsubstituted aryl groups,
42 substituted and unsubstituted alkenyl groups, substituted and
43 unsubstituted alkynyl groups, substituted and unsubstituted
44 heterocyclyl groups, substituted and unsubstituted heterocyclylalkyl
45 groups, $-C(=O)R^{25}$ groups, substituted and unsubstituted aminoalkyl
46 groups, substituted and unsubstituted alkylaminoalkyl groups,
47 substituted and unsubstituted dialkylaminoalkyl groups, substituted
48 and unsubstituted arylaminoalkyl groups, substituted and
49 unsubstituted diarylaminoalkyl groups, substituted and unsubstituted
50 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
51 heterocyclylaminoalkyl groups, substituted and unsubstituted
52 hydroxyalkyl groups, substituted and unsubstituted alkoxyalkyl
53 groups, substituted and unsubstituted aryloxyalkyl groups, and
54 substituted and unsubstituted heterocyclyloxyalkyl groups;

55 R^9 is selected from the group consisting of $-OH$, substituted and
56 unsubstituted alkoxy groups, substituted and unsubstituted aryloxy
57 groups, $-NH_2$, substituted and unsubstituted alkylamino groups,
58 substituted and unsubstituted arylamino groups, substituted and
59 unsubstituted dialkylamino groups, substituted and unsubstituted
60 diarylamino groups, substituted and unsubstituted (alkyl)(aryl)amino
61 groups, substituted and unsubstituted alkyl groups, substituted and
62 unsubstituted aryl groups, $-C(=O)H$, $-C(=O)$ -alkyl groups, and
63 $-C(=O)$ -aryl groups;

64 R¹⁰ is selected from the group consisting of substituted and
65 unsubstituted aryl groups, substituted and unsubstituted heterocyclyl
66 groups, -C(=O)H, -C(=O)-alkyl groups, -C(=O)-aryl groups,
67 -C(=O)O-alkyl groups, -C(=O)O-aryl groups, -C(=O)NH₂,
68 -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups,
69 -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups,
70 -C(=O)N(alkyl)(aryl) groups, -NH₂, -NH(alkyl) groups, -NH(aryl)
71 groups, -N(alkyl)₂ groups, -N(alkyl)(aryl) groups, -N(aryl)₂ groups,
72 -C(=O)NH(heterocyclyl) groups, -C(=O)N(heterocyclyl)₂ groups,
73 -C(=O)N(alkyl)(heterocyclyl) groups, -C(=O)N(aryl)(heterocyclyl)
74 groups, and substituted and unsubstituted heterocyclylalkyl groups;

75 R¹¹ is selected from the group consisting of H, -NH₂, -NH(alkyl)
76 groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(aryl)₂ groups,
77 -N(alkyl)(aryl) groups, -NH(heterocyclyl) groups, -N(heterocyclyl)₂
78 groups, -N(alkyl)(heterocyclyl) groups, -O-alkyl groups, O-aryl
79 groups, substituted and unsubstituted alkyl groups, and substituted
80 and unsubstituted aryl groups;

81 R¹² is selected from the group consisting of H, substituted and
82 unsubstituted alkyl groups, substituted and unsubstituted aryl groups,
83 and substituted and unsubstituted heterocyclyl groups;

84 R¹³ is selected from the group consisting of H, substituted and
85 unsubstituted alkyl groups, substituted and unsubstituted aryl groups,
86 substituted and unsubstituted heterocyclyl groups, -OH, alkoxy
87 groups, aryloxy groups, -NH₂, substituted and unsubstituted
88 alkylamino groups, substituted and unsubstituted arylamino groups,
89 substituted and unsubstituted dialkylamino groups, substituted and
90 unsubstituted diarylamino groups, substituted and unsubstituted
91 (alkyl)(aryl)amino groups, -C(=O)H, -C(=O)-alkyl groups,

-175-

92 -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups,
93 -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups,
94 -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups,
95 -C(=O)N(alkyl)(aryl) groups, substituted and unsubstituted
96 heterocyclalkyl groups, substituted and unsubstituted aminoalkyl
97 groups, substituted and unsubstituted alkylaminoalkyl groups,
98 substituted and unsubstituted dialkylaminoalkyl groups, substituted
99 and unsubstituted arylaminoalkyl groups, substituted and
100 unsubstituted diarylaminoalkyl groups, substituted and unsubstituted
101 (alkyl)(aryl)aminoalkyl groups, -C(=O)-heterocycl groups,
102 -C(=O)-O-heterocycl groups, -C(=O)NH(heterocycl) groups,
103 -C(=O)-N(heterocycl)₂ groups, -C(=O)N(aryl)(heterocycl)
104 groups, -C(=O)-N(alkyl)(heterocycl) groups, substituted and
105 unsubstituted heterocyclaminoalkyl groups, substituted and
106 unsubstituted hydroxyalkyl groups, substituted and unsubstituted
107 alkoxyalkyl groups, substituted and unsubstituted aryloxyalkyl
108 groups, and substituted and unsubstituted heterocycloxyalkyl
109 groups;

110 R¹⁴ is selected from the group consisting of H, -OH, substituted and
111 unsubstituted alkoxy groups, substituted and unsubstituted aryloxy
112 groups, -NH₂, substituted and unsubstituted alkylamino groups,
113 substituted and unsubstituted arylamino groups, substituted and
114 unsubstituted dialkylamino groups, substituted and unsubstituted
115 diarylamino groups, substituted and unsubstituted (alkyl)(aryl)amino
116 groups, substituted and unsubstituted alkyl groups, substituted and
117 unsubstituted aryl groups, -C(=O)H, -C(=O)-alkyl groups, and
118 -C(=O)-aryl groups;

119 R¹⁵ and R¹⁹ may be the same or different and are independently
120 selected from the group consisting of substituted and unsubstituted

-176-

121 alkyl groups, substituted and unsubstituted aryl groups, substituted
122 and unsubstituted heterocyclyl groups, substituted and unsubstituted
123 heterocyclylalkyl groups, -C(=O)H, -C(=O)-alkyl groups,
124 -C(=O)-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups,
125 -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂
126 groups, -C(=O)N(alkyl)(aryl) groups, substituted and unsubstituted
127 aminoalkyl groups, substituted and unsubstituted alkylaminoalkyl
128 groups, substituted and unsubstituted dialkylaminoalkyl groups,
129 substituted and unsubstituted arylaminoalkyl groups, substituted and
130 unsubstituted diarylaminoalkyl groups, substituted and unsubstituted
131 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
132 heterocyclylaminoalkyl, substituted and unsubstituted
133 diheterocyclylaminoalkyl, substituted and unsubstituted
134 (heterocyclyl)(alkyl)aminoalkyl, substituted and unsubstituted
135 (heterocyclyl)(aryl)aminoalkyl, substituted and unsubstituted
136 alkoxyalkyl groups, substituted and unsubstituted hydroxyalkyl
137 groups, substituted and unsubstituted aryloxyalkyl groups, and
138 substituted and unsubstituted heterocyclyloxyalkyl groups;

139 R¹⁶ and R²⁰ may be the same or different and are independently
140 selected from the group consisting of H, substituted and unsubstituted
141 alkyl groups, substituted and unsubstituted aryl groups, and
142 substituted and unsubstituted heterocyclyl groups;

143 R¹⁷ and R²¹ may be the same or different and are independently
144 selected from the group consisting of H, substituted and unsubstituted
145 alkyl groups, substituted and unsubstituted aryl groups, substituted
146 and unsubstituted heterocyclyl groups, -C(=O)H, -C(=O)-alkyl
147 groups, -C(=O)-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl)
148 groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups,
149 -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups,

150 -C(=O)O-alkyl groups, -C(=O)O-aryl groups, substituted and
151 unsubstituted heterocyclalkyl groups, substituted and unsubstituted
152 aminoalkyl groups, substituted and unsubstituted alkylaminoalkyl
153 groups, substituted and unsubstituted dialkylaminoalkyl groups,
154 substituted and unsubstituted arylaminoalkyl groups, substituted and
155 unsubstituted diarylaminoalkyl groups, substituted and unsubstituted
156 (alkyl)(aryl)aminoalkyl groups, -C(=O)-heterocycl groups,
157 -C(=O)-O-heterocycl groups, -C(=O)NH(heterocycl) groups,
158 -C(=O)-N(heterocycl)₂ groups, -C(=O)N(aryl)(heterocycl)
159 groups, -C(=O)-N(alkyl)(heterocycl) groups, substituted and
160 unsubstituted heterocyclaminoalkyl groups, substituted and
161 unsubstituted hydroxyalkyl groups, substituted and unsubstituted
162 alkoxyalkyl groups, substituted and unsubstituted aryloxyalkyl
163 groups, and substituted and unsubstituted heterocycloxyalkyl
164 groups;

165 R¹⁸, R²³, R²⁴, and R²⁵ may be the same or different and are
166 independently selected from the group consisting of H, -NH₂,
167 -NH(alkyl) groups, -NH(aryl) groups, -N(alkyl)₂ groups, -N(aryl)₂
168 groups, -N(alkyl)(aryl) groups, -NH(heterocycl) groups,
169 -N(heterocycl)(alkyl) groups, -N(heterocycl)(aryl) groups,
170 -N(heterocycl)₂ groups, substituted and unsubstituted alkyl groups,
171 substituted and unsubstituted aryl groups, -OH, substituted and
172 unsubstituted alkoxy groups, substituted and unsubstituted
173 heterocycl groups, substituted and unsubstituted aryloxy groups,
174 -NHOH, -N(alkyl)OH groups, -N(aryl)OH groups, -N(alkyl)O-alkyl
175 groups, -N(aryl)O-alkyl groups, -N(alkyl)O-aryl groups, and
176 -N(aryl)O-aryl groups; and

-178-

177 R²² is selected from the group consisting of substituted and
178 unsubstituted alkyl groups, substituted and unsubstituted aryl groups,
179 and substituted and unsubstituted heterocyclyl groups.

1 9. The compound according to claim 8, wherein Y is selected
2 from the group consisting of -OR¹⁰ groups, -NR¹²R¹³ groups, and substituted and
3 unsubstituted alkynyl groups.

1 10. The compound according to claim 8, wherein Z is an -NR¹⁴
2 group.

1 11. The compound according to claim 8, wherein R¹ is selected
2 from the group consisting of -H, substituted and unsubstituted alkoxy groups,
3 substituted and unsubstituted heterocyclylalkoxy groups, substituted and
4 unsubstituted heterocycliloxy groups, and substituted and unsubstituted heterocyclyl
5 groups.

1 12. The compound according to claim 8, wherein R² is selected
2 from the group consisting of H, F, Cl, -NO₂, substituted and unsubstituted
3 heterocyclyl groups, and substituted and unsubstituted heterocyclylalkoxy groups.


1 13. The compound according to claim 8, wherein R⁶ or R⁷ is an
2 alkyl group.

1 14. The compound according to claim 8, wherein R⁶ or R⁷ is an
2 -OR¹⁹ group and R¹⁹ is an alkyl group, an aryl group, a heterocyclyl group, or a
3 heterocyclylalkyl group.

-179-

- 1 15. A compound having the structure I, a tautomer of the
 2 compound, a pharmaceutically acceptable salt of the compound, or a
 3 pharmaceutically acceptable salt of the tautomer

4



5

I

6

wherein,

- 7 Y is selected from the group consisting of -OH, SH, alkylthio
 8 groups, arylthio groups, -OR¹⁰ groups, -C(=O)-R¹¹ groups, -NR¹²R¹³
 9 groups, -CN, substituted and unsubstituted alkyl groups, substituted
 10 and unsubstituted alkenyl groups, substituted and unsubstituted
 11 alkynyl groups, substituted and unsubstituted aralkyl groups,
 12 substituted and unsubstituted heterocyclalkyl groups, substituted
 13 and unsubstituted alkylaminoalkyl groups, substituted and
 14 unsubstituted dialkylaminoalkyl groups, substituted and unsubstituted
 15 arylaminoalkyl groups, substituted and unsubstituted
 16 diarylaminoalkyl groups, substituted and unsubstituted
 17 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
 18 heterocyclaminoalkyl groups, substituted and unsubstituted
 19 heterocycl groups, substituted and unsubstituted aryl groups,
 20 substituted and unsubstituted heterocycloxyalkyl groups, substituted

-180-

21 and unsubstituted hydroxyalkyl groups, substituted and unsubstituted
22 alkoxyalkyl groups, and substituted and unsubstituted aryloxyalkyl
23 groups;

24 Z is selected from the group consisting of O, S, and NR^{14} groups;

25 R^1 , R^2 , R^3 , and R^4 may be the same or different and are
26 independently selected from the group consisting of H, Cl, Br, F, I,
27 $-\text{CN}$, $-\text{NO}_2$, $-\text{OH}$, $-\text{OR}^{15}$ groups, $-\text{NR}^{16}\text{R}^{17}$ groups, substituted and
28 unsubstituted amidinyl groups, substituted and unsubstituted
29 guanidinyl groups, substituted and unsubstituted primary, secondary,
30 and tertiary alkyl groups, substituted and unsubstituted aryl groups,
31 substituted and unsubstituted alkenyl groups, substituted and
32 unsubstituted alkynyl groups, substituted and unsubstituted
33 heterocyclyl groups, substituted and unsubstituted aminoalkyl groups,
34 substituted and unsubstituted alkylaminoalkyl groups, substituted and
35 unsubstituted dialkylaminoalkyl groups, substituted and unsubstituted
36 arylaminoalkyl groups, substituted and unsubstituted
37 diarylaminoalkyl groups, substituted and unsubstituted
38 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
39 heterocyclylalkyl groups, and $-\text{C}(=\text{O})\text{R}^{18}$ groups;

40 R^5 , R^6 , R^7 , and R^8 may be the same or different and are
41 independently selected from the group consisting of H, Cl, Br, F, I,
42 $-\text{NO}_2$, $-\text{OH}$, $-\text{OR}^{19}$ groups, $-\text{NR}^{20}\text{R}^{21}$ groups, $-\text{SH}$, $-\text{SR}^{22}$ groups,
43 $-\text{S}(=\text{O})\text{R}^{23}$ groups, $-\text{S}(=\text{O})_2\text{R}^{24}$ groups, $-\text{CN}$, substituted and
44 unsubstituted amidinyl groups, substituted and unsubstituted
45 guanidinyl groups, substituted and unsubstituted primary, secondary,
46 and tertiary alkyl groups, substituted and unsubstituted aryl groups,
47 substituted and unsubstituted alkenyl groups, substituted and
48 unsubstituted alkynyl groups, substituted and unsubstituted

-181-

49 heterocyclyl groups, substituted and unsubstituted alkylaminoalkyl
50 groups, substituted and unsubstituted dialkylaminoalkyl groups,
51 substituted and unsubstituted arylaminoalkyl groups, substituted and
52 unsubstituted diarylaminoalkyl groups, substituted and unsubstituted
53 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
54 heterocyclylalkyl groups, $-C(=O)R^{25}$ groups, substituted and
55 unsubstituted aminoalkyl groups, substituted and unsubstituted
56 heterocyclylaminoalkyl groups, substituted and unsubstituted
57 hydroxyalkyl groups, substituted and unsubstituted alkoxyalkyl
58 groups, substituted and unsubstituted aryloxyalkyl groups, and
59 substituted and unsubstituted heterocycliloxyalkyl groups;

60 R^9 and R^{14} may be the same or different and are independently
61 selected from the group consisting of H, -OH, substituted and
62 unsubstituted alkoxy groups, substituted and unsubstituted aryloxy
63 groups, $-NH_2$, substituted and unsubstituted alkylamino groups,
64 substituted and unsubstituted arylamino groups, substituted and
65 unsubstituted dialkylamino groups, substituted and unsubstituted
66 diarylamino groups, substituted and unsubstituted (alkyl)(aryl)amino
67 groups, substituted and unsubstituted alkyl groups, substituted and
68 unsubstituted aryl groups, $-C(=O)H$, $-C(=O)$ -alkyl groups, and
69 $-C(=O)$ -aryl groups;

70 R^{10} is selected from the group consisting of substituted and
71 unsubstituted alkyl groups, substituted and unsubstituted aryl groups,
72 substituted and unsubstituted heterocyclyl groups, substituted and
73 unsubstituted heterocyclylalkyl groups, $-C(=O)H$, $-C(=O)$ -alkyl
74 groups, $-C(=O)$ -aryl groups, $-C(=O)O$ -alkyl groups, $-C(=O)O$ -aryl
75 groups, $-C(=O)NH_2$, $-C(=O)NH$ (alkyl) groups, $-C(=O)NH$ (aryl)
76 groups, $-C(=O)N$ (alkyl)₂ groups, $-C(=O)N$ (aryl)₂ groups,
77 $-C(=O)N$ (alkyl)(aryl) groups, $-NH_2$, $-NH$ (alkyl) groups, $-NH$ (aryl)

-182-

78 groups, -N(alkyl)₂ groups, -N(alkyl)(aryl) groups, -N(aryl)₂ groups,
79 -C(=O)NH(heterocyclyl) groups, -C(=O)N(heterocyclyl)₂ groups,
80 -C(=O)N(alkyl)(heterocyclyl) groups, and
81 -C(=O)N(aryl)(heterocyclyl) groups;

82 R¹¹ is selected from the group consisting of H, -OH, alkoxy groups,
83 aryloxy groups, -NH₂, -NH(alkyl) groups, -NH(aryl) groups,
84 -N(alkyl)₂ groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups,
85 substituted and unsubstituted alkyl groups, -NH(heterocyclyl) groups,
86 -N(heterocyclyl)₂ groups, -N(alkyl)(heterocyclyl) groups, and
87 substituted and unsubstituted aryl groups;

88 R¹² is selected from the group consisting of H, substituted and
89 unsubstituted alkyl groups, substituted and unsubstituted aryl groups,
90 and substituted and unsubstituted heterocyclyl groups;

91 R¹³ is selected from the group consisting of H, substituted and
92 unsubstituted alkyl groups, substituted and unsubstituted aryl groups,
93 substituted and unsubstituted heterocyclyl groups, -OH, alkoxy
94 groups, aryloxy groups, -NH₂, substituted and unsubstituted
95 heterocyclylalkyl groups, substituted and unsubstituted aminoalkyl
96 groups, substituted and unsubstituted alkylaminoalkyl groups,
97 substituted and unsubstituted dialkylaminoalkyl groups, substituted
98 and unsubstituted arylaminoalkyl groups, substituted and
99 unsubstituted diarylaminoalkyl groups, substituted and unsubstituted
100 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
101 alkylamino groups, substituted and unsubstituted arylamino groups,
102 substituted and unsubstituted dialkylamino groups, substituted and
103 unsubstituted diarylamino groups, substituted and unsubstituted
104 (alkyl)(aryl)amino groups, -C(=O)H, -C(=O)-alkyl groups,
105 -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups,

106 -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups,
107 -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups,
108 -C(=O)N(alkyl)(aryl) groups, -C(=O)-heterocyclyl groups,
109 -C(=O)-O-heterocyclyl groups, -C(=O)NH(heterocyclyl) groups,
110 -C(=O)-N(heterocyclyl)₂ groups, -C(=O)-N(alkyl)(heterocyclyl)
111 groups, -C(=O)-N(aryl)(heterocyclyl) groups, substituted and
112 unsubstituted heterocyclylaminoalkyl groups, substituted and
113 unsubstituted hydroxyalkyl groups, substituted and unsubstituted
114 alkoxyalkyl groups, substituted and unsubstituted aryloxyalkyl
115 groups, and substituted and unsubstituted heterocyclioxyalkyl
116 groups;

117 R¹⁵ and R¹⁹ may be the same or different and are independently
118 selected from the group consisting of substituted and unsubstituted
119 alkyl groups, substituted and unsubstituted aryl groups, substituted
120 and unsubstituted heterocyclyl groups, substituted and unsubstituted
121 heterocyclylalkyl groups, -C(=O)H, -C(=O)-alkyl groups,
122 -C(=O)-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups,
123 -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂
124 groups, -C(=O)N(alkyl)(aryl) groups, substituted and unsubstituted
125 aminoalkyl groups, substituted and unsubstituted alkylaminoalkyl
126 groups, substituted and unsubstituted dialkylaminoalkyl groups,
127 substituted and unsubstituted arylaminoalkyl groups, substituted and
128 unsubstituted diarylaminoalkyl groups, substituted and unsubstituted
129 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
130 heterocyclylaminoalkyl, substituted and unsubstituted
131 diheterocyclylaminoalkyl, substituted and unsubstituted
132 (heterocyclyl)(alkyl)aminoalkyl, substituted and unsubstituted
133 (heterocyclyl)(aryl)aminoalkyl, substituted and unsubstituted
134 alkoxyalkyl groups, substituted and unsubstituted hydroxyalkyl

135 groups, substituted and unsubstituted aryloxyalkyl groups, and
136 substituted and unsubstituted heterocycloxyalkyl groups;

137 R^{16} and R^{20} may be the same or different and are independently
138 selected from the group consisting of H, substituted and unsubstituted
139 alkyl groups, substituted and unsubstituted aryl groups, and
140 substituted and unsubstituted heterocyclyl groups;

141 R^{17} and R^{21} may be the same or different and are independently
142 selected from the group consisting of H, substituted and unsubstituted
143 alkyl groups, substituted and unsubstituted aryl groups, substituted
144 and unsubstituted heterocyclyl groups, $-C(=O)H$, $-C(=O)$ -alkyl
145 groups, $-C(=O)$ -aryl groups, $-C(=O)NH_2$, $-C(=O)NH$ (alkyl)
146 groups, $-C(=O)NH$ (aryl) groups, $-C(=O)N$ (alkyl) $_2$ groups,
147 $-C(=O)N$ (aryl) $_2$ groups, $-C(=O)N$ (alkyl)(aryl) groups,
148 $-C(=O)O$ -alkyl groups, $-C(=O)O$ -aryl groups, substituted and
149 unsubstituted heterocyclylalkyl groups, substituted and unsubstituted
150 aminoalkyl groups, substituted and unsubstituted alkylaminoalkyl
151 groups, substituted and unsubstituted dialkylaminoalkyl groups,
152 substituted and unsubstituted arylaminoalkyl groups, substituted and
153 unsubstituted diarylaminoalkyl groups, substituted and unsubstituted
154 (alkyl)(aryl)aminoalkyl groups, $-C(=O)$ -heterocyclyl groups,
155 $-C(=O)-O$ -heterocyclyl groups, $-C(=O)NH$ (heterocyclyl) groups,
156 $-C(=O)-N$ (heterocyclyl) $_2$ groups, $-C(=O)-N$ (alkyl)(heterocyclyl)
157 groups, $-C(=O)-N$ (aryl)(heterocyclyl) groups, substituted and
158 unsubstituted heterocyclylaminoalkyl groups, substituted and
159 unsubstituted hydroxyalkyl groups, substituted and unsubstituted
160 alkoxyalkyl groups, substituted and unsubstituted aryloxyalkyl
161 groups, and substituted and unsubstituted heterocycloxyalkyl
162 groups;

-185-

163 R^{18} , R^{23} , R^{24} , and R^{25} may be the same or different and are
164 independently selected from the group consisting of H, $-NH_2$,
165 $-NH(alkyl)$ groups, $-NH(aryl)$ groups, $-N(alkyl)_2$ groups, $-N(aryl)_2$
166 groups, $-N(alkyl)(aryl)$ groups, $-NH(heterocyclyl)$ groups,
167 $-N(heterocyclyl)(alkyl)$ groups, $-N(heterocyclyl)(aryl)$ groups,
168 $-N(heterocyclyl)_2$ groups, substituted and unsubstituted alkyl groups,
169 substituted and unsubstituted aryl groups, $-OH$, substituted and
170 unsubstituted alkoxy groups, substituted and unsubstituted aryloxy
171 groups, substituted and unsubstituted heterocyclyl groups, $-NHOH$,
172 $-N(alkyl)OH$ groups, $-N(aryl)OH$ groups, $-N(alkyl)O-alkyl$ groups,
173 $-N(aryl)O-alkyl$ groups, $-N(alkyl)O-aryl$ groups, and $-N(aryl)O-aryl$
174 groups; and

175 R^{22} is selected from the group consisting of substituted and
176 unsubstituted alkyl groups, substituted and unsubstituted aryl groups,
177 and substituted and unsubstituted heterocyclyl groups;

178 and further wherein at least one of R^5 , R^6 , R^7 , or R^8 is selected from
179 the group consisting of substituted and unsubstituted amidinyl groups,
180 substituted and unsubstituted guanidinyl groups, substituted and
181 unsubstituted saturated heterocyclyl groups, substituted and
182 unsubstituted alkylaminoalkyl groups, substituted and unsubstituted
183 dialkylaminoalkyl groups, substituted and unsubstituted
184 arylaminoalkyl groups, substituted and unsubstituted
185 diarylaminoalkyl groups, substituted and unsubstituted
186 $(alkyl)(aryl)aminoalkyl$ groups, substituted and unsubstituted
187 heterocyclylalkyl groups, substituted and unsubstituted
188 heterocyclylaminoalkyl groups, substituted and unsubstituted
189 hydroxyalkyl groups, substituted and unsubstituted alkoxyalkyl
190 groups, substituted and unsubstituted aryloxyalkyl groups, and
191 substituted and unsubstituted heterocycloxyalkyl groups; $-OR^{19}$

192 groups wherein R^{19} is selected from the group consisting of
193 substituted and unsubstituted aryl groups, substituted and
194 unsubstituted heterocyclyl groups, substituted and unsubstituted
195 heterocyclylalkyl groups, $-C(=O)H$, $-C(=O)$ -aryl groups,
196 $-C(=O)NH_2$, $-C(=O)NH(alkyl)$ groups, $-C(=O)NH(aryl)$ groups,
197 $-C(=O)N(alkyl)_2$ groups, $-C(=O)N(aryl)_2$ groups,
198 $-C(=O)N(alkyl)(aryl)$ groups, substituted and unsubstituted
199 aminoalkyl groups, substituted and unsubstituted alkylaminoalkyl
200 groups, substituted and unsubstituted dialkylaminoalkyl groups,
201 substituted and unsubstituted arylaminoalkyl groups, substituted and
202 unsubstituted diarylaminoalkyl groups, substituted and unsubstituted
203 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
204 heterocyclylaminoalkyl groups, substituted and unsubstituted
205 diheterocyclylaminoalkyl groups, substituted and unsubstituted
206 (heterocyclyl)(alkyl)aminoalkyl groups, substituted and unsubstituted
207 (heterocyclyl)(aryl)aminoalkyl groups, substituted and unsubstituted
208 hydroxyalkyl groups, substituted and unsubstituted alkoxyalkyl
209 groups, substituted and unsubstituted aryloxyalkyl groups, and
210 substituted and unsubstituted heterocyclyloxyalkyl groups; $-NR^{20}R^{21}$
211 groups wherein R^{20} is selected from the group consisting of
212 substituted and unsubstituted heterocyclyl groups; $-NR^{20}R^{21}$ groups
213 wherein R^{21} is selected from the group consisting of substituted and
214 unsubstituted heterocyclyl groups, $-C(=O)H$, $-C(=O)$ -aryl groups,
215 $-C(=O)NH_2$, $-C(=O)NH(alkyl)$ groups, $-C(=O)NH(aryl)$ groups,
216 $-C(=O)N(alkyl)_2$ groups, $-C(=O)N(aryl)_2$ groups,
217 $-C(=O)N(alkyl)(aryl)$ groups, $-C(=O)O$ -alkyl groups,
218 $-C(=O)O$ -aryl groups, substituted and unsubstituted aminoalkyl
219 groups, substituted and unsubstituted alkylaminoalkyl groups,
220 substituted and unsubstituted dialkylaminoalkyl groups, substituted
221 and unsubstituted arylaminoalkyl groups, substituted and
222 unsubstituted diarylaminoalkyl groups, substituted and unsubstituted

-187-

223 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
224 heterocyclylaminoalkyl groups, substituted and unsubstituted
225 hydroxyalkyl groups, substituted and unsubstituted alkoxyalkyl
226 groups, substituted and unsubstituted aryloxyalkyl groups, substituted
227 and unsubstituted heterocyclylalkyl groups, and substituted and
228 unsubstituted heterocyclcyloxyalkyl groups; and $-C(=O)R^{25}$ groups
229 wherein R^{25} is selected from the group consisting of H, $-NH_2$,
230 $-NH(alkyl)$ groups, $-NH(aryl)$ groups, $-N(alkyl)_2$ groups, $-N(aryl)_2$
231 groups, $-N(alkyl)(aryl)$ groups, $-NH(heterocyclyl)$ groups,
232 $-N(heterocyclyl)(alkyl)$ groups, $-N(heterocyclyl)(aryl)$ groups,
233 $-N(heterocyclyl)_2$ groups, substituted and unsubstituted aryl groups,
234 substituted and unsubstituted aryloxy groups, and substituted and
235 unsubstituted heterocyclyl groups.

1 16. The compound according to claim 15, wherein Y is selected
2 from the group consisting of $-OR^{10}$ groups, $-NR^{12}R^{13}$ groups, and substituted and
3 unsubstituted alkynyl groups.

1 17. The compound according to claim 15, wherein Z is an $-NR^{14}$
2 group.

1 18. The compound according to any of claims 40-43, wherein R^1
2 is selected from the group consisting of $-H$, substituted and unsubstituted alkoxy
3 groups, substituted and unsubstituted heterocyclylalkoxy groups, substituted and
4 unsubstituted heterocyclcyloxy groups, and substituted and unsubstituted heterocyclyl
5 groups.


1 19. The compound according to claim 15, wherein R^2 is selected
2 from the group consisting of H, F, Cl, $-NO_2$, substituted and unsubstituted
3 heterocyclyl groups, and substituted and unsubstituted heterocyclylalkoxy groups.

-188-

1 20. The compound according to claim 15, wherein R⁶ or R⁷ is an
2 alkyl group.

1 21. The compound according to claim 15, wherein R⁶ or R⁷ is an
2 -OR¹⁹ group and R¹⁹ is an alkyl group, an aryl group, a heterocyclyl group, or a
3 heterocyclalkyl group.

1 22. A compound having the structure I, a tautomer of the
2 compound, a pharmaceutically acceptable salt of the compound, or a
3 pharmaceutically acceptable salt of the tautomer



4
5 wherein,

6 Y is selected from the group consisting of -OH, SH, alkylthio
7 groups, arylthio groups, -OR¹⁰ groups, -C(=O)-R¹¹ groups, -NR¹²R¹³
8 groups, -CN, substituted and unsubstituted alkyl groups, substituted
9 and unsubstituted alkenyl groups, substituted and unsubstituted
10 alkynyl groups, substituted and unsubstituted aralkyl groups,
11 substituted and unsubstituted heterocyclalkyl groups, substituted
12 and unsubstituted alkylaminoalkyl groups, substituted and
13 unsubstituted dialkylaminoalkyl groups, substituted and unsubstituted
14 arylaminoalkyl groups, substituted and unsubstituted

15 diarylaminoalkyl groups, substituted and unsubstituted
16 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
17 heterocyclylaminoalkyl groups, substituted and unsubstituted
18 heterocyclyl groups, substituted and unsubstituted aryl groups,
19 substituted and unsubstituted heterocycloxyalkyl groups, substituted
20 and unsubstituted hydroxyalkyl groups, substituted and unsubstituted
21 alkoxyalkyl groups, and substituted and unsubstituted aryloxyalkyl
22 groups;

23 Z is selected from the group consisting of O, S, and NR^{14} groups;

24 R^1 , R^2 , R^3 , and R^4 may be the same or different and are
25 independently selected from the group consisting of H, Cl, Br, F, I,
26 -CN, -NO₂, -OH, -OR¹⁵ groups, -NR¹⁶R¹⁷ groups, substituted and
27 unsubstituted amidinyl groups, substituted and unsubstituted
28 guanidinyl groups, substituted and unsubstituted primary, secondary,
29 and tertiary alkyl groups, substituted and unsubstituted aryl groups,
30 substituted and unsubstituted alkenyl groups, substituted and
31 unsubstituted alkynyl groups, substituted and unsubstituted
32 heterocyclyl groups, substituted and unsubstituted aminoalkyl groups,
33 substituted and unsubstituted alkylaminoalkyl groups, substituted and
34 unsubstituted dialkylaminoalkyl groups, substituted and unsubstituted
35 arylaminoalkyl groups, substituted and unsubstituted
36 diarylaminoalkyl groups, substituted and unsubstituted
37 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
38 heterocyclylalkyl groups, and -C(=O)R¹⁸ groups;

39 R^5 , R^6 , R^7 , and R^8 may be the same or different and are
40 independently selected from the group consisting of H, Cl, Br, F, I,
41 -NO₂, -OH, -OR¹⁹ groups, -NR²⁰R²¹ groups, -SH, -SR²² groups,
42 -S(=O)R²³ groups, -S(=O)₂R²⁴ groups, -CN, substituted and

43 unsubstituted amidinyl groups, substituted and unsubstituted
44 guanidinyl groups, substituted and unsubstituted primary, secondary,
45 and tertiary alkyl groups, substituted and unsubstituted aryl groups,
46 substituted and unsubstituted alkenyl groups, substituted and
47 unsubstituted alkynyl groups, substituted and unsubstituted
48 heterocyclyl groups, substituted and unsubstituted alkylaminoalkyl
49 groups, substituted and unsubstituted dialkylaminoalkyl groups,
50 substituted and unsubstituted arylaminoalkyl groups, substituted and
51 unsubstituted diarylaminoalkyl groups, substituted and unsubstituted
52 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
53 heterocyclylalkyl groups, $-C(=O)R^{25}$ groups, substituted and
54 unsubstituted aminoalkyl groups, substituted and unsubstituted
55 heterocyclylaminoalkyl groups, substituted and unsubstituted
56 hydroxyalkyl groups, substituted and unsubstituted alkoxyalkyl
57 groups, substituted and unsubstituted aryloxyalkyl groups, and
58 substituted and unsubstituted heterocycloxyalkyl groups;

59 R^9 and R^{14} may be the same or different and are independently
60 selected from the group consisting of H, -OH, substituted and
61 unsubstituted alkoxy groups, substituted and unsubstituted aryloxy
62 groups, $-NH_2$, substituted and unsubstituted alkylamino groups,
63 substituted and unsubstituted arylamino groups, substituted and
64 unsubstituted dialkylamino groups, substituted and unsubstituted
65 diarylamino groups, substituted and unsubstituted (alkyl)(aryl)amino
66 groups, substituted and unsubstituted alkyl groups, substituted and
67 unsubstituted aryl groups, $-C(=O)H$, $-C(=O)$ -alkyl groups, and
68 $-C(=O)$ -aryl groups;

69 R^{10} is selected from the group consisting of substituted and
70 unsubstituted alkyl groups, substituted and unsubstituted aryl groups,
71 substituted and unsubstituted heterocyclyl groups, substituted and

-191-

72 unsubstituted heterocyclylalkyl groups, -C(=O)H, -C(=O)-alkyl
73 groups, -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl
74 groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl)
75 groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups,
76 -C(=O)N(alkyl)(aryl) groups, -NH₂, -NH(alkyl) groups, -NH(aryl)
77 groups, -N(alkyl)₂ groups, -N(alkyl)(aryl) groups, -N(aryl)₂ groups,
78 -C(=O)NH(heterocyclyl) groups, -C(=O)N(heterocyclyl)₂ groups,
79 -C(=O)N(alkyl)(heterocyclyl) groups, and
80 -C(=O)N(aryl)(heterocyclyl) groups;

81 R¹¹ is selected from the group consisting of H, -OH, alkoxy groups,
82 aryloxy groups, -NH₂, -NH(alkyl) groups, -NH(aryl) groups,
83 -N(alkyl)₂ groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups,
84 substituted and unsubstituted alkyl groups, -NH(heterocyclyl) groups,
85 -N(heterocyclyl)₂ groups, -N(alkyl)(heterocyclyl) groups, and
86 substituted and unsubstituted aryl groups;

87 R¹² is selected from the group consisting of H, substituted and
88 unsubstituted alkyl groups, substituted and unsubstituted aryl groups,
89 and substituted and unsubstituted heterocyclyl groups;

90 R¹³ is selected from the group consisting of H, substituted and
91 unsubstituted alkyl groups, substituted and unsubstituted aryl groups,
92 substituted and unsubstituted heterocyclyl groups, -OH, alkoxy
93 groups, aryloxy groups, -NH₂, substituted and unsubstituted
94 heterocyclylalkyl groups, substituted and unsubstituted aminoalkyl
95 groups, substituted and unsubstituted alkylaminoalkyl groups,
96 substituted and unsubstituted dialkylaminoalkyl groups, substituted
97 and unsubstituted arylaminoalkyl groups, substituted and
98 unsubstituted diarylaminoalkyl groups, substituted and unsubstituted
99 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted

100 alkylamino groups, substituted and unsubstituted arylamino groups,
101 substituted and unsubstituted dialkylamino groups, substituted and
102 unsubstituted diarylamino groups, substituted and unsubstituted
103 (alkyl)(aryl)amino groups, -C(=O)H, -C(=O)-alkyl groups,
104 -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups,
105 -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups,
106 -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups,
107 -C(=O)N(alkyl)(aryl) groups, -C(=O)-heterocyclyl groups,
108 -C(=O)-O-heterocyclyl groups, -C(=O)NH(heterocyclyl) groups,
109 -C(=O)-N(heterocyclyl)₂ groups, -C(=O)-N(alkyl)(heterocyclyl)
110 groups, -C(=O)-N(aryl)(heterocyclyl) groups, substituted and
111 unsubstituted heterocyclylaminoalkyl groups, substituted and
112 unsubstituted hydroxyalkyl groups, substituted and unsubstituted
113 alkoxyalkyl groups, substituted and unsubstituted aryloxyalkyl
114 groups, and substituted and unsubstituted heterocycliloxyalkyl
115 groups;

116 R¹⁵ and R¹⁹ may be the same or different and are independently
117 selected from the group consisting of substituted and unsubstituted
118 alkyl groups, substituted and unsubstituted aryl groups, substituted
119 and unsubstituted heterocyclyl groups, substituted and unsubstituted
120 heterocyclylalkyl groups, -C(=O)H, -C(=O)-alkyl groups,
121 -C(=O)-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups,
122 -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂
123 groups, -C(=O)N(alkyl)(aryl) groups, substituted and unsubstituted
124 aminoalkyl groups, substituted and unsubstituted alkylaminoalkyl
125 groups, substituted and unsubstituted dialkylaminoalkyl groups,
126 substituted and unsubstituted arylaminoalkyl groups, substituted and
127 unsubstituted diarylaminoalkyl groups, substituted and unsubstituted
128 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
129 heterocyclylaminoalkyl, substituted and unsubstituted

- 130 diheterocyclylaminoalkyl, substituted and unsubstituted
131 (heterocyclyl)(alkyl)aminoalkyl, substituted and unsubstituted
132 (heterocyclyl)(aryl)aminoalkyl, substituted and unsubstituted
133 alkoxyalkyl groups, substituted and unsubstituted hydroxyalkyl
134 groups, substituted and unsubstituted aryloxyalkyl groups, and
135 substituted and unsubstituted heterocycloxyalkyl groups;
- 136 R^{16} and R^{20} may be the same or different and are independently
137 selected from the group consisting of H, substituted and unsubstituted
138 alkyl groups, substituted and unsubstituted aryl groups, and
139 substituted and unsubstituted heterocyclyl groups;
- 140 R^{17} and R^{21} may be the same or different and are independently
141 selected from the group consisting of H, substituted and unsubstituted
142 alkyl groups, substituted and unsubstituted aryl groups, substituted
143 and unsubstituted heterocyclyl groups, $-C(=O)H$, $-C(=O)$ -alkyl
144 groups, $-C(=O)$ -aryl groups, $-C(=O)NH_2$, $-C(=O)NH(alkyl)$
145 groups, $-C(=O)NH(aryl)$ groups, $-C(=O)N(alkyl)_2$ groups,
146 $-C(=O)N(aryl)_2$ groups, $-C(=O)N(alkyl)(aryl)$ groups,
147 $-C(=O)O$ -alkyl groups, $-C(=O)O$ -aryl groups, substituted and
148 unsubstituted heterocyclylalkyl groups, substituted and unsubstituted
149 aminoalkyl groups, substituted and unsubstituted alkylaminoalkyl
150 groups, substituted and unsubstituted dialkylaminoalkyl groups,
151 substituted and unsubstituted arylaminoalkyl groups, substituted and
152 unsubstituted diarylaminoalkyl groups, substituted and unsubstituted
153 (alkyl)(aryl)aminoalkyl groups, $-C(=O)$ -heterocyclyl groups,
154 $-C(=O)$ -O-heterocyclyl groups, $-C(=O)NH(heterocyclyl)$ groups,
155 $-C(=O)$ -N(heterocyclyl) $_2$ groups, $-C(=O)$ -N(alkyl)(heterocyclyl)
156 groups, $-C(=O)$ -N(aryl)(heterocyclyl) groups, substituted and
157 unsubstituted heterocyclylaminoalkyl groups, substituted and
158 unsubstituted hydroxyalkyl groups, substituted and unsubstituted

159 alkoxyalkyl groups, substituted and unsubstituted aryloxyalkyl
160 groups, and substituted and unsubstituted heterocycloxyalkyl
161 groups;

162 R^{18} , R^{23} , R^{24} , and R^{25} may be the same or different and are
163 independently selected from the group consisting of H, $-NH_2$,
164 $-NH(alkyl)$ groups, $-NH(aryl)$ groups, $-N(alkyl)_2$ groups, $-N(aryl)_2$
165 groups, $-N(alkyl)(aryl)$ groups, $-NH(heterocyclyl)$ groups,
166 $-N(heterocyclyl)(alkyl)$ groups, $-N(heterocyclyl)(aryl)$ groups,
167 $-N(heterocyclyl)_2$ groups, substituted and unsubstituted alkyl groups,
168 substituted and unsubstituted aryl groups, $-OH$, substituted and
169 unsubstituted alkoxy groups, substituted and unsubstituted aryloxy
170 groups, substituted and unsubstituted heterocyclyl groups, $-NHOH$,
171 $-N(alkyl)OH$ groups, $-N(aryl)OH$ groups, $-N(alkyl)O-alkyl$ groups,
172 $-N(aryl)O-alkyl$ groups, $-N(alkyl)O-aryl$ groups, and $-N(aryl)O-aryl$
173 groups; and

174 R^{22} is selected from the group consisting of substituted and
175 unsubstituted alkyl groups, substituted and unsubstituted aryl groups,
176 and substituted and unsubstituted heterocyclyl groups,

177 and further wherein, at least one of R^1 , R^2 , R^3 , or R^4 is an $-OR^{15}$
178 group and R^{15} is selected from the group consisting of substituted and
179 unsubstituted heterocyclylalkyl groups, substituted and unsubstituted
180 dialkylaminoalkyl groups, substituted and unsubstituted
181 alkylaminoalkyl groups, substituted and unsubstituted aminoalkyl
182 groups, substituted and unsubstituted diarylaminoalkyl groups,
183 substituted and unsubstituted arylaminoalkyl groups, substituted and
184 unsubstituted $(alkyl)(aryl)aminoalkyl$ groups, substituted and
185 unsubstituted heterocyclyl groups, substituted and unsubstituted
186 heterocyclylaminoalkyl groups, substituted and unsubstituted

187 diheterocyclylaminoalkyl groups, substituted and unsubstituted
188 (heterocyclyl)(alkyl)aminoalkyl groups, and substituted and
189 unsubstituted (heterocyclyl)(aryl)aminoalkyl groups.

1 23. The compound according to claim 22, wherein R¹ is an -OR¹⁵
2 group and R¹⁵ is selected from the group consisting of substituted and unsubstituted
3 heterocyclylalkyl groups, substituted and unsubstituted dialkylaminoalkyl groups,
4 substituted and unsubstituted alkylaminoalkyl groups, substituted and unsubstituted
5 aminoalkyl groups, substituted and unsubstituted diarylaminoalkyl groups,
6 substituted and unsubstituted arylaminoalkyl groups, substituted and unsubstituted
7 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted heterocyclyl groups,
8 substituted and unsubstituted heterocyclylaminoalkyl groups, substituted and
9 unsubstituted diheterocyclylaminoalkyl groups, substituted and unsubstituted
10 (heterocyclyl)(alkyl)aminoalkyl groups, and substituted and unsubstituted
11 (heterocyclyl)(aryl)aminoalkyl groups.

1 24. The compound according to claim 22, wherein Z is an -NR¹⁰
2 group.

1 25. The compound according to claim 22, wherein R¹ is selected
2 from the group consisting of -H, substituted and unsubstituted alkoxy groups,
3 substituted and unsubstituted heterocyclylalkoxy groups, substituted and
4 unsubstituted heterocycliloxy groups, and substituted and unsubstituted heterocyclyl
5 groups.


1 26. The compound according to claim 22, wherein R² is selected
2 from the group consisting of H, F, Cl, -NO₂, substituted and unsubstituted
3 heterocyclyl groups, and substituted and unsubstituted heterocyclylalkoxy groups.

-196-

1 27. The compound according to claim 22, wherein R⁶ or R⁷ is an
2 alkyl group.

1 28. The compound according to claim 22, wherein R⁶ or R⁷ is an
2 -OR¹⁹ group and R¹⁹ is an alkyl group, an aryl group, a heterocyclyl group, or a
3 heterocyclylalkyl group.

1 29. A compound having the structure II, a tautomer of the
2 compound, a pharmaceutically acceptable salt of the compound, or a
3 pharmaceutically acceptable salt of the tautomer



II

4

5 wherein,

6 Y is selected from the group consisting of H, -OH, -OR¹⁰ groups,
7 -SH, -SR¹¹ groups, -NR¹²R¹³ groups, -CN, -C(=O)-R¹⁴ groups,
8 substituted and unsubstituted alkyl groups, substituted and
9 unsubstituted alkenyl groups, substituted and unsubstituted alkynyl
10 groups, substituted and unsubstituted aralkyl groups, substituted and
11 unsubstituted heterocyclylalkyl groups, substituted and unsubstituted
12 alkylaminoalkyl groups, substituted and unsubstituted
13 dialkylaminoalkyl groups, substituted and unsubstituted
14 arylaminoalkyl groups, substituted and unsubstituted

15 diarylaminoalkyl groups, substituted and unsubstituted
16 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
17 heterocyclylaminoalkyl groups, substituted and unsubstituted
18 heterocyclyl groups, substituted and unsubstituted aryl groups,
19 substituted and unsubstituted hydroxyalkyl groups, substituted and
20 unsubstituted alkoxyalkyl groups, substituted and unsubstituted
21 aryloxyalkyl groups, and substituted and unsubstituted
22 heterocycloxyalkyl groups;

23 X¹, X², X³, and X⁴ are selected from the group consisting of C and
24 N, wherein at least one of X¹, X², X³, or X⁴ is N;

25 R¹, R², R³, R⁴, R⁵, R⁶, R⁷, and R⁸ may be the same or different and
26 are independently selected from the group consisting of H, Cl, Br, F,
27 I, -NO₂, -CN, -OH, -OR¹⁵ groups, -NR¹⁶R¹⁷ groups, -C(=O)R¹⁸
28 groups, -SH, -SR¹⁹ groups, -S(=O)R²⁰ groups, S(=O)₂R²¹ groups,
29 substituted and unsubstituted amidinyl groups, substituted and
30 unsubstituted guanidinyl groups, substituted and unsubstituted
31 primary, secondary, and tertiary alkyl groups, substituted and
32 unsubstituted aryl groups, substituted and unsubstituted alkenyl
33 groups, substituted and unsubstituted alkynyl groups, substituted and
34 unsubstituted heterocyclyl groups, substituted and unsubstituted
35 alkylaminoalkyl groups, substituted and unsubstituted
36 dialkylaminoalkyl groups, substituted and unsubstituted
37 arylaminoalkyl groups, substituted and unsubstituted
38 diarylaminoalkyl groups, substituted and unsubstituted
39 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
40 heterocyclylalkyl groups, substituted and unsubstituted aminoalkyl
41 groups, substituted and unsubstituted heterocyclylaminoalkyl groups,
42 substituted and unsubstituted hydroxyalkyl groups, substituted and
43 unsubstituted alkoxyalkyl groups, substituted and unsubstituted

- 44 aryloxyalkyl groups, and substituted and unsubstituted
45 heterocycloxyalkyl groups; R⁵ is absent or is H if X¹ is N; R⁶ is
46 absent or is H if X² is N; R⁷ is absent or is H if X³ is N; and R⁸ is
47 absent or is H if X⁴ is N;
- 48 R⁹ is selected from the group consisting of H, -OH, substituted and
49 unsubstituted alkoxy groups, substituted and unsubstituted aryloxy
50 groups, -NH₂, substituted and unsubstituted alkylamino groups,
51 substituted and unsubstituted arylamino groups, substituted and
52 unsubstituted dialkylamino groups, substituted and unsubstituted
53 diarylamino groups, substituted and unsubstituted (alkyl)(aryl)amino
54 groups, substituted and unsubstituted alkyl groups, substituted and
55 unsubstituted aryl groups, -C(=O)H, -C(=O)-alkyl groups, and
56 -C(=O)-aryl groups;
- 57 R¹⁰ is selected from the group consisting of substituted and
58 unsubstituted alkyl groups, substituted and unsubstituted aryl groups,
59 substituted and unsubstituted heterocyclyl groups, substituted and
60 unsubstituted heterocyclylalkyl groups, -C(=O)H, -C(=O)-alkyl
61 groups, -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl
62 groups, -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl)
63 groups, -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups,
64 -C(=O)N(alkyl)(aryl) groups, -NH₂, -NH(alkyl) groups, -NH(aryl)
65 groups, -N(alkyl)₂ groups, -N(alkyl)(aryl) groups, -N(aryl)₂ groups,
66 -C(=O)NH(heterocyclyl) groups, -C(=O)N(heterocyclyl)₂ groups,
67 -C(=O)N(alkyl)(heterocyclyl) groups, and
68 -C(=O)N(aryl)(heterocyclyl) groups;
- 69 R¹¹ and R¹⁹ may be the same or different and are independently
70 selected from the group consisting of substituted and unsubstituted
71 alkyl groups, and substituted and unsubstituted aryl groups;

72 R^{12} is selected from the group consisting of H, substituted and
73 unsubstituted alkyl groups, substituted and unsubstituted aryl groups,
74 and substituted and unsubstituted heterocyclyl groups;

75 R^{13} is selected from the group consisting of H, substituted and
76 unsubstituted alkyl groups, substituted and unsubstituted aryl groups,
77 substituted and unsubstituted heterocyclyl groups, -OH, alkoxy
78 groups, aryloxy groups, -NH₂, substituted and unsubstituted
79 heterocyclylalkyl groups, substituted and unsubstituted aminoalkyl
80 groups, substituted and unsubstituted alkylaminoalkyl groups,
81 substituted and unsubstituted dialkylaminoalkyl groups, substituted
82 and unsubstituted arylaminoalkyl groups, substituted and
83 unsubstituted diarylaminoalkyl groups, substituted and unsubstituted
84 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
85 alkylamino groups, substituted and unsubstituted arylamino groups,
86 substituted and unsubstituted dialkylamino groups, substituted and
87 unsubstituted diarylamino groups, substituted and unsubstituted
88 (alkyl)(aryl)amino groups, -C(=O)H, -C(=O)-alkyl groups,
89 -C(=O)-aryl groups, -C(=O)O-alkyl groups, -C(=O)O-aryl groups,
90 -C(=O)NH₂, -C(=O)NH(alkyl) groups, -C(=O)NH(aryl) groups,
91 -C(=O)N(alkyl)₂ groups, -C(=O)N(aryl)₂ groups,
92 -C(=O)N(alkyl)(aryl) groups, -C(=O)-heterocyclyl groups,
93 -C(=O)-O-heterocyclyl groups, -C(=O)NH(heterocyclyl) groups,
94 -C(=O)-N(heterocyclyl)₂ groups, -C(=O)-N(alkyl)(heterocyclyl)
95 groups, -C(=O)-N(aryl)(heterocyclyl) groups, substituted and
96 unsubstituted heterocyclylaminoalkyl groups, substituted and
97 unsubstituted hydroxyalkyl groups, substituted and unsubstituted
98 alkoxyalkyl groups, substituted and unsubstituted aryloxyalkyl
99 groups, and substituted and unsubstituted heterocycliloxyalkyl
100 groups;

-200-

- 101 R¹⁴ is selected from the group consisting of H, -OH, alkoxy groups,
102 aryloxy groups, -NH₂, -NH(alkyl) groups, -NH(aryl) groups,
103 -N(alkyl)₂ groups, -N(aryl)₂ groups, -N(alkyl)(aryl) groups,
104 substituted and unsubstituted alkyl groups, substituted and
105 unsubstituted aryl groups, -NH(heterocyclyl) groups,
106 -N(heterocyclyl)₂ groups, -N(alkyl)(heterocyclyl) groups, and
107 -N(aryl)(heterocyclyl) groups;
- 108 R¹² and R¹³ may join together to form a 5 to 7 membered saturated or
109 unsaturated, substituted or unsubstituted N-containing ring;
- 110 R¹⁵ is selected from the group consisting of substituted and
111 unsubstituted alkyl groups, substituted and unsubstituted aryl groups,
112 substituted and unsubstituted heterocyclyl groups, substituted and
113 unsubstituted heterocyclylalkyl groups, -C(=O)H, -C(=O)-alkyl
114 groups, -C(=O)-aryl groups, -C(=O)NH₂, -C(=O)NH(alkyl)
115 groups, -C(=O)NH(aryl) groups, -C(=O)N(alkyl)₂ groups,
116 -C(=O)N(aryl)₂ groups, -C(=O)N(alkyl)(aryl) groups, substituted
117 and unsubstituted aminoalkyl groups, substituted and unsubstituted
118 alkylaminoalkyl groups, substituted and unsubstituted
119 dialkylaminoalkyl groups, substituted and unsubstituted
120 arylaminoalkyl groups, substituted and unsubstituted
121 diarylaminoalkyl groups, substituted and unsubstituted
122 (alkyl)(aryl)aminoalkyl groups, substituted and unsubstituted
123 heterocyclylaminoalkyl groups, substituted and unsubstituted
124 diheterocyclylaminoalkyl groups, substituted and unsubstituted
125 (heterocyclyl)(alkyl)aminoalkyl groups, substituted and unsubstituted
126 (heterocyclyl)(aryl)aminoalkyl groups, substituted and unsubstituted
127 alkoxyalkyl groups, substituted and unsubstituted aryloxyalkyl
128 groups, substituted and unsubstituted hydroxyalkyl groups, and
129 substituted and unsubstituted heterocycloxyalkyl groups;

-201-

- 130 R^{16} is selected from the group consisting of H, substituted and
131 unsubstituted alkyl groups, substituted and unsubstituted aryl groups,
132 and substituted and unsubstituted heterocyclyl groups;
- 133 R^{17} is selected from the group consisting of H, substituted and
134 unsubstituted alkyl groups, substituted and unsubstituted aryl groups,
135 substituted and unsubstituted heterocyclyl groups, OH, substituted
136 and unsubstituted alkoxy groups, substituted and unsubstituted
137 aryloxy groups, $-NH_2$, $-C(=O)H$, $-C(=O)$ -alkyl groups, $-C(=O)$ -aryl
138 groups, $-C(=O)NH_2$, $-C(=O)NH$ (alkyl) groups, $-C(=O)NH$ (aryl)
139 groups, $-C(=O)N$ (alkyl)₂ groups, $-C(=O)N$ (aryl)₂ groups,
140 $-C(=O)N$ (alkyl)(aryl) groups, $-C(=O)O$ -alkyl groups,
141 $-C(=O)O$ -aryl groups, substituted and unsubstituted aminoalkyl
142 groups, substituted and unsubstituted alkylaminoalkyl groups,
143 substituted and unsubstituted dialkylaminoalkyl groups, substituted
144 and unsubstituted arylaminoalkyl groups, substituted and
145 unsubstituted diarylaminoalkyl groups, substituted and unsubstituted
146 (aryl)(alkyl)aminoalkyl groups, substituted and unsubstituted
147 heterocyclylalkyl groups, $-C(=O)$ -heterocyclyl groups,
148 $-C(=O)$ -Oheterocyclyl groups, $-C(=O)NH$ (heterocyclyl) groups,
149 $-C(=O)$ -N(heterocyclyl)₂ groups, $-C(=O)$ -N(alkyl)(heterocyclyl)
150 groups, $-C(=O)$ -N(aryl)(heterocyclyl) groups, substituted and
151 unsubstituted heterocyclylaminoalkyl groups, substituted and
152 unsubstituted hydroxyalkyl groups, substituted and unsubstituted
153 alkoxyalkyl groups, substituted and unsubstituted aryloxyalkyl
154 groups, and substituted and unsubstituted heterocycloxyalkyl
155 groups;
- 156 R^{16} and R^{17} may join together to form a 5 to 7 membered saturated or
157 unsaturated, substituted or unsubstituted N-containing ring; and

-202-

158 R¹⁸, R²⁰, and R²¹ may be the same or different and are independently
159 selected from the group consisting of H, -NH₂, -NH(alkyl) groups,
160 -NH(aryl) groups, -N(alkyl)₂ groups, -N(aryl)₂ groups,
161 -N(alkyl)(aryl) groups, substituted and unsubstituted alkyl groups,
162 substituted and unsubstituted aryl groups, -OH, substituted and
163 unsubstituted alkoxy groups, substituted and unsubstituted aryloxy
164 groups, substituted and unsubstituted heterocyclyl groups, -NHOH,
165 -N(alkyl)OH groups, -N(aryl)OH groups, -N(alkyl)O-alkyl groups,
166 -N(aryl)O-alkyl groups, -N(alkyl)O-aryl groups, and -N(aryl)O-aryl
167 groups.

1 30. The compound according to claim 29, wherein Y is selected
2 from the group consisting of H, -OH, -OR¹⁰ groups, and -NR¹²R¹³ groups.

1 31. The compound according to claim 29, at least two of X¹, X²,
2 X³, and X⁴ are C and the corresponding substituents R⁵, R⁶, R⁷, and R⁸ are
3 hydrogen, and at least one of X¹, X², X³, and X⁴ is N.

1 32. The compound according to claim 29, wherein R⁶ or R⁷ is an
2 alkyl group.

1 33. The compound according to claim 29, wherein R⁶ or R⁷ is an
2 -OR¹⁵ group and R¹⁵ is an alkyl, aryl, heterocyclyl, or heterocyclylalkyl group.

1 34. The compound according to claim 29, wherein R¹ is selected
2 from the group consisting of H, substituted and unsubstituted alkoxy groups,
3 substituted and unsubstituted heterocyclylalkoxy groups, substituted and
4 unsubstituted heterocyclyoxy groups, and substituted and unsubstituted heterocyclyl
5 groups.

1 35. The compound according to claim 29, wherein R² is selected
2 from the group consisting of H, F, Cl, -NO₂, substituted and unsubstituted
3 heterocyclyl groups, and substituted and unsubstituted heterocyclalkoxy groups.

1 36. A pharmaceutical formulation, comprising the compound
2 according to any of claims 1, 8, 15, 22, or 29 in combination with a
3 pharmaceutically acceptable carrier.

1 37. A method of treating a patient in need of an inhibitor of
2 vascular endothelial growth factor receptor tyrosine kinase, comprising
3 administering an effective amount of the pharmaceutical formulation according to
4 claim 36 to a patient in need thereof.

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/04 C07D405/04 C07D409/04 A61K31/4704

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, A	WO 01 28993 A (MERCK & CO. INC) 26 April 2001 (2001-04-26) cited in the application page 89 -page 106; claims 1-31 ---	1-37
A	WO 98 13350 A (ZENECA LIMITED) 2 April 1998 (1998-04-02) page 107 -page 122; claims 1-22 ---	1-37
A	WO 97 03069 A (GLAXO GROUP LIMITED) 30 January 1997 (1997-01-30) page 37 -page 43; claims 1-18 ---	1-37
A	EP 0 290 153 A (PFIZER LIMITED) 9 November 1988 (1988-11-09) the whole document ---	1-37
	--- -/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

B document member of the same patent family

Date of the actual completion of the international search

27 February 2002

Date of mailing of the international search report

06/03/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kyriakakou, G

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 414 088 A (WOLFGANG VON DER SAAL ET AL.) 9 May 1995 (1995-05-09) the whole document	1-37
A	IGOR V. UKRAINETZ ET AL.: "Effective Synthesis of 3-(Benzimidazol-2-yl)-4-Hydroxy-2-Oxo-1,2-Dihydroquinolines" TETRAHEDRON LETTERS., vol. 36, no. 42, 1995, pages 7747-7748, XP004027132 ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM., NL ISSN: 0040-4039 cited in the application the whole document	1-35
A	MARTIN P. MAGUIRE ET AL.: "A New Series of PDGF Receptor Tyrosine Kinase Inhibitors: *-Substituted quinoline Derivatives" JOURNAL OF MEDICINAL CHEMISTRY., vol. 37, no. 14, 1994, pages 2129-2137, XP002133783 AMERICAN CHEMICAL SOCIETY. WASHINGTON., US ISSN: 0022-2623 the whole document	1-37
A	LAURENT F. HENNEQUIN ET AL.: "Design and Structure-Activity Relationship of a new class of Potent VEGF Receptor Tyrosine Kinase Inhibitors" JOURNAL OF MEDICINAL CHEMISTRY., vol. 42, no. 26, 1999, pages 5369-5389, XP002164236 AMERICAN CHEMICAL SOCIETY. WASHINGTON., US ISSN: 0022-2623 the whole document	1-37
A	DAVID R STOVER ET AL.: "Recent advances in protein kinase inhibition: Current molecular scaffolds used for inhibitor synthesis" CURRENT OPINION IN DRUG DISCOVERY AND DEVELOPMENT, vol. 2, no. 4, 1999, pages 274-285, XP000926237 CURRENT DRUGS, LONDON, GB ISSN: 1367-6733 the whole document	1-37

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0128993	A	26-04-2001	AU 1091301 A WO 0128993 A2	30-04-2001 26-04-2001
WO 9813350	A	02-04-1998	AU 733551 B2 AU 4313797 A CN 1237963 A EP 0929526 A1 WO 9813350 A1 JP 2001500890 T NO 991423 A	17-05-2001 17-04-1998 08-12-1999 21-07-1999 02-04-1998 23-01-2001 11-05-1999
WO 9703069	A	30-01-1997	AU 6613996 A WO 9703069 A1 EP 0843671 A1 HR 960316 A1 JP 11508906 T ZA 9605935 A	10-02-1997 30-01-1997 27-05-1998 28-02-1998 03-08-1999 12-02-1998
EP 290153	A	09-11-1988	DK 214088 A EP 0290153 A1 FI 881825 A JP 63280078 A PT 87273 A US 4898872 A	07-12-1988 09-11-1988 22-10-1988 17-11-1988 01-05-1988 06-02-1990
US 5414088	A	09-05-1995	DE 3932953 A1 AU 6526190 A CA 2066690 A1 WO 9104974 A1 EP 0496753 A1 JP 5501249 T	11-04-1991 28-04-1991 04-04-1991 18-04-1991 05-08-1992 11-03-1993

THIS PAGE BLANK (USPTO)

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
21 March 2002 (21.03.2002)

PCT

(10) International Publication Number
WO 02/022598 A1(51) International Patent Classification⁷: C07D 401/04,
405/04, 409/04, A61K 31/4704

(21) International Application Number: PCT/US01/42131

(22) International Filing Date:
11 September 2001 (11.09.2001)

(25) Filing Language: English

(26) Publication Language: English


(30) Priority Data:
60/232,159 11 September 2000 (11.09.2000) US(71) Applicant (for all designated States except US): CHI-
RON CORPORATION [US/US]; 4560 Horton Street,
Emeryville, CA 94608-2916 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): RENHOWE, Paul
[US/US]; 262 Stetson Drive, Danville, CA 94506 (US).
PECCHI, Sabina [IT/US]; 5834 Merriewood Drive,
Oakland, CA 94611 (US). MACHAJEWSKI, Tim
[US/US]; 2514 Norwalk Court, Martinez, CA 94553 (US).
SHAFFER, Cynthia [US/US]; 4868 El Grande Place, El
Sobrante, CA 94803 (US). TAYLOR, Clarke [US/US];
612 Adams Street, Albany, CA 94706 (US). McCREA,
Bill [US/US]; 1040 Amato Drive, Berkeley, CA 94705
(US). McBRIDE, Chris [US/US]; 4504 Lawton Avenue,
Oakland, CA 94609 (US). JAZAN, Elisa [US/US]; 520
McLaughlin Avenue, Richmond, CA 94805 (US). WER-
NETTE-HAMMOND, Mary-Ellen [US/US]; 17893
Center Street, Castro Valley, CA 94546 (US). HARRIS,
Alex [US/US]; 14 MacPherson Place, Danville, CA 94526
(US).


[Continued on next page]

(54) Title: QUINOLINONE DERIVATIVES AS TYROSINE KINASE INHIBITORS



I

(57) Abstract: Organic compounds having the formulas I and II are provided where the variables have the values described herein. Pharmaceutical formulations include the organic compounds or pharmaceutically acceptable salts thereof and a pharmaceutically acceptable carrier and may be prepared by mixing the organic compounds or pharmaceutically acceptable salts of the organic compounds with a carrier and water. A method of treating a patient includes administering a pharmaceutical formulation according to the invention to a patient in need thereof.



II

WO 02/022598 A1



(74) **Agent:** FRIEDRICHSEN, Bernard; Foley & Lardner,
150 East Gilman Street, P.O. Box 1497, Madison, WI
53701-1497 (US).

IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

(81) **Designated States (national):** AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU,
ZA, ZW.

Published:

— with international search report

(48) **Date of publication of this corrected version:**

21 November 2002

(15) **Information about Correction:**

see PCT Gazette No. 47/2002 of 21 November 2002, Sec-
tion II

(84) **Designated States (regional):** ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)